STATISTICS

1. **Two or more group 0f subjects follow for 5 yrs to compare treatment of sideroblastic anemia. The most analysis of difference to be use is?**
    a) regression analysis
    b) chi square test
    c) ANOVA
    d) Association curve
2. **2group of patients followed for period of 10 yrs to study the Effect of cigarette smoking in lung cancer, type of study?**
    a) Cohort study
    b) Cross sectional study
    c) Case control study
    d) Experimental study
    e) Descriptive study
3. **Which of the following statements about sampling distribution of the sample mean is incorrect?**
    a) stander error of sample mean decreases as sample increase
    b) stander error of sample mean measures the difference between the sample mean and the population mean
    c) sample mean is unbiased estimator of the true (unknown) population mean
4. **Which of the following is correct?**
    a) a large sample always ensure that sample is represent of population
    b) in random samples the randomization ensure that we have precise and accurate estimate
    c) we do not need to randomize if our sample size is sufficiency enough
    d) if all other things are equal, we need a larger sample size
5. **The median is:**
    a) The most frequently occurring value
    b) The average value
    c) The middle value when data is ordered is ascending manner
    d) the difference between the maximum and minimum values in a data set

6. **Null Hypothesis is:**
    a) there is no significant association between the variables
    b) there is a significant association between the variables
    c) Subjective in nature
    d) The same as research hypothesis
7. **Type I error in testing hypothesis is:**
    a) Rejects null hypothesis when it is true
    b) Accepts null hypothesis when it is false
    c) The sample mean differs from the population mean
    d) The test is biased
8. **Standard deviation:**
    a) The Sum. Of Deviation from the mean
    b) The square root of variance
    c) can be calculated for both sample and population data
    d) The data set is perfectly distributed
9. **The confidence interval is:**
    a) A point estimate of the population parameter
    b) A range of values within which the population parameter is expected to fall
    c) A range of values within which the mean is expected to fall
    d) The total number of observations in a sample
10. **To estimate a population parameter based on sample statistics, this is called:**
    a) Simple Random Sampling
    b) Sample Distribution
    c) Point Estimation
    d) Inferential Statistics
11. **Probably distribution of variables including all values in the sample:**
    a) Sample statistic
    b) Simple random sampling
    c) Sample distribution
    d) Parameter
12. **P-value:**
    a) The probability that the null hypothesis is true
    b) The probability of rejecting the alternative hypothesis
    c) probability of observing data assuming null hypothesis is true
    d) probably the data results by chance