EXAM 3

RECALL

MARCH 2022

casein is the milk protein

1. Renin converts casein to paracasein in presence of

a. Ca

b. Mg

c. Na

d. K

Ca > Na

2. Hypokalemia diagnose clinically mainly by:

a. Cardiac arrhythmia

b. Hyperreflexes

c. Diarrhea

d. Antagonised digoxin effect

3. Which of the following test the most sensitive for diagnosis of iron deficiency anemia

a. serum iron

b. Serum transferrin

c. serum ferritin most sensitive is the storage form

d. serum erythropoietin

4. Which of the following represent the correct order of events of menstrual cycle ?

a. FSH rise , LH surge, ovulation , Luteal phase, menstruation.

b. Luteal phase, menstruation, FSH rise, ovulation ,follicular phase

c. follicular phase , Luteal phase ,FSH rise, ovulation, menstruation

d. Menstruation ,FSH rise ,LH surge , ovulation, luteal phase

1st day of cycle is 1st day of bleeding
FSH starts follicular growth (arises)
Estrogen is responsible for maturation and growth of the follicle
LH surge then ovulation
Corpus luteum form > Progesterone
I fertilization occurs > hCG
If not > shedding of endometrium

5. Which of the following represents the typical order of physical and physiologic changes of puberty in females?

a. Menarche ,thelarche, andrenarche, peak growth velocity

b. Thelarche, andrenarche , peak growth velocity, Menarche

c. peak growth velocity, adrenarche , Menarche, thelarche

d. adrenarche , menarche, thelarche, peak growth velocity

9 years 10 years for hair growth 11 - 13 years

6. What is the correct order to the phases of the sexual response cycle?

a. Sexual attraction or arousal, excitement phase, plateau phase, orgasm, resolution phase

> in female > in male

b. Sexual attraction or arousal, excitement phase, orgasm, plateau phase, resolution phase

c. Sexual attraction or arousal, excitement phase, orgasm, resolution phase , plateau phase

d. plateau phase, excitement phase, orgasm, resolution phase, Sexual attraction or arousa

7. During the follicular phase, the dominant follicle arise in response to which of the following?

a) Matrix –degrading proteases

if maturation: estrogen

b) HCG

c) FSH

d) Androgens

8. Which of the following statement correctly describes one of the three physiologic differences between male and female orgasms?

a. The male is capable of rapidly successive orgasms

b. The female is capable of rapidly successive orgasms

c. The duration of the female orgasm is shorter than that of the male orgasm Longer

d. The female experiences emission and ejaculation ,but to lesser extent than the male

9. The posterior pituitary releases two hormones ,arriving directly from the hypothalamus:

a. Insulin and thyroxine

b. Oxytocin and insulin

c. Oxytocin and ADH

d. ACTH

In hypothalamus:
Paraventricular nucleus: Oxytocin
Supraoptic nucleus: ADH
Suprachiasmatic nucleus: ACTH

10. All of the following are true about human HCG hormone except:

a. In early pregnancy , the level are expected to double every 2_3 days

b. it is structurally related to oxytocin , and bind to receptors to support the corpus luteum

hCG is related to: FSH, LH, TSH

c. It is initially produced by the trophoblast cells of the embryo

d. During the first trimester, it stimulates the fetal gonads to make steroid hormones responsible for sexual differentiation

11. There are at least four mechanisms involved in transporting critical molecules in matched with the appropriate mechanism?

a. Respiratory gases are transported by active transport passive diffusion

Ca and Glc by facilitated diffusion
AA by active diffusion

b. Immunoglobulin G (IgG) bind s a membrane receptor and enters via endocytosis

c. Calcium cross the membrane by diffusion down a concentration gradient

d. Glucose is bind to membrane receptors and enters the placental tissue by endocytosis

12.S.D is a 25 years old woman who has been attempting to conceive for 1 year. She presented to an infertility doctor , as her efforts have been unsuccessful . she reported that she has been timing intercourse based on the change in her vaginal discharge , she noted that midcycle , her discharge becomes thin and watery, high level of which hormone account for this change along with softening of the cervical stroma? = thin cervix, used in HSG

a. Estrogen + decrease osteoclast (bone resorption) vasodilation + Breast ductal proliferation + Decrease cholesterol synthesis

b. Progesterone

c. Prolactin

d. Antidiuretic hormone

13. Which of the following test the most sensitive for diagnosis of iron deficiency anemia in pregnancy

a. serum iron

b. Serum transferrin

c. serum ferritin

d. Total iron binding capacity

14. Failure to correct hypokalemia is concomitant caused by:

a. Decrease in magnesium count

b. Decrease in sodium count

c. Decrease in chloride count

15. Bacterial cells are at their metabolic peak during

a. lag phase

b. log

c. stationary

d. decline

Phases of bacterial growth:
- Lag: inactive, no growth
- Log: growth, metabolic active
- Stationary: stable no increase number
- Decline: loss number

Type of bacterial reproduction
Binary fission

Viruses have complicated growth with the host DNA

Fungal growth by budding

16. Viral DNA is resistant to DNA Of the host cell because it contains

a. 5-HMC 5-Hydroxymethylcytosine

b. 5-HMA

c. 5-CHM

d. 5-MHC

17. Mycoplasma are bacterial cells that

has no cell wall, MCC of bacterial Atypical pneumonia Tx by Tetracycline and Erythromycin Resistant also to vancomycin

a. fail to reproduce on artificial media

b. have a rigid cell wall

c. are resistant to penicillin

d. stain well with gram s stain

18. which of the following was non-neural vaccine for rabies

a. HEPV

b. Card vaccine

c. BPL (beta-propiolactone)

d. Simple

19. Example for live vaccine is

Vaccines:
1- Live attenuated: BCG. Sabin oral polio, MMR, Yellow fever
Risk in immunodeficiency, pregnancy, elderly
2- Toxoid (subcellular): Polio IM Salk, DTaP, H.inf B, HBV, PCV
3- Recombinant reassortment in viruses: HPV, Rota-rex

a. Rubella and BCG

b. Polio and TAB

c. Diphtheria and Tetanus

d. Hepatitis A and Rabies.

20. Lyme disease is caused by

caused by Borrelia Burgdorferi (Atypical bacteria), transmitted by tick bites

a. bacteria

b. Fungi

c. Spirochaete

d. Virus

21. The mode of reproduction which occurs in mycoplasma is

a. Budding

b. Bursting

c. Binary fission

d. Binary fusion

22. Streptococcus fecalis , the conjugation take place at

a. pili

b. Cell membrane

c. cell wall

d. flagella.

23. Endotoxin produced by gram negative bacteria is present in

a. Peptidoglycan

b. Lippolysacharide

c. Theichoic acid stained by gram stain/ not available in gram-negative

d. Inner membrane

24. which one of the following is about herpes viruses

a. Icosahedral ,with envelope,ds DNA

b. Polyhedral with envelope ,dsDNA

c. RNA ,helical with envelope

d. Ds DNA , brick shape

DNA viruses:
1- Non-enveloped: adeno v, HPV, Parvo v.
2- Enveloped: HSV + HBV + Pox v.

Icosahedral viruses: by Capsule shape
HPV, HBV, HSV, Rhino virus

25. Cold like symptoms are caused by which bacteria

a. Pseudomonas

b. E.coli

c. Haemphilus influenza chocolate agar for culture

d. Haemphilus streptococcus.

* Bacterial culture media:
Neisseria: Chocolate or Thayer–Martin
Strep and staph: Blood agar
Enterobacteria: MacConkey
Legionella: Charcoal
TB: Löwenstein–Jensen (LJ)
Diphtheria: Löffler
Bordetella: Regan-Lowe
Vibrio: Thiosulfate-Citrate-Bile Salts-Sucrose (TCBS)

26. AIDS disease is caused by a virus which belongs to

a. Retro virus = Lentivirus

b. Rhabdo virus

c. Hepatitis virus

d. Adeno virus

27. Incubation period for infective hepatitis disease = HAV

a. 45-80 days

b. 15-35 days

c. 35-50 days

d. 5-15 days

Incubation period:
HBV: 2 - 5 months
HCV: 2 weeks to 6 months
HEV: 15 to 70 days

28. Vaccines prepared from toxins and chemicals are

a. Cellular vaccines

b. Sub-cellular vaccines

c. Attenuated vaccines

d. Heterologous vaccines

29. Most common pneumonia in HIV patients due to = AIDS patients

a. Pneumocystitis carinii

b. Cryptospordidum

c. S. pneumoiae MC Bacterial pneumonia MC Viral pneumonia is CMV

d. Toxoplasma

30. Pregnant lady in her first trimester developed skin rash and fever for few days .. after delivery her baby presented with small head and skin rash , what is your diagnosis? Blue berry muffin rash is CMV

a. T. pallidum

b. Measles

c. Rubella

d. CMV

31. Genetic material of influenza virus is?

a. Single helix DNA only Parvo virus

b. Single helix RNA

c. Double strand RNA only Rota virus

d. Single strand RNA

32. Mutation if which the following corresponding gene can result in resistant to HIV infection:

a. T cell receptor

b. Chemokine receptor CC-R5 receptor

c. Fc receptor

d. CD28 receptor

33. Proteins are soluble in

a. anhydrous acetone.

b. aqueous alcohol = Universal solvent

c. anhydrous alcohol

d. benzene

34. The hydrogen bond between peptide linkages of a protein molecules are interfered by

a. guanidine break peptide bonds

b. uric acid

c. oxalic acid

d. salicylic acid

35. A lipoprotein inversely related to the incidence of coronary artherosclerosis is

b) VLDL TAG

c) LDL Directly: move cholesterol from liver to blood

d) IDL TAG

e) HDL Move cholesterol from blood to liver

35. The amino acid with a nonpolar side chain is = no charge

a. Serine Glutamine + Arginine is polar

b. Valine + Alanine, leucine, proline, isoleucine, tryptophan, phenylalanine

c. Asparagine

d. Threonine

36. In case of repeated contraction , muscle cells regenerate NAD By converting pyruvate to

a. Glycogen

b. Lactate in anaerobic glycolysis by lactate dehydrogenase.
Consume ATP and regenerate NAD+

c. Starch

d. Cellulose

37. what is the total gain of energy during anaerobic respiration

a. Two ATP Aerobic: 6 or 8 ATP
TCA: 12 ATP

b. one ATP

c. Four ATP

d. Three ATP

38. Conversion of alanine to carbohydrate is termed

a. Glycogenesis

b. Gluconeogensis alanine (AAs) + lactate + glycerol + propionate

c. Glycogenolysis

d. Photosynthesis

39. The following is an enzyme required for glycolysis

a. Pyruvate kinase

b. Pyruvate carboxylase

c. Gluose -6-phosphatase b + c + d is for Gluconeogenesis

d. Glycerokinase

40. Aketogenic amino acid is

B. valine

a. Cysteine

b. Leucine

c. Threonine

41. The synthesis of glucose from pyruvate by gluconeogenesis

a. Requires the participation of biotin

b. Occurs exclusively in the cytosol

c. Is inhibited by elevated level of insulin

d. Requires oxidation /reduction of FAD.

42. glycogen is converted to glucose-1-phosphate by Break down glycogen by

a. UDPG transferase

b. Branching enzyme

c. Phosphorylase

d. Phosphatas control by insulin to increase synthesis unlike kinase by glucagon

43. Sucrose is composed of

a. Glucose

b. Glucose and Glycogen

c. Two molecules of glucose

d. Glycogen and fructose

44. The intermediate in hexose monophosphate shunt is

a. D-Ribulose

b. D-Arabinose

c. D-Xylose

d. D-lyxose

Pentose phosphate pathway
It is regulated by glucose 6-phosphate dehydrogenase
It gives mainly:
- Ribulose 5-phosphate: which used in DNA synthesis
- NADPH: which is used in the synthesis of fatty acids and eliminates free radicals

45. Which of the following is a mitochondrial enzyme that mediates all steps in the conversion of cholesterol to pregnenolone and rate limiting step for the entire steroid synthetic pathway

a. P450P10

b. P450scc SCC= Side chain cleavage

c. p450mwc

d. p450ACC

46. Two condition in which gluconeogenesis is increased are

a. Diabetes mellitus and atherosclerosis

b. Fed condition and thyrotoxicosis Thyrotoxicosis is antagonizing Insulin

c. Diabetes mellitus and starvation

d. alcohol intake and cigarette smoking

47. Oxidative decarboxylation of pyruvate requires

a. NADP+

C. Cytichromes

b. Pyridoxal phosphate

c. COASH

48. Pyruvate dehydrogenase complex and ketoglutarate dehydrogenase complex require the following for their oxidative decarboxylation:

a. COASH and Lipoic acid Co-factors are important

b. NAD+ and FAD

c. COASH and TPP

d. COASH, TPP, NAD, FAD, Lipoate. TPP= Thiamine pyrophosphate

49. High energy molecule needed to convert oxaloacetete to phosphoenoyl pyruvate

Pyruvate to oxaloacetate = need ATP

a. GTP

b. ITP

c. ITP and GTP

d. None of the above

50. Monosaccharide are

a. glucose and fructose + Galactose

b. aldose and hexose

c. maltose

51. Ehlers –Danlos syndrome characterized by hypermobile joints and skin is due to

a. Abnormality in gene for pro collagen

b. Deficiency of lysyl oxidase

c. Deficiency of prolyl hydroxylase

d. Deficiency of lysyl hydroxylase

52. If you examine primordial germ cells in a prepuberal male under the microscope, in which stage of cell cycle will you find them ?

e. Prophase of meiosis 1 if female before puberty
If female after puberty = Meiosis 2

f. Anaphase

g. Telophase

h. Interphase G1, S, G 2 = No M

- Spermatogenesis is continuous after puberty, every 2 months
- Oogenesis: Begins in the fetal period and is a discontinuous process, start mitosis in week 5 till 5 months = 7 million (max) > apoptosis till menopause
- Meiosis 1 starts at labor until puberty (ovulation)
- Thin stop in Meiosis 2 until fertilization

53. Which of the following is a true statement in an otherwise normal gestation ?

a. in the presence of SRY , a female phenotype will develop Male will develop

b. in the absence of SRY , a female phenotype will develop

c. SRY is located on the X chromosome

d. Only one X chromosome is necessary for normal ovarian development.

54. 20 years old girl attends the gynaecology clinic with her mother , she presents with primary amenorrhea, on examination she is tall with BMI 19Kg/m2 Blind end vagina . there is no axillary or pubic hair. What is the most likely diagnosis ?

- **Complete androgen insensitivity syndrome**
- Klinefelter syndrome
- Mullerain agenesis
- Turner syndrome

55. M.T. is a 27 year old white woman who has been attempting pregnancy for 15 months . she is monitoring her ovulation with LH testing . her test today indicated that she will ovulate tomorrow . in what phase will the ovulated germ cell be arrested?

a. Prophase of first mitotic division

If female after puberty = Meiosis 2

b. Anaphase of first meitotic division

c. Metaphase of second meitotic division complete by fertilization > lead to ovum + 3 polar bodies

d. Zygotic phase

56. pregnant lady carrier for gene responsible for Duchenne muscular dystrophy her children will

a. 50% of sons and 50 % of daughters will affected

b. 50% of sons will be affected

c. 100% of sons will be affected

d. 100% of daughters will be affected.

X linked recessive if mother carrier
Females: 50% carriers + 50% healthy
Males: 50% diseased + 50% healthy

X linked recessive if mother diseased
Females: 100% carriers
Males: 100% diseased

X linked recessive if father diseased
Females: 100% carriers
Males: 100% healthy

57. Which type of antibodies will associate in blood cell coagulation

a. IgE

b. IgM

c. IgA

d. IgG

58. P53

a. Mutated in tumor cell

b. Oncogen

c. Used in cell replication

d. DNA repair

59. Arterial infraction is least likely to be occur in:

a. Intestine

b. Spleen

c. Liver

d. Kidney

60. Creatine phosphate elevated level marker for + Myoglobin + Lactate dehydrogenase

a. Muscle disease

b. Cardiac disease

c. Kidney disease

d. Liver disease

61. All of the following are true about osteoporosis except

a. Can be primary or secondary

b. Increased osteoblast activity affect osteoclast

c. occur in vitamin D deficiency

d. occur in vitamin C deficiency

62. Vaso occlusive crisis in patient with sickle cell anemia is

a. Hand and foot syndrome RBC due to dehydration or hypoxia or acidosis = sickling and occlusion lead to ischemia and severe pain

b. Auto splenectomy

63. Drug induce asthma by..

a. Type I hypersensitivity reaction = Immediate

b. Induce immune reaction

c. act as hapten

64. Excision of small subcutaneous nodule after appendectomy by 28 days ,microscopic biopsy shows macrophages , collagen, lymphocytes, which retractile material are seen in nodule and giant cells and surgical complication

a. Chronic inflammation

b. Suture granuloma foreign body

c. Abscess formation

65. The substance synthesized by fibroblast and connect the cellular intergrin to exteracellular collagen and to basement membrane

a. Fiberonectin

b. Procollagen

c. Dermatan sulphate

d. Hyalouronic acid

66. Calf muscle affected after long period of immbolization by :

a. Hypertropy

b. Atrophy

c. Dystrophy

67. Esiophilic staining with loss of nuclei and striation in myocardial tissue ?

a. Coronary artery thrombosis

b. Viral infection

c. Chest trauma

68. After along period of corticosteroid therapy , wound healing is delayed due to :

a. Collagen damage

b. Fibroblast elaboration

69. 8 years old female bend thumb to touch her forearm, skin pull out about 8 cm , she has difficulty in healing of skin cuts. The underlying disease is caused by defect in:

a. Collagen

Healing or connective tissue disease = Collagen defect

b. Factor VIII

c. Antitrypsin

d. Retinoblastoma protein.

70. pyogenic bacterial infection and ischemic brain infarcts most characteristically produce which one of the listed types of necrosis

a. Caseous necrosis

b. Coagulation

c. Fat necrosis

d. liquefactive necrosis

71. Anasarca occur due to

a. Complicated pregnancy induce hypertension

b. surgical removal of lymph node

c. Nephrotic syndrome

Causes of Generalized edema
- CHF
- Renal failure
- Nephrotic syndrome
- Malnutrition

72. Generalized edema in congestive heart failure is due to

a. Increased hydrostatic pressure

b. Decreased oncotic pressure

c. Lymphatic obstruction

d. DVT

73. Metastatic calcification :

a. In TB

b. Vitamin D intoxication

c. Amylodosis

d. Acute haemorrhgic pancretitis

74. ciprofloxacin acts by inhibiting

b) cell wall synthesis

c) RNA synthesis

d) folate synthesis

e) DNA gyrase

75. Drug of choice for dermal , oral ,vaginal candidiasis is

a. Griseofulvin

b. Amphotericin B MoA: DNA disruption, toxic to humans and lead to renal and liver failure (eukaryotic)

c. Gentian violet

d. Nystatin

76. B- Lactum ring is present in

a. Erythromycin

b. Penicillin

c. Tetracyclins

d. Chromphenical

Beta-lactamase bacteria resistance
Inhibitor by Clavulanic acid given in combination with Amoxicillin = Amoclan

- Inhibit cell wall: Cephalosporins + Vancomycin + Bacitracin
- 30S: Aminoglycosides, Gentamicin, Tetracyclines
- 50S: Erythromycin, Clindamycin, Chloramphenicol

Quinolones: DNA gyrase and Topoisomerase
Rifampin: DNA-dependent RNA polymerase
Metronidazole: Disrupts DNA
Sulfonamides: Dihydropteroate synthetase
Trimethoprim: Dihydrofolate reductase

77. Erythromycin is obtained from

a. s. griseus

b. s. rimosus

c. s. scabies

d. s. erythraeus.

78. Mode of action of quinolone antibiotics on growing bacteria was thought to be

a. Inhibition of lactamase

b. Prevention of cross linking of glycine

c. Inhibition of DNA gyrase

d. Inhibition of reverse transcriptase

79. Drug of choice for methicillin resistant staph .Aureus is

a. Ampicillin

b. Erythromycin

c. Neomycin

d. Vancomycin

80. The anti tumor antibiotic act by inhibiting

a. Cell wall synthesis

b. RNA synthesis

c. Cell membrane synthesis

d. The DNA structure and function

81. Which of the following does not inhibit cell wall synthesis ?

a. Penicillin

b. Carbenicillin

c. Amikacin 30S protein

d. Vancomycin

82. Which of the following is penicillinase resisting acid labile penicillin ?

a. Amoxycillin

b. Cloxacillin

c. Carbenicillin

d. Methicillin

83. Muscle of urogenital diaphragm include

a. Sphincter urethrae

b. Sphincter vesicae

c. Levator ani

d. Ischiocavernosus

84. Cancer from pelvic viscera can deposits cancerous cells in the vertebrae this communication occur between the internal iliac veins and the vertebral veins through the:

a. Lateral sacral vein

b. Lumbar vein

c. Prostatic plexus

d. Umbilical veins

85. lymph node of lower part of uterine body is + cervix

a. Para aortic lymph node

b. Internal iliac lymph node

c. External iliac lymph node

d. Deep group

86. What is the glistening membrane that covers the fetal surface of the placenta?

a. Chorion

b. Decidua mother direction

c. Amnion

d. Cotyledon

87. 19 years old college student , majoring in biology ,just deliverd her first baby, she is very interested in studying early pregnancy and asks about the origin of the placental tissue .which of the following develop into the placenta?

a. Trophectoderm

b. Inner cell mass of the blastocyst

c. Decidua

d. Myoblast cells.

Decidua basalis end in menopause
- D & C: D. functionalis only
- Ablation: D. basalis and functionalis

88. Inguinal canal have four wall which of the following from the posterior wall:

a. Internal oblique

b. External oblique

c. Transvers abdominus

d. Transversalis fascia

89. Iliac crest at level of

a. L3

b. L4

a. L5 iliac tubercle

b. S1

90. The dermatomes of umbilical region ?

a. T7 xiphisternum

b. T10

c. T12 symphysis pubis

L2: thigh
L3: Medial knee + leg
L4: Medial foot
L5: Dorsum of foot
S1: Lateral foot + Posterior leg and knee
S2: Posterior thigh
S3: Perianal area

d. L1 Gentelia

91. The superficial inguinal ring from

a. Transverses abdominus

b. Aponeurosis external oblique — Deep ring in transversalis fascia

c. Aponeurosis internal oblique

d. conjoint tendon

92. Main muscle responsible for abduction of hip joint

a. Iliopoas muscle

b. Gluteus medius — support the pelvis in gait. Defect lead to Trendelenburg gait
N by superior gluteal N

c. Gluteus minimus

93. Pudendal nerve before enter the pudendal canal lie on which of the following structure

a. Piriformis

b. Obturator internus muscle and fascia

c. Sacrospinatous ligament

d. Sacrotuberous ligament

94. Which of the following operations involve in cutting rectus sheath anteriorly and posteriorly

a. paramedian

b. Subcostal

c. Transverse suprapubic

d. Vertical incision

Abdominal wall muscles
* Upper 1/3: only External oblique aponeurosis in front
* Middle 1/3: Transversalis fascia posteriorly and Internal oblique aponeurosis bisect half ant + post
* Lower 1/3: All Ms anteriorly. Only Transversalis fascia posteriorly

95. Medial part of the knee joint supply by

a. Femoral nerve anterior part — lateral cutaneous nerve of thigh: Laterally

b. Sciatic nerve posterior part

c. tibial nerve

d. Obturator nerve

96. Which of the following nerve supply the anterior 2/3 of labia majora and clitoris

a. Ilioinguinal nerve — pass in inguinal canal

b. Genitofemoral nerve cremasteric reflex

c. perineal branch of pudendal nerve

d. lateral sacral nerve

97. which of the following structure lies medial to femoral artery within femoral canal

a. femoral nerve outside the sheath

b. femoral vein

c. femoral sheath contents = femoral A + V + LN

d. lymph nodes

sheath

From medial to lateral
Canal, V, A, N

Femoral triangle boundaries:
Inguinal ligament superiorly
adductor longus muscle medially
sartorius muscle laterally.

98. Vagina covering by peritoneum in

a. anterior wall no peritoneum

b. posterior fornix

c. anterolateral wall

d. lateral fornix

99. Which of the following is non glandular organ

a. Uterus

b. Vagina stratified squamous epithelium

c. Fallopian tube

d. Cervix

100. Primary support of uterus by:

a. Cardinal ligament

b. Round ligament

c. Pubocervial ligament.

101. 35 years old grandmultipara has had a major postpartum haemorrhage (PPH) following normal delivery, mechanical and pharmacological measure have failed to control the bleeding, examination has confirmed that there are no retained placental tissue in uterine cavity and absence of trauma to genital tract, what is the most appropriate first line surgical management?

a. Balloon Tamponade

b. ligation of internal iliac artery 2nd option

c. postpartum hysterectomy

d. lynch or modified compression suture. if CS

102. D.E.is a 19 years old college student who presented to the ER for vaginal spotting. She reported that she had a positive home pregnancy test 2 days ago, her last menstrual period was 5 weeks ago. The ER physician ordered a quantitative human chorionic gonadotropin (HCG) test that was reported as 213 mIU/ml . what is the source of this hormone?

a. Fetal pituitary gland

b. Developing trophoblast

c. Ovarian corpus luteum

increase in HCG every 48 hours
increase > 60% is normal
increase < 60% in ectopic pregnancy
decrease = non-viable fetus
Doubling = Twins or molar

103. Most common causes of secondary post partum hemorrhage occur due to

a. Infection or endometritis

b. uterine atonia

c. uterine rupture

d. Endometriosis ✗

104. primary postpartum hemorrhage occur with in

a. 12 hours

b. 24 hours

c. 48 hours

d. 72 hours

105. You are attending the labor of patient who has prolonged 2nd stage of labor .you note the fetal head start to retract after being tightly applied to the vulva , what is the next most appropriate management step?

a. Apply fundal pressure

b. C-section

c. All fours manoeuvre

d. MC-roberts manoeuvre

= Turtle sign of shoulder dystocia
Mx by HELPER
1- Help
2- Episiotomy
3- Leg elevation (Mc Robert)
4- Pubic pressure
5- Enter maneuver
6- Role on 4 and repeat enter m
7- Methods of last resort

106. prgnant lady has hyperthyroidism on carbimazole developed sore throat investigation should be done

a. FBC cause Agranulocytosis

b. TFT confirm control then do CBC

c. Throat swab

d. CRP

107. 24 year old lady delivered by C/S Section in day 7 she developed SOB on climb stairs , what is causing her condition

a. PE

b. Amniotic fluid embolism

c. Hypertrophic cardiomyopathy SOB on exertion is a cardiac cause

d. Congestive heart failure

108. 48 year old lady presented with heavy menstrual bleeding and intermenstrual bleeding her HB is 112% and US showed no endometrial abnormalitiy next step

a. coagulation screening if HMB since menarche or with other source of bleeding

b. Diagnostic hysteroscopy

c. endometrial sampling if menopause go for hysteroscopy

d. pelvic MRI

109. 31 year old woman presents in preterm labour at 34 weeks of gestation , her labour progresses quickly and she delivers baby boy. Both mother and baby appear to be in good health , the woman requested delayed cord clamping. What time frame would be recommended for delayed cord clamping in this situation?

a. Clamp cord as soon as possible At term clamp the cord as soon as possible

b. 30 second - 3 minute

c. 10-30 second

d. 4-6 minutes

110. 19 years old female presented with amenorrhea for 5 week pregnancy test was positive and HCG level 7000 , uterus empty on U/S how to treat her

a. Conservative Typical presentation of ectopic pregnancy

b. Medical ttt with methotrexate

c. Mesoprestol oral

d. Surgery

111. Mid wife examine pregnant lady in 21 week of her gestation and she found her has fused labium minora , what is the type of FGM in this lady?

a. Type 1 cutting the clitoris only

b. Type 2 cutting the clitoris + labia minora with no suture

c. Type 3 cutting the clitoris + labia minora + majora with suture and vagina narrowing

d. Type 4 cauterization and piercing (harmful)

112. Study of VBAC in woman had previous vaginal delivery after one C/S section , type of study is?

a. Cohart study follow up study

b. cross sectional

c. Survey

d. Case control

113. Pregnant lady in her 38 week of gestation ,presented with labour pain and on PV there is palpable nasal bridge and palpebral fissure , what is diameter of this presentation?

a. Submentobregmatic = Face = 9.5 cm

b. Mentovertical = Brow presentation = 13 cm

c. Suboccipitiobregmatic = Vertex = 9.5 cm

d. Occipitofrontal = Deflexion vertex = 11.5 cm

114. Pregnant lady in 28 week of her gestation ,high vaginal swab for B. streptoccoi is positive .what you will do?

a. Give her course of oral AB

b. Intrapartum injectable AB

c. If not treated her baby will get severe neonatal sepsis

d. Reassurance

115. 32 year old primagravida in her 36 week of gestation presented with generalized tonic clonic convulsion , her BP 150/100 , urine contain ++ protein. What is the initial step in management?

a. Diazepam

b. Mg sulphate

c. Hydralizine

d. Nifedipie