EXAM 5

RECALL

MARCH 2023

1- amount of oxytocin used in cs

a-5IU

b-10IU in NVD + risk of PPH in CS

c-15IU

c-40IU

2- massive bleeding is or = 150ml/ min
or = more than 50% in less than 3 hours

a-loss 1 volume in 24 h

b-loss 1 volume in 2 h

c-loss 50% in 24 h

d- loss 50% in 2 h

Tx my Massive blood transfusion protocol:
Replacement of >1 blood volume in 24 hrs
or >50% of blood volume in 4 h
or > 5 units in 4 hours
Ratio: 1 packed RBC + 1 FFP + 1 Platelets
* Need monitoring ..... Risks:
- volume overload + e disturbances
- over-transfusion (monitor Hb)
- hypothermia (monitor temp)
- hyperkalemia
- disease transmission

3- amount of fresh frozen plasma used in massive bleeding

a-2 unit

b-4unit

c-6unit

d-10_15 ml True is 15-20 ml / kg

4- in which period uplistrail contraceptive is effective

a-24 h MoA: selective progesterone receptor modulator

b-72h

c-120h 5 days

d-48 h

5- side effect of warfarin in pregnancy :

a-nasal hypoplasia

b-renal agenesis

c- intracranial hemorrhage

7- about bacterial vaginosis

Answer diagnosed by hay\ison test used in classification of vaginal normal flora

8- after how long we consider 3rd stage of labor is abnormal

- 2nd stage
Prima: 2 hours
Multipra: 1 hour

- 1st stage latent / active
Prima: 1 cm in 2 h /
Multipra: 1 cm in 1 h /

Monitor every 4 hours
Assessment and reass
ARM
CS

a-20 min

b-30 min

c- 60 min

9- vaginal tear include rectal mucosa is

a- 1st degree Skin and mucosa

b- 2nd degree Skin and mucosa and superficial transverse perineal muscle

c- 3rd degree 3A <50% EAS, 3B >50% EAS, 3C EAS + IAS

d- 4th degree reach rectal mucosa

10- in suspected PPROM in speculum examination if no liquor appear next step is

a-send her home

b-repeat test after 4h c-pad test

d- PBL test

11- prim gravide in 40 week with spontaneous rupture membrane and 2 contraction per 10 min your action

a- give oxytocin

b- cs c-

d-

12- long diameter presentation is

Answer mentovertical

13- prim gravida with cervix 2 cm and intact membrane your action is

a. a-cs
b. b-check after 4 h
b. give oxytocin

14- pregnant lady with twin and sudden rupture of membrane your action

a. a-cs
b. vaginal examination to check liquor
c. vaginal exam to check cord prolapse 15-about using partogram
a. a-who suggest b-

16-amount of body iron in hemoglobin

a-50%

b-60%

c-75%

d-90%

18- female with primary ammenorrhoea and absent uturs and vagina cause is

a-gonadal agenesis b-mullurian agenesis c-pcos Ovarian cause: anovulation

Turner syndrome MRKH Syndrome

19- female with 28 day cycle omit her contraceptive in day 17 and 18 now she is on day 20 she worry about being pregnant your advice

a-reassuring that no thing will happen b-start new vail of contraceptive

c-take the 2 missing pill as fast as possible and continue in her contraceptive d-it is too late

20 -multipara under went previous cs then has successful VBAC after to years now pregnant after 9 years from the last pregnancy with fetus with wt 3500g fetus position…….factor that suggest she will now has successful vaginal delivery is

a-previous VBAC b-fetal size

c-period between pregnancy d- fetal position

1st = Engagement
2nd = Descent
3rd = Flexion
4th = Internal rotation
5th = Extension
6th = External rotation
7th = Delivery of anterior shoulder

21- 7 stage of normal labor the 6th stage is External rotation

22- contraindication to use instrumental delivery 23-In shoulder dystocia your action:

a-MC Robert and fundal pushing MC Robert + Supra pubic pressure

b-MC Robert and axial traction c-

1-which of the following was non-neural vaccine for rabies

a. HEPV

b. Card vaccine

c. BPL (beta-propiolactone)

d. Simple

22- Contraindication to use instrumental delivery
- Not engaged
- Not fully dilated
- Less than 34 week/ SGA < 1 kg/ IUGR
- Bleeding disorders
- Unknown position or transverse
- Large head = hydrocephalus
- HIV patients

2. Which type of antibodies will associate in blood cell coagulation

a. IgE

b.IgM

c.IgA

d.IgG

3- ciprofloxacin acts by inhibiting

b)cell wall synthesis

c)RNA synthesis

d)folate synthesis

e)DNA gyrase

4- Cancer from pelvic viscera can deposits cancerous cells in the vertebrae this communication occur between the internal iliac veins and the vertebral veins through the:

a. Lateral sacral vein

b. Lumbar vein

c. Prosatatic plexus

d. Umbilical veins

5- Which of the following nerve supply the anterior 2/3 of labia majora and clitoris

a. Ilioinguinal nerve Anterior 1/3 only

b. Genitofemoral nerve

c. perineal branch of pudendal nerve

d. lateral sacral nerve

6- Infectious mononucleosis, a viral disorder that can be debilitating, is characterized by which of the following statements?

A. It is most prevalent in children less than 14 years old

B. It is caused by a rhabdovirus

C. The causative pathogen is an Epstein-Barr virus

D. Ribavirin is the treatment of choice

7- A Lipoprotein lipase:

a. Hydrolyses cholesteryl ester

b. Is present in the cytosol of adipocytes

c. Is active during starvation

d. Hydrolyses TAG in chylomicrons and VLDL

8- Free redcal cause cell death by

A. denatured of protein contain sh group

B. apptosis

9- year old female known case of ovarian CA presented with Medial thigh Parasethia the nerve involved by CA is

a. obturator N

b. medial cutaneous N of the thigh

c. intermediate cutaneous N of the thigh

d. femoral trunk

e. lumbosacral trunk

10- about left ureter

A. has lumenal construction when enter the pelvic brim

11- Ingunal LN mean Mets from which internal organ

A. overy

B. uterus

C. falobian tube

12- Most precancerous organ and can predictable by screening?

A. Ovary

B. Vulva

C. Cervix.

D. Endometrium

13- Breakdown of 1gm Haemoglobin produces:

a. 20 mg of bilirubin

b. 35 mg of bilirubin

c. 50 mg of bilirubin

d. 70 mg of bilirubin

14- The active transport system for hepatic uptake of bilirubin is congenitally defective in:

a. Gilbert's disease

b. Crigler-Najjar syndrome type 1

c. Rotor's syndrome

d. Crigler-Najjar syndrome type 2

15- Excretion of conjugation bilirubin from liver cells into biliary canaliculi is defective in:

a. Gilbert's disease

b. Crigler-Najjar syndrome

c. Lucey-Driscoll syndrome

d. Rotor's syndrome

16- In iron deficiency anaemia:

a. mobilization of hemosiderin reserves.

b. activation of ferritin phage

c. elevated intestine apo ferritin

17. Vasopressin and oxytocin circulate unbound to proteins and have very short plasma half-lives on the order of :

a. 1-2 minutes

b. 2-4 minutes

c. 5-8 minutes

d. 10-12 minutes

18-A study involving analysis of risk for genetic defects in a family is:

a. Genetics Engineering

b. Genetics counselling

d. Genetic equilibrium

c. Genetics drift

19-Biological activity of ACTH is completely lost on removal of :

a. 10 N-terminal amino acids

b. 24 N-terminal amino acids

c. 24 C-terminal amino acids

d. 15 C-terminal amino acids

20-Hyperglycaemic effect of glucocorticoids is due to:

a. inactivation of protein phosphatase

b. inactivation of fructose 1,6-bisphosphatase

c. stimulation of synthesis of pyruvate carboxylase

d. stimulation of synthesis of eltrexykinase

21-Cortisol is synthesized in:

a. zona fasiculata

b. zona glomerulosa

c. zona reticularis

d. chromaffin cells

22-Mineralocorticoids are steroid hormones produced from:

a. zona fasiculata

b. zona glomerulosa

c. zona reticularis

d. zona pellucida

23-Catecholamine are released from:

a. zona fasiculata

b. zona glomerulosa

c. zona reticularis

d. chromaffin cells

24-Measles is characterized by:

a. Negribodies

b. Babes-Ernest granules

c. Koplik's spots

d. Fever

25-Assessment of fetal anemia :

a. Doppler umbilical artery = Fetal well being

b. Doppler umbilical vein

c. middle cerebral artery = Fetal anemia

d. fetal heart rate by pinard

26-Red infarction develop in:

a. Spleen

b. Liver

c. Kidney

d. Intestine

27-The number of amino acids in the hormone oxytocin is:

a. 7

b. 9

c. 14

d. 18

28-very efficient inhibitor of steroid biosynthesis is:

a. Aminoglutethimide

b. Aminoimidazole

c. Aminoimidazolesuccinyl carboxamine

d. Aminopterin

29-In the synthesis of Angiotensin I, rennin acts on Angiotensinogen and cleaves the:

a. Leucine - leucine at 10 and 11 position

b. Valine - tyrosine at 3 and 4 position

c. Isoleucine - histidine at 5 and 6 position

d. Proline - histidine at 7 and 8 position

30-Which of the normal range of ionized calcium in plasma is:

a. 2-4 mg/dl

b. 2-4 mEq/L

c. 4-5 mg/dl

d. 4-5 mEq/L

31-Lymphogranuloma venerum (LGV) is sexually transmitted disease is caused by:

a. C.opthalmia

b. C.trachomatis

c. C.pneumonias

d. C.psittasi

32-Koplic's spots will develop in:

a. HIV

b. Measles

c. Mumps

d. Rubella

33-Scrup typhus is caused by:

a. R.Quintana

b. R.rickettsii

c. R.orinetalis

d. R.prowazekii

34-grading of tumour is best described as:

a. degree of differentiation

b. size of tumour

c. lymph nodes

d. metastasis

35-pure hypertrophy is seen in:

a. breast of pregnant women

b. uterus of pregnant women

c. muscle of athletes

36- female came to your clinic has hirsutism and acne, her BMI 37, which means she is obese and has DM, what is the Pathophysiology of PCO:

a. conversion of oestrogen into androgen

b. hyperinsulinemia

c. increase level of FSH

37- Healing by first intention:

a. the surface of wound is rapidly covered by epithelium

b. excessive granulation tissue

c. wound will covered by fibroblast

38- Tetanus is caused by spread of:

a. exotoxin in sympathetic system

b. exotoxin in parasympathetic system

c. endotoxin in sympathetic system

d. endotoxin in parasympathetic system

39- Phenotype is determined by:

a. sex chromosome (XX- XY) karyotyping

b. gonadal differentiation

c. steroid hormones

40- increase volume in pregnancy due to:

a. increase thirst threshold b.inhibition of ADH

c. inhibition of ANP d.stimulation of aldosterone

41- After treatment of myocardial infarction, the injury occurs instead of restore blood supply by:

a. free radical

42- mature placenta composed of:

a. syncytiotrophoblast, cytotrophoblast

b. maternal blood vessels, syncytiotrophoblast, fetal blood vessels

c. syncytiotrophoblast, fetal blood vessels

d. syncytiotrophoblast, cytotrophoblast, mesenchyma

43-external genitalia development:

a. genital tubercle give rise to clitoris

b. clitoris between two labia majora

c. genital folds become labia majora

d. glans derived from genital swelling

44-Complement based agglutination reaction is known as:

a. haem agglutination

b. complement fixation

c. conglutination

d. Schultz dale phenomenon

45-Theca cell produce primerly:

a. progesterone

b. androsterone

c. estrogen

d. DHEAS

46-Mechanism of action of Tamoxifen:

a. selective oestrogen receptive modulator

b. aromatase inhibitor

c. progesterone receptor antagonist

47-About the anterior abdominal wall: Except

a. Median umbilical ligament represented by urachus

c. ligamentum of teres remnant of umbilical vein

d. pyramidalis m. presented in 20% of population

48-Poly A tail is frequently found in:

a. Histone in RNA

b. Bacterial RNA

c. Eukaryotic RNA

d. tRNA

49. Wound healing by regeneration invariable in:

a. heart

b. renal tubules

c. brain

d. intestine + Stem cells .... Labile cells

50- anterior division of internal iliac artery:

a. lateral sacral artery

b. iliolumbar artery

c. obturator artery

51- When total lung capacity = 5L, tidal volume = 2.5L, IVR= 1, ERV= 1 what is residual volume?

a. 1.5 L

b. 2 L

c. 2.5 L

d. 0.5 L

52- About spermatogenesis:

a. nucleus form the cap

b. mitochondria make the neck sheath

c. the centriole elongated to form axial filament

d. tail have two and nine microtubules

53- Epithelium cell tumour:

a. Papilloma

b. Cystadenoma

c. Choriocarcinoma

d. Nephroblastoma

54- The technique involved in comparing the DNA component of two sample is known as:

a. monoclonal Ab technique

b. genetic finger printing

c. recombinant DNA technology

d. PCR

55- At what stage of the following life is compatible Extra uterine:

a. glandular

b. canalicular

c. scavular

d. after birth

56- Which of the following statement regarding TFT in pregnancy is true:

a. normal total, normal T4

b. normal total, increase T4

c. increase total, increase T4

d. increase total, normal freeT4

57- Thyroperoxidase used H2O2 as oxidative agent, H2O2 come from:

a. FAD dependent enzyme

b. NAD dependent enzyme

c. NADPH dependent enzyme

58- In the congenital syphilis:

a. snuffles nose + Hutchison teeth

b. ascending infection through vagina

c. chancer in tertiary syphilis

d. severe if occurs in first trimmest

Effect of increase cortisol in pregnancy:
decrease placental blood flow and decrease birth weight
- Increase gluconeogenesis and glycogenolysis
Decrease liver protein contents

59- Effect of cortisol in pregnancy:

a. increase the liver protein binding protein b.

c. d

60- Regarding early stages of development:

a. zygote contain blastomer b.

c. implantation occur after 10 days

d. inner cell mass becomes placenta

61- Innervation of majority of Vulva?

a. Ilioinguinal nerve

b. Pudendal nerve

c. Inferior rectal nerve

d. Perineal branch of S4

62- Central chemoreceptor occurs by:

a. CO2

b. O2

c. PH

d. H

63- Cell of chronic inflammation is:

a. macrophages with giant cell

b. polymorph nuclear

c. monocyte

d. plasma cell

64- Which of the following is true about exudate:

a. occurs in heart failure

b. due to chronic inflammation

c. used in cytology cultures

d. has low specific gravity

65- leukocyte adhesion by:

a. selectin

b. integrin

c. mucin

d. fibronectin

66- Progesterone : Glandular stage

a. convert estradiol to estron b.

c. causes uterine contraction

d. decrease thickness of the cervical mucosa

68- Which of the following viruses causes an acute febrile rash and pro- duces disease in immunocompetent children but has been associated with transient aplastic crises in persons with sickle cell disease?

a. Rubeola

b. Varicella-zoster

c. Parvovirus

d. Rubella

e. Herpes simplex

69- The latest and most effective therapy for AIDS patients includes azidothymidine (AZT), dideoxyinosine (DDI), and saquinavir or similar agents. Use of these three drugs would inhibit which of the following viral processes?

a. RNase, DNase

b. gp120 formation

c. p24 antibody expression

d. All membrane synthesis

e. Reverse transcriptase, protease

70-N. gonorrhoeae is a fastidious pathogen and found in sites often contaminated with normal flora. The best medium for isolation is

a. Sheep blood agar

b. Löffler's medium

c. Thayer-Martin agar

d. Thiosulfate citrate bile salts sucrose medium

e. Löwenstein-Jensen medium

71- The class of antibiotics known as the quinolones are bactericidal. Their mode of action on growing bacteria is thought to be?

a. Inhibition of DNA gyrase

b. Inactivation of penicillin-binding protein II

c. Inhibition of B-lactamase

d. Prevention of the cross-linking of glycine

e. Inhibition of reverse transcriptase

72- Krukenberg Tumor of the Ovary is an example of:

a) Lymphatic Spread

b) Vascular Spread

c) Transcolomic Spread

d) Intraductal Spread

e) Transluminal Spread

73- Pyruvate carboxylase

a. Converts acetyl CoA to pyruvate

b. Requires carnitine

c. Converts pyruvate to oxaloacetate

d. Is activated by high fructose 1,6 bisphosphate

e. Requires pyridoxal phosphate

74- newborn with periventricular calcification commonly associated with:-

a-Toxoplasmosis

b-Cytomegalovirus

c-rubella d-syphilis

75- patient with HBV +ve and have cor infection with HDV what true indicator for patient coinfection of HDV :

a. HBsAg

b. HBc IgM

c. HBeAg

d. HBs IgM

76- infection of women reproductive organs its often caused by STDs its spread from vagina to uterus to fallopian tube or ovary :

a.PID

b. trachomonus vaginalis

c.c.UTI

77- Glomerular filtration rate would be increased by:

a. increase in afferent arteriolar pressure

b. decrease in efferent arteriolar resistance

c. increase in the concentration of plasma proteins

d. compression of renal capsule

78- In the ECG the p wave denotes:

a. atrial contraction

b. atrial depolarization

c. ventricular repolarization

d. SA node depolarization

79- which substance increased effective renal surface area ;

a. ADH

b. histamine

c. Endothelin

d. dopamine

80- ammonia is a good buffer in urine because

a. renal produce large amounts of NH3 in chronic acidosis

b. renal wall impermeable to NH3

c. renal wall permeable to NH4

d. reabsorbed in thick ascending loop

81- the lower limb..

a. Femoral ring is 1.25 inch lateral and below the pubic tubercle

b. The anterior superior illiac spine is at spinous level of second sacral vertebra

c. The greater sciatic notch lies at the level of third sacral vertebra

d. The line that meet the anterior superior iliac spine with most prominent point in ischial tuberosity pass directly through the acetabulum

81- malignant tumor in trigone of the bladder metastasize to the following lymph nodes:

a.lumbar

b.sacral

c.external iliac only

d.external and internal iliac

82- reversible inhibition of cycloxgenase in platelet is caused by?

A-alprostadil

B-asprin

C-paracetamol

D-ibuprofien

83- Cardiotoxicity limits the clinical usefulness of which one of the following antitumor antibiotics :

a.Doxorubicin

b. Bleomycin

c. Dactinomycin

d. Plicamycin

84-A nucleophilic attack on deoxyribonucleic acid which cause disruption of base pairing occurs as a result :

a) Cyclophosphamide

b) Thioguanine

c) Prednisone

d) Methotrexate

85-9 yrs female diagnosed with TB. before prescribing treatment you should be aware of that one antituberculus induce cytochrome P450 enzyme in the liver :

a. rifampicin

b. isoniazid

c. pyrazinamide

d. ethambiotol

86-Coombs test is positive and associated with heamolytic anaemia in :

a. methyledopa

b. captopril

c. clonidine

87-45 years old treated for ovarian cancer developed difficulty hearing the cause is:

a. paclitaxil

b. bleomycin

c. cisplatin

88-benzodiazipine action:

a. enhance GABA B neurotransmitters

b. enhance GABA A in spinal cord

c. enhanced Cl channels that act on GABA neurotransmitters 89-amiloride potent diuretic action by:

a. aldesteron antagonist

b. acts on ENaC

c. Na -K blocker

90-primary mechanism of flurosemide:

a. inhibition of sodium potassium ATPase

b. inhibtion of Na k cl co-transporter

c. inhibtion of Na Cl co-transporter

d. e- inhibtion of Cl transporter