

---

Homework for  
**MA 346 Numerical Methods**  
Spring 2022 — Homework 6

---

**Submit via Canvas before April 17, 2022, 11:59 p.m..**

**Exercise 1 (Numerical Integration)**

Consider the following ordinary differential equation (ODE):

$$\frac{du(t)}{dt} = f(t, u(t)) \quad \text{on } [0, T] \quad \text{with } u(0) = u_0, \quad (1)$$

where  $u_0 \in \mathbb{R}$  is given and  $f$  being continuously differentiable with respect to both arguments. To find an approximation to the function  $u(t)$  that solves (1), one can proceed as follows: First one applies the Fundamental Theorem of Calculus (see first lecture) to obtain:

$$u(t) = u(0) + \int_0^t f(s, u(s)) ds. \quad (2)$$

We may iterate this procedure as follows:

$$u(t_i) = u(t_{i-1}) + \int_{t_{i-1}}^{t_i} f(s, u(s)) ds, \quad (3)$$

where  $t_i = i\Delta t$ ,  $i = 1, \dots, n$ , and  $\Delta t = T/n$ ; we decomposed the interval  $[0, T]$  in  $n$  small subintervals  $[t_{i-1}, t_i]$  of length  $\Delta t$ .

As a second step one approximates the integral on the right-hand side of (3) with a quadrature formula.

- a) Employ the following quadrature rule

$$\int_a^b g(x) dx \approx (b-a)g(a)$$

to approximate the integral on the right-hand side in (3) and therefore obtain a numerical method that approximates  $u(t_i)$  for  $t_i = i\Delta t$ ,  $i = 1, \dots, n$ .

- b) Employ the following quadrature rule

$$\int_a^b g(x) dx \approx (b-a)g(b)$$

to approximate the integral on the right-hand side in (3) and therefore obtain a numerical method that approximates  $u(t_i)$  for  $t_i = i\Delta t$ ,  $i = 1, \dots, n$ .

c) What difference do you observe between the numerical methods you obtained in a) and b)?

d) Employ the trapezoidal rule

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{2} (g(a) + g(b))$$

to approximate the integral on the right-hand side in (3) and therefore obtain a numerical method that approximates  $u(t_i)$  for  $t_i = i\Delta t$ ,  $i = 1, \dots, n$ .

e) Which method (a), b), or d)) do you think will converge the fastest to the true values of  $u(t_i)$  for a decreasing  $\Delta t$ ? Give a brief justification.