

An Introduction to the equivariant Tamagawa Number Conjecture

Harry Spencer

University College, London

17th May 2023

Introduction/Structure

The eTNC is a prediction about the behaviour at $s = 0$ of motivic L -functions.

It is an attempt to generalise both the analytic class number formula and the Birch–Swinnerton-Dyer conjecture.

Introduction/Structure

The eTNC is a prediction about the behaviour at $s = 0$ of motivic L -functions.

It is an attempt to generalise both the analytic class number formula and the Birch–Swinnerton-Dyer conjecture.

Its general statement is unwieldy and opaque, so we will discuss one of the ‘simplest’ special cases and end with some discussion on numerical evidence for the eTNC.

Introduction/Structure

The eTNC is a prediction about the behaviour at $s = 0$ of motivic L -functions.

It is an attempt to generalise both the analytic class number formula and the Birch–Swinnerton-Dyer conjecture.

Its general statement is unwieldy and opaque, so we will discuss one of the ‘simplest’ special cases and end with some discussion on numerical evidence for the eTNC.

- 1 Determinant Modules
- 2 (A very small amount of) Algebraic K -Theory
- 3 BSD & the eTNC
- 4 Numerical Examples

References etc.

The first part of this talk and some of the general philosophy are based on notes of David Burns.

The latter part is all based on Werner Bley's paper 'Numerical evidence for the equivariant Birch and Swinnerton-Dyer conjecture, part I'.

Determinant modules

We give a brief introduction to *determinant modules*, restricting to the case of free R -modules M of finite rank r :

Definition

$$[M]_R = \bigwedge^r M \cong R \quad \text{and} \quad [M]_R^{-1} = \operatorname{Hom}_R([M]_R, R).$$

Determinant modules

We give a brief introduction to *determinant modules*, restricting to the case of free R -modules M of finite rank r :

Definition

$$[M]_R = \bigwedge^r M \cong R \quad \text{and} \quad [M]_R^{-1} = \text{Hom}_R([M]_R, R).$$

This extends to finitely generated R -modules M for $R = \mathbb{Q}[G], \mathbb{C}[G]$, etc. for finite groups G by writing

$$R = \prod_i F_i \quad \text{and} \quad M = \bigoplus_i M_i$$

for F_i fields and M_i a free F_i -module, and taking

$$[M]_R = \prod_i [M_i]_{F_i}.$$

Determinant modules

This construction has the following properties:

- 1 Given $\mathcal{E} : 0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0$, we obtain canonical

$$\iota(\mathcal{E}) : [N]_R \xrightarrow{\sim} [M]_R \otimes_R [P]_R.$$

Determinant modules

This construction has the following properties:

- 1 Given $\mathcal{E} : 0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0$, we obtain canonical

$$\iota(\mathcal{E}) : [N]_R \xrightarrow{\sim} [M]_R \otimes_R [P]_R.$$

- 2 We have canonical isomorphism

$$\mathrm{ev}_M : [M]_R \otimes_R [M]_R^{-1} \rightarrow R$$

by $m \otimes f \mapsto f(m)$.

Determinant modules

This construction has the following properties:

- 1 Given $\mathcal{E} : 0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0$, we obtain canonical

$$\iota(\mathcal{E}) : [N]_R \xrightarrow{\sim} [M]_R \otimes_R [P]_R.$$

- 2 We have canonical isomorphism

$$\text{ev}_M : [M]_R \otimes_R [M]_R^{-1} \rightarrow R$$

by $m \otimes f \mapsto f(m)$.

- 3 Given $f : M \xrightarrow{\sim} N$, we obtain canonical isomorphism

$$t(f) : [M]_R \otimes_R [N]_R^{-1} \xrightarrow{[f]_R \otimes 1} [N]_R \otimes_R [N]_R^{-1} \xrightarrow{\text{ev}_N} R,$$

where $[f]_R$ is the map induced by f .

Determinant modules

To explicate the link to determinants, we note that for $f : M \xrightarrow{\sim} N$ the following commutes

$$\begin{array}{ccc} [M]_R [N]_R^{-1} & \xrightarrow{t(f)} & R \\ \beta_M \otimes \beta_N \downarrow & & \uparrow \times \det(\Phi) \\ R \otimes R & \xrightarrow{\text{id}} & R \end{array}$$

where the maps β_\bullet are given by a choice of basis and Φ is the matrix of f with respect to the chosen bases.

Determinant modules

To explicate the link to determinants, we note that for $f : M \xrightarrow{\sim} N$ the following commutes

$$\begin{array}{ccc} [M]_R [N]_R^{-1} & \xrightarrow{t(f)} & R \\ \beta_M \otimes \beta_N \downarrow & & \uparrow \times \det(\Phi) \\ R \otimes R & \xrightarrow{\text{id}} & R \end{array}$$

where the maps β_\bullet are given by a choice of basis and Φ is the matrix of f with respect to the chosen bases.

We can do pretty much the same thing for other rings (e.g. $R = \mathbb{Z}[G]$), although we lose the fact that $[M]_R$ is a free rank one R -module in general.

Some algebraic K -theory

We now give a little background on algebraic K -theory.

Some algebraic K -theory

We now give a little background on algebraic K -theory.

For a ring R , we denote by $K_0(R)$ the free abelian group on isomorphism classes $[M]$ of projective R -modules modulo the relations

$$[M] = [N] + [P]$$

whenever there exists

$$0 \rightarrow N \rightarrow M \rightarrow P \rightarrow 0.$$

Some algebraic K -theory

We also have a 'relative K_0 '.

Some algebraic K -theory

We also have a 'relative K_0 '. We'll just define $K_0(\mathbb{Z}[G], F)$, where F is a characteristic 0 field.

Some algebraic K -theory

We also have a 'relative K_0 '. We'll just define $K_0(\mathbb{Z}[G], F)$, where F is a characteristic 0 field.

This is an abelian group whose elements can be represented by tuples $[P, \phi, Q]$ up to isomorphism, where P and Q are projective $\mathbb{Z}[G]$ -modules and

$$\phi : P \otimes_{\mathbb{Z}[G]} F[G] \xrightarrow{\sim} Q \otimes_{\mathbb{Z}[G]} F[G].$$

Some algebraic K -theory

We also have a 'relative K_0 '. We'll just define $K_0(\mathbb{Z}[G], F)$, where F is a characteristic 0 field.

This is an abelian group whose elements can be represented by tuples $[P, \phi, Q]$ up to isomorphism, where P and Q are projective $\mathbb{Z}[G]$ -modules and

$$\phi : P \otimes_{\mathbb{Z}[G]} F[G] \xrightarrow{\sim} Q \otimes_{\mathbb{Z}[G]} F[G].$$

As before, we can view this as a free abelian group on isomorphism classes modulo the relations generated by (pairs of) short exact sequences.

Some algebraic K -theory

We also have a 'relative K_0 '. We'll just define $K_0(\mathbb{Z}[G], F)$, where F is a characteristic 0 field.

This is an abelian group whose elements can be represented by tuples $[P, \phi, Q]$ up to isomorphism, where P and Q are projective $\mathbb{Z}[G]$ -modules and

$$\phi : P \otimes_{\mathbb{Z}[G]} F[G] \xrightarrow{\sim} Q \otimes_{\mathbb{Z}[G]} F[G].$$

As before, we can view this as a free abelian group on isomorphism classes modulo the relations generated by (pairs of) short exact sequences.

We have a canonical homomorphism

$$\delta : \zeta(F[G])^\times \rightarrow K_0(\mathbb{Z}[G], F).$$

Some algebraic K -theory

We can also consider elements of $K_0(\mathbb{Z}[G], F)$ as a tuple

$$((\hat{X}, Y, \hat{\theta}), \theta_\infty),$$

where:

Some algebraic K -theory

We can also consider elements of $K_0(\mathbb{Z}[G], F)$ as a tuple

$$((\hat{X}, Y, \hat{\theta}), \theta_\infty),$$

where:

- \hat{X} is of the form $[X_1]_{\hat{\mathbb{Z}}[G]}[X_2]_{\hat{\mathbb{Z}}[G]}^{-1}$;

Some algebraic K -theory

We can also consider elements of $K_0(\mathbb{Z}[G], F)$ as a tuple

$$((\hat{X}, Y, \hat{\theta}), \theta_\infty),$$

where:

- \hat{X} is of the form $[X_1]_{\hat{\mathbb{Z}}[G]}[X_2]_{\hat{\mathbb{Z}}[G]}^{-1}$;
- Y is of the form $[Y_1]_{\mathbb{Q}[G]}[Y_2]_{\mathbb{Q}[G]}^{-1}$;

Some algebraic K -theory

We can also consider elements of $K_0(\mathbb{Z}[G], F)$ as a tuple

$$((\hat{X}, Y, \hat{\theta}), \theta_\infty),$$

where:

- \hat{X} is of the form $[X_1]_{\hat{\mathbb{Z}}[G]}[X_2]_{\hat{\mathbb{Z}}[G]}^{-1}$;
- Y is of the form $[Y_1]_{\mathbb{Q}[G]}[Y_2]_{\mathbb{Q}[G]}^{-1}$;
- $\hat{\theta} : \hat{X} \otimes_{\hat{\mathbb{Z}}[G]} \hat{\mathbb{Q}}[G] \xrightarrow{\sim} Y \otimes_{\mathbb{Q}[G]} \hat{\mathbb{Q}}[G]$;

Some algebraic K -theory

We can also consider elements of $K_0(\mathbb{Z}[G], F)$ as a tuple

$$((\hat{X}, Y, \hat{\theta}), \theta_\infty),$$

where:

- \hat{X} is of the form $[X_1]_{\hat{\mathbb{Z}}[G]}[X_2]_{\hat{\mathbb{Z}}[G]}^{-1}$;
- Y is of the form $[Y_1]_{\mathbb{Q}[G]}[Y_2]_{\mathbb{Q}[G]}^{-1}$;
- $\hat{\theta} : \hat{X} \otimes_{\hat{\mathbb{Z}}[G]} \hat{\mathbb{Q}}[G] \xrightarrow{\sim} Y \otimes_{\mathbb{Q}[G]} \hat{\mathbb{Q}}[G]$;
- $\theta_\infty : Y \otimes_{\mathbb{Z}[G]} F[G] \xrightarrow{\sim} F[G]$.

Some algebraic K -theory

We can also consider elements of $K_0(\mathbb{Z}[G], F)$ as a tuple

$$((\hat{X}, Y, \hat{\theta}), \theta_\infty),$$

where:

- \hat{X} is of the form $[X_1]_{\hat{\mathbb{Z}}[G]}[X_2]_{\hat{\mathbb{Z}}[G]}^{-1}$;
- Y is of the form $[Y_1]_{\mathbb{Q}[G]}[Y_2]_{\mathbb{Q}[G]}^{-1}$;
- $\hat{\theta} : \hat{X} \otimes_{\hat{\mathbb{Z}}[G]} \hat{\mathbb{Q}}[G] \xrightarrow{\sim} Y \otimes_{\mathbb{Q}[G]} \hat{\mathbb{Q}}[G]$;
- $\theta_\infty : Y \otimes_{\mathbb{Z}[G]} F[G] \xrightarrow{\sim} F[G]$.

This works by mapping $[P, \phi, Q]$ to

$$((\prod_p [P_p]_{\mathbb{Z}_p[G]}[Q_p]_{\mathbb{Z}_p[G]}^{-1}, [P \otimes \mathbb{Q}[G]]_{\mathbb{Q}[G]}[Q \otimes \mathbb{Q}[G]]_{\mathbb{Q}[G]}^{-1}, \prod_p \theta_p), t(\phi)).$$

BSD & the eTNC

Now fix an elliptic curve E/\mathbb{Q} , a Galois extension K/\mathbb{Q} with Galois group G and set $S = \{\text{bad primes}\} \cup \{\text{ramified primes}\}$. We state the eTNC for E/K .

BSD & the eTNC

Now fix an elliptic curve E/\mathbb{Q} , a Galois extension K/\mathbb{Q} with Galois group G and set $S = \{\text{bad primes}\} \cup \{\text{ramified primes}\}$. We state the eTNC for E/K .

We define

$$\Xi = [E(K) \otimes_{\mathbb{Z}} \mathbb{Q}]_{\mathbb{Q}[G]}^{-1} [(E(K) \otimes_{\mathbb{Z}} \mathbb{Q})^*]_{\mathbb{Q}[G]} [H_B^+]_{\mathbb{Q}[G]}^{-1} [H_{\text{dR}}/H_{\text{dR}}^0]_{\mathbb{Q}[G]}.$$

BSD & the eTNC

Now fix an elliptic curve E/\mathbb{Q} , a Galois extension K/\mathbb{Q} with Galois group G and set $S = \{\text{bad primes}\} \cup \{\text{ramified primes}\}$. We state the eTNC for E/K .

We define

$$\Xi = [E(K) \otimes_{\mathbb{Z}} \mathbb{Q}]_{\mathbb{Q}[G]}^{-1} [(E(K) \otimes_{\mathbb{Z}} \mathbb{Q})^*]_{\mathbb{Q}[G]} [H_B^+]_{\mathbb{Q}[G]}^{-1} [H_{\text{dR}}/H_{\text{dR}}^0]_{\mathbb{Q}[G]}.$$

We have an isomorphism $(E(K) \otimes_{\mathbb{Z}} \mathbb{Q})^* \xrightarrow{\sim} E(K) \otimes_{\mathbb{Z}} \mathbb{Q}$.

BSD & the eTNC

Now fix an elliptic curve E/\mathbb{Q} , a Galois extension K/\mathbb{Q} with Galois group G and set $S = \{\text{bad primes}\} \cup \{\text{ramified primes}\}$. We state the eTNC for E/K .

We define

$$\Xi = [E(K) \otimes_{\mathbb{Z}} \mathbb{Q}]_{\mathbb{Q}[G]}^{-1} [(E(K) \otimes_{\mathbb{Z}} \mathbb{Q})^*]_{\mathbb{Q}[G]} [H_B^+]_{\mathbb{Q}[G]}^{-1} [H_{\text{dR}}/H_{\text{dR}}^0]_{\mathbb{Q}[G]}.$$

We have an isomorphism $(E(K) \otimes_{\mathbb{Z}} \mathbb{Q})^* \xrightarrow{\sim} E(K) \otimes_{\mathbb{Z}} \mathbb{Q}$. Combining with the period isomorphism (from Deligne's conjecture), we obtain

$$\theta_{\infty} : \Xi \otimes_{\mathbb{Q}[G]} \mathbb{R}[G] \rightarrow \mathbb{R}[G].$$

Define $T_\ell = \mathbb{Z}_\ell[G] \otimes_{\mathbb{Z}_\ell} T_\ell(E)$.

Define $T_\ell = \mathbb{Z}_\ell[G] \otimes_{\mathbb{Z}_\ell} T_\ell(E)$.

There exists a complex $R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)$ with isomorphisms

$$\theta_\ell : \Xi \otimes_{\mathbb{Q}[G]} \mathbb{Q}_\ell[G] \xrightarrow{\sim} [R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)] \otimes_{\mathbb{Q}[G]} \mathbb{Q}_\ell[G],$$

Define $T_\ell = \mathbb{Z}_\ell[G] \otimes_{\mathbb{Z}_\ell} T_\ell(E)$.

There exists a complex $R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)$ with isomorphisms

$$\theta_\ell : \Xi \otimes_{\mathbb{Q}[G]} \mathbb{Q}_\ell[G] \xrightarrow{\sim} [R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)] \otimes_{\mathbb{Q}[G]} \mathbb{Q}_\ell[G],$$

from which we obtain

$$R\Omega := ((\prod_{\ell} [R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)], \Xi, \prod_{\ell} \theta_\ell^{-1}), \theta_\infty) \in K_0(\mathbb{Z}[G], \mathbb{R}).$$

Define $T_\ell = \mathbb{Z}_\ell[G] \otimes_{\mathbb{Z}_\ell} T_\ell(E)$.

There exists a complex $R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)$ with isomorphisms

$$\theta_\ell : \Xi \otimes_{\mathbb{Q}[G]} \mathbb{Q}_\ell[G] \xrightarrow{\sim} [R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)] \otimes_{\mathbb{Q}[G]} \mathbb{Q}_\ell[G],$$

from which we obtain

$$R\Omega := ((\prod_\ell [R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)], \Xi, \prod_\ell \theta_\ell^{-1}), \theta_\infty) \in K_0(\mathbb{Z}[G], \mathbb{R}).$$

Our hope is that this element will encode data about the leading terms at $s = 1$ of $L(E/\mathbb{Q}, \chi, s)$ for characters χ of G .

Define $T_\ell = \mathbb{Z}_\ell[G] \otimes_{\mathbb{Z}_\ell} T_\ell(E)$.

There exists a complex $R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)$ with isomorphisms

$$\theta_\ell : \Xi \otimes_{\mathbb{Q}[G]} \mathbb{Q}_\ell[G] \xrightarrow{\sim} [R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)] \otimes_{\mathbb{Q}[G]} \mathbb{Q}_\ell[G],$$

from which we obtain

$$R\Omega := ((\prod_\ell [R\Gamma_c(\mathbb{Z}_{S_\ell}, T_\ell)], \Xi, \prod_\ell \theta_\ell^{-1}), \theta_\infty) \in K_0(\mathbb{Z}[G], \mathbb{R}).$$

Our hope is that this element will encode data about the leading terms at $s = 1$ of $L(E/\mathbb{Q}, \chi, s)$ for characters χ of G . In particular,

Conjecture (eTNC)

$$T\Omega := R\Omega + \delta((L^*(E/\mathbb{Q}, \chi, 1))_{\chi \in Irr(G)}) = 0.$$

Conjecture (Rationality)

$$u := \frac{R}{\Omega_{Reg}} \cdot (L^*(E/\mathbb{Q}, \chi, 1))_{\chi} \in \zeta(\mathbb{Q}[G])^{\times}.$$

Conjecture (Rationality)

$$u := \frac{R}{\Omega \text{Reg}} \cdot (L^*(E/\mathbb{Q}, \chi, 1))_{\chi} \in \zeta(\mathbb{Q}[G])^{\times}.$$

Here Ω is a 'period', Reg is an 'equivariant regulator', and R is a resolvent.

Conjecture (Rationality)

$$u := \frac{R}{\Omega \text{Reg}} \cdot (L^*(E/\mathbb{Q}, \chi, 1))_{\chi} \in \zeta(\mathbb{Q}[G])^{\times}.$$

Here Ω is a 'period', Reg is an 'equivariant regulator', and R is a resolvent.

Explicitly:

$$\Omega_{\chi} = \Omega_{+}(E/K)^{d^{+}(\chi)} \Omega_{-}(E/K)^{d^{-}(\chi)}.$$

Conjecture (Rationality)

$$u := \frac{R}{\Omega \text{Reg}} \cdot (L^*(E/\mathbb{Q}, \chi, 1))_{\chi} \in \zeta(\mathbb{Q}[G])^{\times}.$$

Here Ω is a 'period', Reg is an 'equivariant regulator', and R is a resolvent.

Explicitly:

$$\Omega_{\chi} = \Omega_{+}(E/K)^{d^{+}(\chi)} \Omega_{-}(E/K)^{d^{-}(\chi)}.$$

For ψ an abelian character and $\chi = \sum_{\sigma \in \text{Gal}(\mathbb{Q}(\psi)/\mathbb{Q})} \psi^{\sigma}$, we consider the $e_{\chi} \mathbb{Q}[G]$ -vector space $e_{\chi}(E(K) \otimes_{\mathbb{Z}} \mathbb{Q})$.

Conjecture (Rationality)

$$u := \frac{R}{\Omega \text{Reg}} \cdot (L^*(E/\mathbb{Q}, \chi, 1))_{\chi} \in \zeta(\mathbb{Q}[G])^{\times}.$$

Here Ω is a 'period', Reg is an 'equivariant regulator', and R is a resolvent.

Explicitly:

$$\Omega_{\chi} = \Omega_{+}(E/K)^{d^{+}(\chi)} \Omega_{-}(E/K)^{d^{-}(\chi)}.$$

For ψ an abelian character and $\chi = \sum_{\sigma \in \text{Gal}(\mathbb{Q}(\psi)/\mathbb{Q})} \psi^{\sigma}$, we consider the $e_{\chi} \mathbb{Q}[G]$ -vector space $e_{\chi}(E(K) \otimes_{\mathbb{Z}} \mathbb{Q})$. Pick an $e_{\chi} \mathbb{Q}[G]$ -basis P_1, \dots, P_d .

Numerical evidence

Conjecture (Rationality)

$$u := \frac{R}{\Omega \text{Reg}} \cdot (L^*(E/\mathbb{Q}, \chi, 1))_{\chi} \in \zeta(\mathbb{Q}[G])^{\times}.$$

Here Ω is a 'period', Reg is an 'equivariant regulator', and R is a resolvent.

Explicitly:

$$\Omega_{\chi} = \Omega_{+}(E/K)^{d^{+}(\chi)} \Omega_{-}(E/K)^{d^{-}(\chi)}.$$

For ψ an abelian character and $\chi = \sum_{\sigma \in \text{Gal}(\mathbb{Q}(\psi)/\mathbb{Q})} \psi^{\sigma}$, we consider the $e_{\chi} \mathbb{Q}[G]$ -vector space $e_{\chi}(E(K) \otimes_{\mathbb{Z}} \mathbb{Q})$. Pick an $e_{\chi} \mathbb{Q}[G]$ -basis P_1, \dots, P_d . We have

$$\text{Reg}_{\chi} = (\det(\langle P_i, e_{\psi^{\sigma}} P_j^* \rangle))_{\sigma \in \text{Gal}(\mathbb{Q}(\chi)/\mathbb{Q})}.$$

Numerical evidence

Lastly, the resolvent R is defined as follows:

Numerical evidence

Lastly, the resolvent R is defined as follows: pick a normal basis element α_0 for K/\mathbb{Q} and set

$$R_\chi = R_\chi(\alpha_0) = \det\left(\sum_{\sigma \in G} \sigma(\alpha_0) \chi(\sigma^{-1})\right).$$

Numerical evidence

Lastly, the resolvent R is defined as follows: pick a normal basis element α_0 for K/\mathbb{Q} and set

$$R_\chi = R_\chi(\alpha_0) = \det\left(\sum_{\sigma \in G} \sigma(\alpha_0) \chi(\sigma^{-1})\right).$$

We now take the following hypotheses:

- The rationality conjecture holds;
- $\text{III}(E/K)$ is finite;
- All the ramified primes in K are primes of good reduction for E .

Numerical evidence

Assuming these, Bley shows:

Theorem

There exists a(n explicit) finite set of primes HP such that, for $\ell \notin HP$,

$$T\Omega_\ell = 0 \in K_0(\mathbb{Z}_\ell[G], \mathbb{C}_\ell) \iff u \text{ has support in } HP.$$

Numerical evidence

Assuming these, Bley shows:

Theorem

There exists a(n explicit) finite set of primes HP such that, for $\ell \notin HP$,

$$T\Omega_\ell = 0 \in K_0(\mathbb{Z}_\ell[G], \mathbb{C}_\ell) \iff u \text{ has support in } HP.$$

Here, $u = (u_1, \dots, u_r) \in \zeta(\mathbb{Q}[G])^\times$ 'has support in HP ' if $(u_i, p) = 1$ for all i and all $p \notin HP$.

Numerical evidence

Assuming these, Bley shows:

Theorem

There exists a(n explicit) finite set of primes HP such that, for $\ell \notin HP$,

$$T\Omega_\ell = 0 \in K_0(\mathbb{Z}_\ell[G], \mathbb{C}_\ell) \iff u \text{ has support in } HP.$$

Here, $u = (u_1, \dots, u_r) \in \zeta(\mathbb{Q}[G])^\times$ 'has support in HP ' if $(u_i, p) = 1$ for all i and all $p \notin HP$.

HP should contain the primes dividing $|\text{III}(E/K)|$, but we have no good way to compute this.

Numerical evidence

Assuming these, Bley shows:

Theorem

There exists a(n explicit) finite set of primes HP such that, for $\ell \notin HP$,

$$T\Omega_\ell = 0 \in K_0(\mathbb{Z}_\ell[G], \mathbb{C}_\ell) \iff u \text{ has support in } HP.$$

Here, $u = (u_1, \dots, u_r) \in \zeta(\mathbb{Q}[G])^\times$ 'has support in HP ' if $(u_i, p) = 1$ for all i and all $p \notin HP$.

HP should contain the primes dividing $|\text{III}(E/K)|$, but we have no good way to compute this. Hence we will have to assume BSD for our numerical computations.

Numerical evidence

To compute u , we need to compute the (equivariant) regulator;
hence we need to compute $E(K)$.

Numerical evidence

To compute u , we need to compute the (equivariant) regulator; hence we need to compute $E(K)$.

We use:

Theorem (Longo, Tian-Zhang)

Suppose E/K is a modular elliptic curve over a totally real number field. Then

$$\mathrm{ord}_{s=1} L(E/K, s) = 0 \implies \mathrm{rk}(E/K) = 0.$$

Example

Example

Consider the elliptic curve

$$y^2 + y = x^3 - x^2 - 10x - 20$$

over the splitting field K of $f(x) = x^3 - 4x + 1$, an S_3 -extension of \mathbb{Q} .

Example

Consider the elliptic curve

$$y^2 + y = x^3 - x^2 - 10x - 20$$

over the splitting field K of $f(x) = x^3 - 4x + 1$, an S_3 -extension of \mathbb{Q} .

Using Magma, one can easily compute that $\text{rk}(E/K) = 0$ and $E(K) = E(\mathbb{Q}) \cong C_5$.

Example

Consider the elliptic curve

$$y^2 + y = x^3 - x^2 - 10x - 20$$

over the splitting field K of $f(x) = x^3 - 4x + 1$, an S_3 -extension of \mathbb{Q} .

Using Magma, one can easily compute that $\text{rk}(E/K) = 0$ and $E(K) = E(\mathbb{Q}) \cong C_5$. We also find

$$(L^*(E/\mathbb{Q}, \chi, 1))_{\chi} = (0.253842, 0.419359, 2.66127).$$

Example

Consider the elliptic curve

$$y^2 + y = x^3 - x^2 - 10x - 20$$

over the splitting field K of $f(x) = x^3 - 4x + 1$, an S_3 -extension of \mathbb{Q} .

Using Magma, one can easily compute that $\text{rk}(E/K) = 0$ and $E(K) = E(\mathbb{Q}) \cong C_5$. We also find

$$(L^*(E/\mathbb{Q}, \chi, 1))_{\chi} = (0.253842, 0.419359, 2.66127).$$

Moreover, from BSD we predict $|\text{III}(E/K)| = 5^3$.

Example

Consider the elliptic curve

$$y^2 + y = x^3 - x^2 - 10x - 20$$

over the splitting field K of $f(x) = x^3 - 4x + 1$, an S_3 -extension of \mathbb{Q} .

Using Magma, one can easily compute that $\text{rk}(E/K) = 0$ and $E(K) = E(\mathbb{Q}) \cong C_5$. We also find

$$(L^*(E/\mathbb{Q}, \chi, 1))_{\chi} = (0.253842, 0.419359, 2.66127).$$

Moreover, from BSD we predict $|\text{III}(E/K)| = 5^3$. We take $HP = \{2, 3, 5, 11, 229\}$.

Example

After computing the resolvent, we find

$$u = (0.200000, -5.00000, -25.0000),$$

Example

After computing the resolvent, we find

$$u = (0.200000, -5.00000, -25.0000),$$

so we take

$$u' = \left(\frac{1}{5}, -5, -25\right).$$

We see that u' has support in HP , so it is left to consider the primes of HP .

Example

After computing the resolvent, we find

$$u = (0.200000, -5.00000, -25.0000),$$

so we take

$$u' = \left(\frac{1}{5}, -5, -25\right).$$

We see that u' has support in HP , so it is left to consider the primes of HP .

For $\ell = 11, 229$, the group $K_0(\mathbb{Z}_\ell[G], \mathbb{Q}_\ell)$ is trivial.

Example

After computing the resolvent, we find

$$u = (0.200000, -5.00000, -25.0000),$$

so we take

$$u' = \left(\frac{1}{5}, -5, -25\right).$$

We see that u' has support in HP , so it is left to consider the primes of HP .

For $\ell = 11, 229$, the group $K_0(\mathbb{Z}_\ell[G], \mathbb{Q}_\ell)$ is trivial. For $\ell = 3$, we find u is $K_0(\mathbb{Z}_\ell[G], \mathbb{Q}_\ell)_{\text{tors}} \cong C_2$ and can compute that it is trivial.

Example

After computing the resolvent, we find

$$u = (0.200000, -5.00000, -25.0000),$$

so we take

$$u' = \left(\frac{1}{5}, -5, -25\right).$$

We see that u' has support in HP , so it is left to consider the primes of HP .

For $\ell = 11, 229$, the group $K_0(\mathbb{Z}_\ell[G], \mathbb{Q}_\ell)$ is trivial. For $\ell = 3$, we find u is $K_0(\mathbb{Z}_\ell[G], \mathbb{Q}_\ell)_{\text{tors}} \cong C_2$ and can compute that it is trivial.

Hence we have verified the eTNC for E/K away from $\ell = 2, 5$.

Closing remarks

- This computational approach can also tell us about the Galois module structure of $\mathrm{III}(E/K)$.

Closing remarks

- This computational approach can also tell us about the Galois module structure of $\mathrm{III}(E/K)$.
- I lied to you!

Closing remarks

- This computational approach can also tell us about the Galois module structure of $\mathrm{III}(E/K)$.
- I lied to you! To properly do all this, we need a more nuanced definition of determinant modules (Picard categories, virtual objects, and more...).

Closing remarks

- This computational approach can also tell us about the Galois module structure of $\mathrm{III}(E/K)$.
- I lied to you! To properly do all this, we need a more nuanced definition of determinant modules (Picard categories, virtual objects, and more...).
- To state the eTNC for general motives, we need to replace $E(K) \otimes_{\mathbb{Z}} \mathbb{Q}$ by motivic cohomology.