

the 1990s, the incidence of *S. flexneri* serotype 3 infections has increased in the United Kingdom [10]. In the United States, the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].

There is a paucity of data on the incidence of *S. flexneri* serotype 3 infections in the United Kingdom. In the 1980s, *S. flexneri* serotype 3 was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [12]. In the 1990s, *S. flexneri* serotype 3 was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [13]. In the 2000s, *S. flexneri* serotype 3 was the most commonly isolated serotype from patients with acute bacterial dysentery in the United Kingdom [14].

The purpose of this study was to determine the incidence of *S. flexneri* serotype 3 infections in the United Kingdom in the 2000s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11]. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11]. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11]. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11]. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11]. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11]. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11]. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11]. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* serotype 3 infections has increased since the 1970s [11].