

Venus Photochemistry

M.Tech. Project, Phase 1 Presentation



Suhas H

Roll Number: 25M0328

Under the Supervision of

Prof. Deepak Singh

Geoinformatics and Natural Resources Engineering
Centre of Studies in Resources Engineering


INDIAN INSTITUTE OF TECHNOLOGY BOMBAY

Mumbai – 400076, India

Outline

1. Introduction
2. The Model and the Method
3. Experiments 1 to 7
4. Experiments 8 to 14
5. Findings and Verification
6. Phase 2
7. Conclusion
8. References and Sources


Venus and its clouds



Dai et al. (2025)

- Surface: 92 bar, about 750 K.
- Sulfuric acid clouds, 47 to about 70 km.
- Cloud top: 65 to 72 km.
- Grey band: where the model responds, 69 to 93 km.

The unknown ultraviolet absorber



Observed: MESSENGER spectrum, digitised from Pérez-Hoyos et al. (2018), Figure 8. Grey band: 320 to 400 nm.

Pérez-Hoyos et al. (2018); Dai et al. (2025); Titov et al. (2008)

- Dark at 320 to 400 nm; SO_2 cannot explain it.
- About half the solar heating at the cloud top.
- Amount varied twofold, 2006 to 2017.
- Its markings trace vertical motion.

The candidates proposed so far

Candidate	For	Against
Sulfur dioxide, SO_2	Matches below 320 nm	Negligible beyond 320 nm
Polysulfur, S_x	Early spectral candidate	No form matches
Ferric chloride, $FeCl_3$	Dissolves in the acid	Band too narrow and weak
Disulfur monoxide, S_2O	Matches 400 to 500 nm	Fades at 320 to 340 nm
OSSO, S_2O_2	Matches 330 to 420 nm	Far too little at cloud top
Iron and sulfur minerals	Absorb at 300 to 500 nm	Presence unknown
Chlorine and sulfur compounds	Spectra close	No abundance given
Organic “red oil”	Dissolves in acid	Speculative
A mixture of absorbers	Explains the band’s width	Not yet tested

Almost every case rests on the spectrum alone.

Dai et al. (2025); Pérez-Hoyos et al. (2018)

Literature survey: twelve papers, three tests

Paper	Focus	Spectrum	Amount	Altitude
Dai et al. (2024)	The model, OSSO	◔	◑	◔
Pérez-Hoyos et al. (2018)	Observed absorber	◑	○	◔
Bierson and Zhang (2020)	Mixing, OSSO	◔	◑	◑
Rimmer et al. (2021)	Droplet chemistry	◑	◔	◔
Lefèvre et al. (2022)	Cloud convection	○	◔	○
Lefèvre et al. (2024)	With chemistry	○	○	○
Jiang et al. (2024)	Iron sulfates	◑	◑	○
Trabelsi and Francisco (2026)	Chlorine and sulfur	◑	○	○
Egan et al. (2025)	OSSO, end to end	●	●	◑
Spacek et al. (2026)	Strength needed	◑	●	◔
Egan, James and Plane (2025)	$FeCl_3$, laboratory	◑	○	○
Egan et al. (2026)	$FeCl_3$, 3D model	●	◑	◔
This project	**All in one model**	◑	◑	●

Key

- ○ Not addressed
- ◔ Assumed or set
- ◑ Partly tested
- ● Tested in full

Partly: shape only, an estimate, or no light calculation.
Full: absolute, against observation.


The twelve papers reviewed; five more references are background.

In the literature, altitude is never tested in full: it is assumed, set, or left out.

Research gap and objective


Gap: no study lets a model place a candidate and measures the effect of where it sits.

This project: all three tests at once, inside one model



The model: VULCAN-Venus

How an absorber acts on the model




Dai, Shao and Sheng (2024); Bierson and Zhang (2020)

Set-up of every Phase 1 run

Grid	0 to 112 km, 57 layers
Chemistry	62 entries, 838 reaction slots
Wavelengths	1670 bins, 96 to 698 nm
Sunlight	One angle, 58 degrees
Mixing	Bierson and Zhang
Off	Vertical wind


The groundwork: an absorber that was switched off



Control and treatment differ in one term only.

What was done, in four stages

Fourteen experiments; each stage builds on the one before.



Measuring a change, and where it happens

One number for a whole field

$$\frac{\text{Size of the change in sunlight}}{\text{Size of the reference sunlight}}$$

Size: root of summed squares over 57 layers by 1670 bins; in per cent

Absorber moved up	2.2498%
Absorber moved down	0.3067%

The response band: narrowest range holding a share

Share asked	Range, km	Width, km	Held
50 per cent	73 to 83	10	54.567%
80 per cent	71 to 89	18	81.479%
90 per cent	**69 to 93**	**24**	**91.370%**
99 per cent	65 to 103	38	99.138%

- Centre of the response: 82.044 km.
- Absorber particles sit lower: 90.194% at 58 to 69 km.

The reproducibility floor

Experiments 9 and 10 should agree; their gap is numerical noise.


Quantity	Experiment 9	Experiment 10	Verdict
Field of sunlight	differ by 0.0017% of the change		Resolved
SO inside the band	0.7384%	2.5278%	Numerical

The rule

In-band abundance changes below 2.5278% are upper bounds, not measurements.


The two floors

The chemistry floor applies to abundance changes only.



Experiments 1 and 2: does the absorber do anything?


Switched on, the absorber changes sunlight strongly and the sulfur species barely.



	Largest change	Verdict
Sunlight	87.404%	resolved
SO_2	0.2243%	below floor
SO	1.7399%	below floor

Experiment 1: the control matches the supplied model, mean change 0.013553%.

Experiment 3: is the response proportional?




Half, nominal and twice the amount: the response scales with it.

Test, where 1 means proportional	Ratio
Optical depth, half amount	1.00002
Sunlight, half amount	1.00208
Whole field, half amount	1.010795
Whole field, twice the amount	0.979304

- Chemistry: within the floor, not resolved.

Experiment 4: which wavelengths?




Each run limited to one window changes sunlight only in that window.

Molecule	Main window, nm	Share
S_2O	300 to 320	93.689%
SO_3	300 to 320	80.754%
S_3	380 to 420	49.605%
S_4	420 to 620	100%

- Each molecule draws on its own part of the spectrum.

Experiments 5 and 6: which reactions, which altitudes?




The response sits in one band, 69 to 93 km, for every wavelength window tested.

Reaction	In the band	Peak
SO_2 photolysis	99.128%	71 km
SO_3 photolysis	99.992%	73 km
COS + 2 SO_3	almost none	35 km

- Direct effects sit in the band; knock-on effects lie lower.


Experiment 5: the reactions behind the SO_2 change


The largest absolute change is not the most affected reaction.



Experiment 7: does placement matter?

Moved up, the same absorber has 7.34 times more effect than moved down.






Move	Change
Up, +17.8439 km	2.2498%
Down, −18.0000 km	0.3067%
Ratio	**7.34**

- Same amount, same spectrum.
- Moved up, SO in the band: 8.404%.

Experiment 8: spectrum against placement


Same amount, same height, three spectra: shape matters, placement matters more.



- Spectra scaled to one total; optical depth above 69 km within 0.6461%.

Experiment 9: the chemistry places the candidate

Placed by its own chemistry, ClS_2 falls out of the response band.




	Set	Chemistry
Centre, km	63.419	13.656
In band	9.1940%	0.0000%

- Closest in amount: short by 121.914.
- Optical depth above 69 km: 2.0×10^8 times less.
- Same as no absorber, to 0.001487%.


Experiment 10: a replicate, and a retraction

S_3 on the ClS_2 profile gave the same optical depth: a replicate of Experiment 9.



Experiment 11: two computed isomer spectra

The authors' cross sections run as absorbers; neither isomer can be made in the amount needed.




Trabelsi and Francisco (2026)

	$SSCl_2$	ClSSCl
Peak height	0.114%	0.079%
Peak position	0.156 nm	0.331 nm
Field change	0.19465%	0.30144%
Short by	4.266×10^5	6.668×10^6

- $SSCl_2$: 85.49% of its absorption at 320 to 400 nm, the most tested.
- ClSSCl peaks below 300 nm; the model sees its tail.

Experiment 12: all three tests on every species


Twenty-three species screened; none passes all three tests.



Every pass mark is one the empirical absorber, which reproduces the observed brightness of Venus, itself meets.

Experiment 12: the three tests disagree

Amount is a gate; on the other two tests each picks a different best species.



Best eight on each test, best first


smallest shortfall	most column in the band	best spectral match
NO_2	NO_3	Cl_2
SO_2	NO_2	OSSO
S_3	$ClCO_3$	SCl_2
OSSO	SO_3	HOCl
HNO_2	ClO	HNO_2
O_3	S_2	S_2Cl_2
Cl_2	HOCl	H_2O_2
ClS_2	OSSO	NO_2

Each test picks a different winner: the rankings disagree.

no absorption at this wavelength, so not placed: $ClCO_3$, ClO, H_2O_2, H_2SO_4, HNO_3, NO_3, S_2, S_2O, S_4, SO_2Cl_2, SO_3

Experiment 13: iron and sulfur minerals

Even a droplet made wholly of either mineral could not absorb enough at 375 nm.




- Requirement: 1278 per cm at 375 nm; coefficients as measured by Jiang et al.

Jiang et al. (2024); Spacek et al. (2026)

Experiment 14: ferric chloride in acid

Measured in hydrochloric acid, ferric chloride needs 2.82% to 17.05% of the droplet, not 71.0%.



Most chloride-rich curve: a factor of 2.6 from the independent Egan et al. (2026) fit, against 64.5 before the correction

- The rejection rested on water; altitude is still untested.

Spacek et al. (2026); Egan, James and Plane (2025); Egan et al. (2026)

What Phase 1 establishes

Result	Basis
The absorber acts within one band, 69 to 93 km	Measured
Placement beats spectrum: 7.34 against 4.95	Measured
Placed by its own chemistry, ClS_2 equals no absorber	Measured
No species in the model passes all three tests	Calculated, model output
Minerals fall short; the ferric chloride rejection is reopened	Calculated, published data
Chemical responses mostly below the floor	Bounded

- Measured: radiative, reproducible to 0.0017% of the change.
- Bounded: abundances, reproducible only to 2.5278%.

The limits

Limit	What it means
One column, one sun angle	No latitude, no local time, steady states only
Weak vertical mixing	Placement tested at one mixing strength
Simplified droplets	Fitted cross sections; full droplet calculation off
Empirical reference	Spectral rankings measure resemblance to a fit
Only what the network holds	23 of 62 species could be screened
Floor from one pair of runs	A lower bound on the numerical scatter

Every run shares these limits, so comparisons stand; Phase 2 takes up the first two.

Verification: no number typed by hand


Every number is recomputed from the saved output and checked on the page.



- The checker proves a number is in the registry, not that it is the right one.
- So every number was also checked by hand against its own entry.

Verification: reading one registry entry


Each name reads from left to right, from the general to the particular.



Value 2.4563×10^5: the model holds that many times too little OSSO above 69 km.


Phase 2: fast winds, slow rising

Mixing needs 552 days to cross the 55 to 69 km gap; the winds circle Venus in 4 to 6.



Limaye et al. (2018); Bertaux et al. (2016)


Phase 2: where the layers sit



Lefèvre et al. (2022)

- Deep convective layer: 50 to 60 km.
- Cloud-top layer near 70 km, at noon only.
- It overlaps the response band.
- Its authors call it still speculative.

Phase 2: a second mixing profile, never used




Bierson and Zhang (2020); Dai et al. (2023); Lefèvre et al. (2024)

- Every Phase 1 run used the weak profile.
- The unused file: 8716 to 14011 times stronger at 47 to 53 km.
- Crossing that layer: 118 days becomes 17 minutes.
- It stops at 53 km, below the band.

Phase 2: six work packages

Does placement survive realistic transport, and can transport lift a candidate into the band?



- WP1 and WP5 come first; together they can give a result on their own.
- WP3 runs alongside WP2; ferric chloride comes last.

Conclusion

- Switched on, the absorber acts in one band: 69 to 93 km.
- Where an absorber sits matters more than its spectrum.
- The chemistry, not the modeller, decides where a candidate sits.
- No species in the model passes all three tests.
- Minerals fall short; ferric chloride is reopened, its altitude untested.
- Every number traces back to the saved output.

Next: can realistic mixing carry a candidate into the band?

References

Bertaux, J.-L., et al. (2016). Influence of Venus topography on the zonal wind and UV albedo at cloud top level: the role of stationary gravity waves. *J. Geophys. Res. Planets* 121, 1087 to 1101.

Bierson, C. J., and Zhang, X. (2020). Chemical cycling in the Venusian atmosphere: a full photochemical model from the surface to 110 km. *J. Geophys. Res. Planets* 125, e2019JE006159.

Dai, L., Shao, W., Gu, H., and Sheng, Z. (2023). Determination of the eddy diffusion in the Venusian clouds from VeRa sulfuric acid observations. *Astron. Astrophys.* 679, A155.

Dai, L., Shao, W., and Sheng, Z. (2024). An investigation into Venusian atmospheric chemistry based on an open-access photochemistry-transport model at 0 to 112 km. *Astron. Astrophys.* 689, A55.

Dai, L., et al. (2025). Venus cloud research: progress and perspectives. *Space Sci. Rev.* 221, 51.

Egan, J. V., et al. (2025). Is OSSO a significant contributor to the unknown UV absorber in Venus' atmosphere? *Geophys. Res. Lett.* 52, e2024GL113090.

Egan, J. V., James, A. D., and Plane, J. M. C. (2025). Laboratory measurements of ferric chloride ($FeCl_3$) under Venusian conditions. *ACS Earth Space Chem.*

Egan, J. V., et al. (2026). Ferric chloride: the Venusian unknown UV absorber? Preprint, ESS Open Archive, not peer reviewed.

Jiang, C. Z., et al. (2024). Iron-sulfur chemistry can explain the ultraviolet absorber in the clouds of Venus. *Sci. Adv.* 10, eadg8826.

References

Lefèvre, M., Marcq, E., and Lefèvre, F. (2022). The impact of turbulent vertical mixing in the Venus clouds on chemical tracers. *Icarus* 386, 115148.

Lefèvre, M., et al. (2024). Impact of the turbulent vertical mixing on chemical and cloud species in the Venus cloud layer. *Geophys. Res. Lett.* 51, e2024GL108771.

Limaye, S. S., et al. (2018). Venus' spectral signatures and the potential for life in the clouds. *Astrobiology* 18(9), 1181 to 1198.

Pérez-Hoyos, S., et al. (2018). Venus upper clouds and the UV absorber from MESSENGER/MASCS observations. *J. Geophys. Res. Planets* 123, 145 to 162.

Rimmer, P. B., et al. (2021). Hydroxide salts in the clouds of Venus: their effect on the sulfur cycle and cloud droplet pH. *Planet. Sci. J.* 2, 133.

Spacek, J., Rimmer, P. B., Petkowski, J. J., and Lee, Y. J. (2026). A model of UV-blue absorbance in bulk liquid of Venusian cloud aerosols is consistent with efficient organic absorbers at high concentrations. *Astrobiology* 26(9), 727 to 739.

Titov, D. V., et al. (2008). Atmospheric structure and dynamics as the cause of ultraviolet markings in the clouds of Venus. *Nature* 456, 620 to 623.

Trabelsi, T., and Francisco, J. S. (2026). Chlorine-sulfur isomers as parents of ClS_2 and SCl_2 on Venus: spectroscopy and photochemistry of ClSSCl, $SSCl_2$, and $(ClS)_2$. *J. Chem. Phys.* 164, 084302.

Numbers and their sources (1 of 4)

Number	What it is	Registry entry	Report
1.095	Optical depth needed at 320 nm	`const.tau_req`	3.6
2.2498%	Field change, absorber moved up	`e7a.L2pct`	5.6
0.3067%	Field change, absorber moved down	`e7b.L2pct`	5.6
7.34	Ratio of the two moves	`e7.ratio_7A_over_7B`	5.6
69 to 93 km	Response band, 90 per cent share	`e6.interval.90.lo, .hi`	4.5
91.370%	Share held by that band	`e6.interval.90.contained`	4.5
54.567, 81.479, 99.138%	Shares held by the 50, 80, 99 per cent bands	`e6.interval.<share>.contained`	4.5
82.044 km	Centre of the response	`e6.A1.meanz`	4.5
90.194%	Absorber particles at 58 to 69 km	`pnom.cloud`	4.5
0.0017%	Radiation floor, share of the change	`e10.fielddiff_pct`	6.3
0.7384, 2.5278%	SO in the band, Experiments 9 and 10	`res.e9.SO.inband, res.e10.SO.inband`	6.3
0.013553%	Experiment 1, mean change	`e1.aflux.meanrel`	5.1

Set-up values come from the model files (report Table 3.1); literature values are cited on their slides.

Numbers and their sources (2 of 4)

Number	What it is	Registry entry	Report
87.404%	Largest change in sunlight	`e3fp.bnd.A1.aflux.max_pct`	5.1
0.2243, 1.7399%	Largest SO_2 and SO changes	`e2.SO2.maxrel, e2.SO.maxrel`	5.1
1.00002, 1.00208	Proportionality, optical depth and flux	`e3.lin.tau.inband.median, e3.lin.aflux.inband.median`	5.2
1.010795, 0.979304	Proportionality at half and twice the amount	`e3fp.aflux.ratio.A0p5, .A2p0`	5.2
93.689, 80.754, 49.605, 100%	Main-window shares, S_2O, SO_3, S_3, S_4	`e4a.<species>_1.<window>.pct`	5.3
99.128%, 71 km	SO_2 photolysis in the band, its peak	`e6.R781.pct_in_band, .peak_z`	5.5
99.992%, 73 km	SO_3 photolysis in the band, its peak	`e6.R787.pct_in_band, .peak_z`	5.5
35 km	Peak of the knock-on reaction	`e6.R593.peak_z`	5.5
17.8439, 18.0000 km	Centre moved up, moved down	`e7.7a.centroid_shift, e7.7b.centroid_shift`	5.6
8.404%	SO in the band, absorber moved up	`res.e7a.SO.inband`	5.6
0.6461%	Optical depth difference between spectra	`tau.spec_change_pct`	6.1
4.95	Largest over smallest spectral effect	`e8c.L2pct over e8b.L2pct`	6.1

Numbers and their sources (3 of 4)

Number	What it is	Registry entry	Report
121.914	ClS_2 shortfall, Experiment 8	`scr.ClS2.short`	6.1
63.419, 13.656 km	Centre, set and by chemistry	`pnom.centroid, p9.centroid`	6.2
9.1940, 0.0000%	Share in the band, set and by chemistry	`pnom.inband, p9.inband`	6.2
2.0×10^8	Fall in optical depth above 69 km	`tau.reduction`	6.2
0.001487%	Difference from no absorber	`id.l2_ratio_pct`	6.2
1.2365, 1.000003	S_3 over ClS_2, in and out of the band	`twoby2.nominal_ratio, .cls2_ratio`	6.3
1.0474, 1.0473	Withdrawn constant, flux ratio	`chr.mean, chr.flux_ratio`	6.3
0.114%, 0.156 nm	$SSCl_2$ peak match	`iso.SSCl2.val_pct, .val_nm`	6.4
0.079%, 0.331 nm	ClSSCl peak match	`iso.ClSSCl.val_pct, .val_nm`	6.4
0.19465, 0.30144%	Field change, $SSCl_2$ and ClSSCl	`e11a.L2pct, e11b.L2pct`	6.4
4.266×10^5, 6.668×10^6	Isomer shortfalls	`iso.SSCl2.short, iso.ClSSCl.short`	6.4
85.49%	$SSCl_2$ share at 320 to 400 nm	`share320400.11A`	6.4

Numbers and their sources (4 of 4)

Number	What it is	Registry entry	Report
23, 16, 2, 0	Screened, placement, both, all three	`e12.n_screened`, `e12.gate.n_placement`, `.n_place_and_spec`, `.n_all`	6.5
62	Species the model tracks	`e12.n_tracked`	6.5
2.4563×10^5	OSSO shortfall at 360 nm	`e12.S2O2.short.360`	8.1
1278 per cm	Absorption the droplets need at 375 nm	`e13.A_required`	6.6
71.0%	Ferric chloride need, aqueous	`e14.aqueous_pct_of_droplet`	6.7
2.82, 17.05%	Ferric chloride need, hydrochloric acid	`e14.ClA.pct_of_droplet`, `e14.ClE.pct_of_droplet`	6.7
552 days	Mixing across the 55 to 69 km gap	`p2.gap.Bkzz.t_days`	9.1
118 days, 17 minutes	Crossing 47 to 53 km, weak and raised	`p2.raised.Bkzz.t_days`, `p2.raised.BDkzz.t_minutes`	9.3
8716 to 14011	How much stronger the unused profile is	`p2.kzz.raised.ratio_min`, `.ratio_max`	9.3
10830	Values in the registry	all entries of `verify/registry.json`	8.1
407, 129	Numbers checked, report and figures	checker output	8.2

Thank you