

## Quiz #2 - Solution

2015/10/21

1. (8, 7 pts) Determine the constant  $c$  so that  $p(x)$  or  $f(x)$  satisfies the conditions of being a probability mass function (p.m.f) or a probability density function (p.d.f) for some random variable.

(i)  $p(x) = \frac{x}{c}$  ,  $x = 1, 2, \dots, n$ .

$$\sum_{x=1}^n \frac{x}{c} = \frac{1}{c} \frac{n(n+1)}{2} = 1 \Leftrightarrow c = \frac{2}{n(n+1)}$$

(ii)  $f(x) = \frac{3}{16}x^2$  ,  $-c \leq x \leq c$

$$\int_{-c}^c \frac{3}{16}x^2 dx = \frac{3}{16} \frac{1}{3} x^3 \Big|_{-c}^c = \frac{c^3}{8} = 1 \Leftrightarrow c = 2$$

2. (5, 8, 7 pts) The pdf of  $X$  is given by

$$f(x) = \frac{e^{-x}}{(1+e^{-x})^2} , \quad -\infty < x < \infty$$

(i) Verify that  $f(x)$  is a pdf.

$$\int_{-\infty}^{\infty} \frac{e^{-x}}{(1+e^{-x})^2} dx = \int_1^{\infty} \frac{1}{t^2} dt = 1 \quad (1+e^{-x} = t)$$

(ii) Show that

$$Y = \frac{1}{1+e^{-X}}$$

has a  $U(0,1)$  distribution.

$$F(x) = \frac{1}{1+e^{-x}} \quad \therefore F(X) = \frac{1}{1+e^{-X}} \sim U(0,1)$$

(iii) Find a function  $G$  such that

$$P(G(U) \leq x) = \frac{1}{1+e^{-x}} , \quad -\infty < x < \infty , \quad U \sim U(0,1)$$

$$P(F^{-1}(U) \leq x) = F(x) = \frac{1}{1+e^{-x}}$$

$$\therefore G(x) = F^{-1}(x) = \log\left(\frac{x}{1-x}\right)$$

3. (10, 10 pts) (i) Let the random variable  $X$  be equal to the number of days that it takes a high-risk driver to have an accident. Assume that  $X$  has an exponential distribution. If  $P(X < 50) = 0.25$ , compute  $P(X > 100 | X > 50)$ .

From the memoryless property of the exponential distribution,

$$P(X > 100 | X > 50) = P(X > 50) = 1 - P(X < 50) = 0.75$$

(ii) Suppose the birth weight ( $X$ ) in grams of U.S. infants has the pdf

$$f(x) = \frac{3x^2}{3500^3} e^{-(x/3500)^3}, \quad 0 < x < \infty.$$

Compute  $P(X > 4000 | X > 3000)$ .

$$F(x) = 1 - e^{-(x/3500)^3}$$

$$\therefore P(X > 4000 | X > 3000) = \frac{P(X > 4000)}{P(X > 3000)} = e^{(6/7)^3 - (8/7)^3}$$

4. (10 pts) Let  $X$  have an exponential distribution with  $\lambda = 1$ , that is, the pdf of  $X$  is  $f(x) = e^{-x}$ ,  $0 < x < \infty$ . Let  $T = \ln(X)$ . Find a p.d.f of  $T$ .

$$g(x) = \ln(x), \quad g^{-1}(x) = (g^{-1})'(x) = e^x$$

$$\therefore f_T(t) = f_X(e^t) |e^t| = e^{-e^t} e^t, \quad -\infty < t < \infty$$

5. (8, 7 pts) 어느 고속도로 톨게이트에는 10분당 평균 5대의 차가 지나가고, 지나가는 차의 대수는 포아송분포를 따른다고 한다.

(i) 차가 1대 지나가기까지 5분 이상 기다릴 확률은 무엇인가?

1분당 통과대수는 평균이 0.5인 포아송분포를 따르므로, 대기시간을  $Y$ 라 하면,  $Y \sim \text{Exp}(2)$ .

$$P(Y > 5) = e^{-\frac{1}{2} \cdot 5} = e^{-5/2}$$

(ii) 8대의 차가 지나가기까지 26.3분 이상 기다릴 확률은 무엇인가? ( $\chi_{0.05}^2(16) = 26.3$ )

8대의 차가 지나가기까지의 대기시간을  $Y'$ 이라 하면,  $Y' \sim \text{Gamma}(8, 2) = \chi^2(16)$

$$\therefore P(Y' > 26.3) = P(\chi^2(16) > \chi_{0.05}^2(16)) = 0.05$$

6. (5, 8, 7 pts) (5번에서 계속) 1분 간격으로 차가 톨게이트를 통과하는지 여부를 기록하였다 (적어도 한 대 이상이 통과하면 1, 한 대도 통과하지 않으면 0).

(i) 1분 내에 적어도 한 대 이상의 차량이 통과할 확률은 얼마인가?

$$\text{대기시간이 1분보다 짧을 확률이므로 } p = P(Y \leq 1) = 1 - e^{-\frac{1}{2}}$$

(ii) 총 20분 동안 기록한 결과, 차량이 통과한 시간대의 수를  $T$ 라 하면,  $P(T \geq 19)$ 를 계산하여라.

$$T \sim B(20, p), \quad \therefore P(T \geq 19) = P(T = 19) + P(T = 20) = \binom{20}{19} p^{19} (1-p)^1 + \binom{20}{20} p^{20} (1-p)^0 = p^{19} (20 - 19p)$$

(iii) 차량이 통과한 시간대가 3회 이상 발생할 때까지 기록을 계속한다고 할 때, 기록한 횟수가 5회 이상일 확률을  $\sum$  형태로 표현하여 보아라.

기록한 횟수를  $S$ 라 하면,  $S \sim NB(3, p)$

$$\therefore P(S \geq 5) = \sum_{k=5}^{\infty} \binom{k-1}{2} p^3 (1-p)^{k-3}$$