

Short Paper

Subtitle

Tran Quoc Hoang^{a,1,*}, Nguyen Hoang Tuong Vy^a, Nguyen Vy Van^a, Nguyen Le Ngoc Vy^a, Nguyen Hoang Quynh Anh^a

^aInternational University - VNU-HCM, School of Biotechnology, Quarter 33, Linh Xuan Ward, Ho Chi Minh City, Vietnam, 71309
^b,

Abstract

This is the abstract. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Vestibulum augue turpis, dictum non malesuada a, volutpat eget velit. Nam placerat turpis purus, eu tristique ex tincidunt et. Mauris sed augue eget turpis ultrices tincidunt. Sed et mi in leo porta egestas. Aliquam non laoreet velit. Nunc quis ex vitae eros aliquet auctor nec ac libero. Duis laoreet sapien eu mi luctus, in bibendum leo molestie. Sed hendrerit diam diam, ac dapibus nisl volutpat vitae. Aliquam bibendum varius libero, eu efficitur justo rutrum at. Sed at tempus elit.

Keywords: keyword1, keyword2

Submission Note: Paper prepared for the Microbiology (BT321IU) course, International University - VNU-HCM.

1. Introduction

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Integer nec odio. Praesent libero. Sed cursus ante dapibus diam. Sed nisi. Nulla quis sem at nibh elementum imperdiet. Duis sagittis ipsum. Praesent mauris. Fusce nec tellus sed augue semper porta. Mauris massa. Vestibulum lacinia arcu eget nulla. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos himenaeos.

2. Microbes that Changed History

This section examines the all-encompassing influence of pathogenic microbes on human civilization, focusing on two specific agents that altered demographic, economic, and social structures globally.

2.1. The Black Death and *Yersinia pestis*

The Black Death, a devastating pandemic caused by the bacterium *Y. pestis* that peaked between 1346 and 1353, serves as a primary example of microbial impact on human history. The disease is believed to have traveled along the Silk Road from Central Asia to Europe, facilitated by

the expansion of trade routes and the movement of armies [1, 2]. Upon reaching Europe, the plague caused a demographic collapse of unprecedented scale, killing an estimated 50% of the population in affected regions of Europe and Asia [3, 4].

The massive loss of life triggered profound societal and economic shifts. The sudden scarcity of labor empowered the surviving peasantry, which led to the decline of the feudal system as workers demanded higher wages and better conditions [5, 6]. The psychological trauma of the pandemic also reshaped religious and social attitudes, manifesting in extreme movements such as the flagellants and the persecution of minority groups [7, 8].

The causative agent of the plague is the bacterium *Yersinia pestis*. The transmission cycle historically involved the black rat (*Rattus rattus*) and the rat flea (*Xenopsylla cheopis*) [9]. When the flea bites an infected host, the bacteria block the flea's digestive tract, causing it to regurgitate the pathogen into the bloodstream of a new host during subsequent feeding attempts [10]. While bubonic plague was the most common form, the pneumonic form allowed for direct person-to-person transmission—thereby accelerating the spread [11, 12]. Fig. 1 demonstrates in detail how the plague is spread.

2.2. Smallpox and Global Eradication

Shifting focus to another microbial agent that fundamentally altered human history: Smallpox. Caused by the *Variola major* virus and carrying a fatality rate $\approx 30\%$, the disease played a decisive role in the colonization of the Americas. Following the arrival of Europeans, smallpox decimated indigenous populations, such as the Aztecs and Incas, who possessed no prior immunity [14]. This “virgin soil” epidemic killed tens of millions—perhaps up to 95% of

*Corresponding author

Email addresses: tranquochuang@protonmail.com (Tran Quoc Hoang), BTBTIU24133@student.hcmiu.edu.vn (Nguyen Hoang Tuong Vy), BTBTIU24126@student.hcmiu.edu.vn (Nguyen Vy Van), BTBTIU24131@student.hcmiu.edu.vn (Nguyen Le Ngoc Vy), BTBTIU24007@student.hcmiu.edu.vn (Nguyen Hoang Quynh Anh)

¹Student ID: BTBTWE24036




Fig. 1: Mechanisms of Plague (*Y. pestis*) Transmission and Disease Progression. The cycle begins with (1) the Flea Life Cycle: Eggs, laid in moist soil/burrows, hatch into larvae that feed on rodent feces and develop into pupae and adult fleas. Infected adult fleas parasitize rodents. (2) The Enzootic Cycle maintains *Y. pestis* among rodent hosts and reservoirs, primarily vectored by fleas (*X. cheopis*, *X. brasiliensis*, and *S. fonquerniei*). (3, 4) Epizootic Spills occur when fleas infect non-host mammals, which often lack resistance, rapidly succumb to the disease, and spread the plague geographically. (5) Human Zoonotic Transmission: Humans are typically infected via the bite of an infected flea from a rodent or secondary mammal host. (6) Human-to-Human Spread: Bubonic plague may progress to a highly contagious pneumonic plague via lung infection, enabling *Y. pestis* transmission through infectious respiratory droplets generated by coughing. Reproduced from Mackay-Alderson J, Quastel M, Wilson E, Bellamy D [13].

the indigenous population of the Americas—far exceeding deaths caused by warfare and facilitating the collapse of indigenous empires [15, 16]. Due to a lack of reliable data, historians continue to debate the true mortality rates and extent of casualties.

The fight against smallpox marked the birth of immunology. In 1796, Edward Jenner observed that milkmaids who contracted cowpox, a mild disease, were immune to smallpox. He tested this hypothesis by inoculating a young boy with material from a cowpox lesion and subsequently exposing him to smallpox; the boy remained healthy [17, 18]. Jenner’s work established the principle of vaccination, a term derived from *vacca* (Latin for cow), replacing the riskier practice of variolatio, an earlier technique that intentionally induced a mild smallpox infection to confer immunity [19].

Smallpox holds the distinction of being the first and only human infectious disease to be eradicated. Following a massive global campaign by the World Health Organiza-

tion (WHO) involving vaccination, surveillance, and quarantine, the disease was declared eradicated in 1980 [20, 21]. This achievement stands as one of the greatest triumphs of public health and microbiology.

3. Famous Microbiologists and Groundbreaking Discoveries

This section highlights the contributions of three pivotal figures whose work transitioned medicine from superstition to the scientific germ theory of disease.

3.1. Louis Pasteur: The Father of Modern Microbiology

3.1.1. Germ Theory

Louis Pasteur, a French chemist, fundamentally changed the understanding of biology by disproving the theory of spontaneous generation. Through his elegant experiments using swan-necked flasks, he demonstrated that microorganisms in the air, not a “vital force,” caused contamination in sterile broth [22]. Fig. 11 shows the setup of

this experiment. Pasteur’s work laid the foundation for the germ theory of disease, which establishes that specific invisible microbes are responsible for specific illnesses [23].

3.1.2. Pasteurization

In addition to establishing the germ theory, Pasteur applied his knowledge of fermentation to solve problems in the French wine and silk industries. He discovered that heating liquids to a specific temperature (60–90°C) killed the bacteria responsible for spoilage without destroying the product. This process, known as pasteurization, revolutionized food safety and is still widely used today for milk and other beverages [24].

3.1.3. Vaccine Development

Pasteur also pioneered the development of laboratory-attenuated vaccines. He created vaccines for chicken cholera and anthrax by using weakened strains of the bacteria [25]. His most famous medical achievement was the development of the rabies vaccine in 1885, which he successfully used to save the life of Joseph Meister, a boy bitten by a rabid dog [26, 27].

3.2. Alexander Fleming: The Serendipitous Discovery

3.2.1. Discovery of Penicillin

In 1928, Scottish bacteriologist Alexander Fleming made a serendipitous discovery that launched the antibiotic era. Upon returning from a holiday, he noticed that a mold, *Penicillium notatum*, had contaminated a Petri dish of *Staphylococcus* bacteria. Crucially, he observed that the bacteria surrounding the mold had been destroyed [28].

3.2.2. Impact on Medicine

Fleming identified the mold’s active substance as penicillin. Although mass production was later achieved by Howard Florey and Ernst Chain during World War II, Fleming’s discovery provided the first effective treatment for bacterial infections that had previously been fatal [29, 30]. Penicillin was hailed as a “miracle drug,” saving millions of lives from wound infections and diseases like syphilis and pneumonia, though Fleming himself warned early on about the potential for antibiotic resistance [31].

3.3. Robert Koch and Koch’s Postulates

3.3.1. Methodological Rigor

Robert Koch, a German physician, is often considered the father of medical bacteriology alongside Pasteur. He introduced rigorous scientific methods to the field, most notably “Koch’s Postulates.” This set of four criteria is used to establish a causal relationship between a specific microbe and a specific disease: (1) the microbe must be present in every case of the disease; (2) it must be isolated and grown in pure culture; (3) inoculation of the culture into a healthy host must reproduce the disease; and (4) the microbe must be recovered again from the experimentally infected host [32, 33].

3.3.2. Key Discoveries

Using these methods, Koch identified the specific causative agents of several deadly diseases, including *Bacillus anthracis* (anthrax), *Vibrio cholerae* (cholera), and *Mycobacterium tuberculosis* (tuberculosis) [34, 35, 36]. His work provided the definitive proof for the germ theory of disease and established the standards for diagnostic microbiology [37].

4. Microbes in Art and Culture

This section explores the scientific mechanisms underlying microbial involvement in aesthetic and cultural domains, focusing on bacterial pigment production as a medium for art and the role of fungi and yeasts in pigmentation for textiles and sustainable dyes.

4.1. Bacterial Art (Agar Art/Bio-Painting)

Bacterial art is a form of art resulting from the special combination of microbiology with the power of visual expression by utilizing the Petri dish as a living canvas. This form of art relies entirely on the underlying biological mechanisms by which certain bacterial species synthesize brightly colored pigments [38].

The medium consists of nutrient-rich agar plates, where different bacterial strains are inoculated and cultured to grow into intricate images. The colors are generated by chromogenic bacterial secondary metabolites produced through distinct metabolic and biosynthetic pathways [source: 39]. The strong and visible colors of these pigments make it easier for scientists and artists to isolate and identify the strains that produce the compounds in sufficient amounts.

Two examples of bacterial species employed in biopainting are widely studied for their distinctive pigment production. The first is *Serratia marcescens*, a Gram-negative bacterium that is also a producer of prodigiosin. Prodigiosin is a red pigment belonging to the prodiginine group, characterized by a tripyrrole chemical structure, and has been successfully used in textile coloring processes [40]. The second example is *Chromobacterium violaceum*. This strain produces violacein, a distinctive bisindole-violet or blue pigment [41]. Violacein and prodigiosin are well-known hydrophobic bacterial chromogenic pigments, and these two are responsible for the purple and red color phenotype of the bacterial strain. Fig. 2 shows the chemical structure and the colored phenotypes of the bacterial strains that produce these compounds.

By integrating microbiological techniques with the principles of aesthetic design, bacterial art blurs the traditional boundaries between science and art. Fig. 3 shows the representation of various colors producing microorganisms on a Petri plate. Fig. 8 shows agar art from living microbes. Those are examples of colorful bacteria applied to agar in an artistic manner (A-C) ASM Agar Art Contest winners.




Fig. 2: Chemical structures of the bacterial pigments violacein (produced by *Chromobacterium violaceum*) and prodigiosin (produced by *Serratia marcescens*), shown alongside the respective colored phenotypes of the producing bacterial strains. Reproduced from Choi SY, Lim S, Yoon K, Lee JI, Mitchell RJ [39].

Fig. 9 shows The 2016 American Society for Microbiology Agar Art Contest winner, The First Race, illustrated fertilization using four pigment-producing bacterial species grown on selective agar. Created by Md Zohurul Islam (University of Copenhagen), the artwork used *Staphylococcus aureus* (red), *S. xylosus* (green), *S. hyicus* (white), and *Corynebacterium glutamicum* (yellow), with additional hues produced by mixing these microbes.

4.2. Fungi in Pigmentation and Textiles

The rapid development of greener approaches in industrial processes, such as textile dyeing, has been driven by the environmental concern of synthetic dyes [42]. In this context, fungi have become a promising sources of natural pigments.

Fungi have shaped the history of natural dyes for centuries throughout history, with uses like ancient lichen

dyes extracted from *Rocella* and *Ochrolechia* species for purples and violets. There were many traditional dyes made from lichens in the Scottish Highlands, including red dyes from the cudbear lichen (*Lecanorartartarea*), the common orange lichen (*Xanthoriaparietina*), and several species of leafy *Parmelia* lichens. During the 15th and 17th centuries, purple lichen dyes continued to play an important role in the European dyeing industry [43]. And today, the traditional practices have evolved into modern bio-dyes via fermentation to serve the demand of today's sustainable textile production demand [44].

From an industrial perspective, most fungal pigments are water-soluble, making production and extraction relatively straightforward. They can be easily cultivated and scaled up in industrial fermentation systems and can be harvested without the need for organic solvents, enhancing their sustainability and environmental compatibility (see Fig. 4).



Fig. 3: Representation of various colors producing microorganisms on a Petri plate. Reproduced from Tuli HS, Chaudhary P, Beniwal V, Sharma AK [40].

For reference, Table 1 shows List of fungal pigments produced on an industrial scale.

Fungal pigments cover a broad color spectrum, such as yellow, orange, red, purple, blue, brown, and black. Key producers are *Monascus* spp. (red, yellow, orange), *Penicillium* spp. (red, yellow, brown), *Aspergillus* spp. (yellow, brown), *Fusarium* spp. (pink, violet, red), and *Blakeslea trispora* (-carotene) [43]. Numerous fungal genera have demonstrated efficient pigment production [46]. For instance, *Penicillium brevicompactum* can produce a mixture of yellow, orange, and red pigments [47]. While *Penicillium purpurogenum* generates red pigments with strong dyeing performance on fabrics such as wool, silk and polyester. Fig. 5 shows 3 effect of red pigment produced by *Penicillium purpurogenum* application on different fabrics (wool, silk, polyester, viscose). Furthermore, different fungal species with their active pigment for application in the textile industry are shown in Fig. 10. Fungal pigments have demonstrated successful dyeing performance on a wide range of textile materials, including cotton, linen, silk, and several synthetic fibers [48, 49].

4.3. Microbes in Fiction and Media (Optional)

5. Bacteria in Space (Astromicrobiology)

Astromicrobiology investigates how microorganisms such as bacteria and archaea to survive, adapt, and behave under the extreme physical and chemical conditions found beyond Earth. This is crucial because many of Earth's microorganism, known as extremophiles already capable of thriving in hostile environments such as boiling hot springs, deep-sea hydrothermal vents, permafrost, acidic rivers, highly radioactive sites [50]. Their survival depend on their metabolisms and cellular structures evolved specifically for these extremes. Accurately simulating extraterrestrial ultraviolet (UV) radiation, which is significantly

more intense and has a wider wavelength range than what reaches Earth's surface, is one of the main challenges in astromicrobiology [51]. The solar UV spectrum cannot be completely replicated in space by even the most powerful artificial UV generators on Earth. Simulation studies are nevertheless useful in spite of this drawback. As an illustration of how extremophiles might take advantage of the chemical environment of other planets, the acidophilic bacteria *Acidithiobacillus ferrooxidans* was found to grow successfully in simulated Martian regolith, even without additional nutrients. In true space conditions, however, the most informative results come from orbital experiments such as the ADAPT study, which exposed genetically UV-resistant *Bacillus subtilis* spores to space radiation in Low Earth Orbit (LEO) [52].

5.1. Extremophiles Defined

Extremophiles are organisms, mostly bacteria and archaea, that not only survive but flourish in environments and conditions that were previously thought to be incompatible with life. The last point can be applied to a wide range of settings, including high pressure, temperature, pH, radiation, salinity, and nutrient availability, since they exist in severe habitats [53]. Extremophiles offer crucial insights about the boundaries of life on Earth and assist scientists in determining the potential habitability of alien settings since they live in such harsh circumstances.

Extreme settings can be either naturally occurring or man-made; these animals are just the result of evolution and the best adapted to their surroundings. Thousands of new species are found and named by scientists each year. Microorganisms have played a significant role in this massive expansion of species discoveries in recent years. Other terms that are used to categorize particular kinds of extremophiles include: Low pH acidophiles, high pH alkaliphiles, anaerobic extremophiles (anti-oxygen), cryophiles (thrive in cold temperatures), piezophiles (high pressure), psychrophiles (thrive in low temperatures), thermophiles (live in temperatures above 40°C), hyperthermophiles (above 80°C), xerophiles, methanogens, Toxitol-erance Deep-sea hydrothermal vents, ice sheets and permafrost, volcanic systems and hot springs, salt pans and hypersaline lakes, and extremely acidic or alkaline waters are just a few examples of the various types of extreme conditions found in nature [54, 55]. According to National Geographic, around 1.2 million species have been identified around the planet, but some experts speculate that 8,7 million or more may exist [56].

By surviving in such environments, extremophiles reveal the true boundaries of Earth-based life. This helps astrobiologists evaluate the potential habitability of extraterrestrial environments such as Mars, Europa, Enceladus, or exoplanets, and understand the biochemical strategies that life might use elsewhere. Extremophilic habitats may occur naturally (e.g., hydrothermal vents, volcanoes, permafrost) or be produced by human activity (industrial

brines, highly acidic mining runoffs). Regardless of origin, extremophiles are the best-adapted products of evolution within these conditions [57].

5.2. Survival on the International Space Station (ISS)

In the space experiment “Molecular adaption strategies of microorganisms to different space and planetary UV climate conditions” (ADAPT), bacterial endospores of the highly UV-resistant *Bacillus subtilis* strain MW01 were exposed to low-Earth orbit (LEO) and simulated martian surface conditions for 559 days on board the European Space Agency’s exposure facility EXPOSE-E, mounted outside the International Space Station. These facilities allow biological samples to be exposed directly to Space vacuum, Full-spectrum solar UV radiation (>110 nm), Cosmic ionizing radiation, Rapid temperature cycling, Simulated Martian surface conditions [58].

One of the main goals of Mars exploration space missions is to find organic compounds on the planet’s surface. Therefore, comprehending the preservation of organic matter in the Martian environment is a crucial step in interpreting future data obtained by these missions. The organic substances (glycine, serine, phthalic acid, phthalic acid in the presence of a mineral phase, and mellitic acid) completely degraded after a 1.5 year exposure to Mars-like surface UV radiation conditions in space. Under Martian surface conditions, their half-lives ranged from 50 to 150 hours. Amino acids and a dipeptide in pure form and embedded in meteorite powder were exposed to space conditions for eighteen months in order to study the chemical behavior of organic molecules in the space environment. The samples were then brought back to Earth and examined in a lab for reactions brought on by solar UV and cosmic radiation [59].

6. Conclusion

7. Abbreviations


| Abbreviation | Full Term |
|--------------|-----------------|
| PGx | Pharmacogenomic |
| DSA | Dolor sit amet |

8. Declaration of Competing Interest

The authors declare no competing interests.

9. Data/Code Availability

The Quarto project underlying this paper—including the manuscript, figures, and bibliography—is available at <https://github.com/ht2905/historical-microbes>.



A) Growth of pigmented fungi in agar medium



B) Maintenance of pigmented fungi

1. Control
2. *Penicillium purpurogenum*
3. *Paecilomyces farinosus*
4. *Emericella nidulans*
5. *Fusarium moniliforme*
6. *Monascus purpureus*



C) Scale-up of pigment producing fungi in fermenter

Fig. 4: Scale-up process for pigment-producing fungi from laboratory culture to industrial fermentation. The process moves from initial isolation and growth in a Petri dish to larger-scale production. (A) Growth and visualization of pigmented fungi on solid agar medium (Petri dish). (B) Maintenance and preservation of the fungal strain for long-term use. (C) Scale-up of the culture in a liquid medium within a fermenter, which is a critical step for sustainable and high-volume industrial pigment production. Reproduced from Venil CK, Velmurugan P, Dufossé L, Renuka Devi P, Veera Ravi A [45].



Fig. 5: 3 Effect of red pigment produced by *Penicillium purpogenum* application on different fabrics (wool, silk, polyester, viscose). Reproduced from Elkhateeb W, Elnahas M, Daba G [49].




Fig. 6: Representative idealized cross section of Earth's crust showing the diversity of extreme environments and their approximate location. Reproduced from [54].

10. Appendix A: Supplementary Figures




Fig. 7: Pasteur’s swan-necked flask experiment illustrating how airborne microorganisms, rather than a “vital force,” contaminate sterile broth. Reproduced from Shutterstock [60].

Table 1: List of fungal pigments produced on an industrial scale

| Fungal species | Pigments |
|------------------------------------|--|
| <i>Monascus</i> species | Ankaflavin (yellow), monascorubramine (red), rubropunctatin (orange) |
| <i>Ophiocordyceps unilateralis</i> | Erythrostominone (red), 3,5,8-TMON (red) |
| <i>Blakeslea tripora</i> | β -Carotene (yellow-orange), lycopene (red) |
| <i>Ashbya gossypii</i> | Riboflavin (yellow) |
| <i>Penicillium oxalicum</i> | Anthraquinone derivative (red), anthraquinones (red and other hues) |
| | Arpink red TM , secalonic acid D (yellow) |

Note. List of fungal pigments produced on an industrial scale, synthesized from Caro Y, Venkatachalam M, Lebeau J, Fouillaud M, Dufossé

L [63].




Fig. 8: Examples of Agar Art from Living Microbes. Panels (A-C) display winning entries from the ASM Agar Art Contest. (A) "Cell to cell" (2015) using yellow *Nesterenkonia*, orange *Deinococcus*, and *Sphingomonas*. (B) "Neurons" (2015) using red *Serratia*, yellow *Nesterenkonia*, orange *Deinococcus*, and *Sphingomonas*. (C) "Sustenance" (2018): (Top) pink colonies, orange *Nesterenkonia*, and *Deinococcus radiodurans*. (Bottom) Dark-violet recombinant *E. coli* (violacein pathway), red *Serratia marcescens*, and white *Bacillus*. (D) "Remainders": microbial paintings on food-colored agar plugs. (E) "Boston skyline" (2019): sculpted and collaged agar and bacteria. (F) "Bacillus surprise": dark-violet recombinant *E. coli* and white *Bacillus*. (G) A palette of colored recombinant yeast (*Saccharomyces cerevisiae*) expressing chromogenic proteins (black/grey/purple from violacein, blue from anemone gene, pink from RFP, and various yellows/oranges from beta-carotene). (H) "Puzzle Pieces" (2017) and (I) "Sunset at the End" (2016): patterns printed using engineered *S. cerevisiae*. Reproduced from Frankel E, Temple J, Dikener E, Berkmen M [61].

Table 2: Fungal pigments and their application in the textile industry

| Fungi | Pigment | Color | Fabrics |
|--|-------------------------|---------------|------------------------------------|
| <i>Alternaria alternata</i> | Anthraquinones | Reddish-brown | Cotton |
| <i>Penicillium Oxalium</i> | Anthraquinones | Arpink red | Wool |
| <i>Chlorociboria aeruginosa</i> | Quinones | Green | Bleached cotton, spun polyacrylic, |
| <i>Scytalidium cuboideum</i> | | Red | spun polyamide, spun polyester, |
| <i>Scytalidium ganodermorphothorum</i> | | Yellow | worsted wool |
| <i>Aspergillus</i> sp. | Quinones | Brown cotton | Silk, silk cotton |
| <i>Acrostalagmus</i> (NRC 90) | Quinones | Brown | Wool |
| <i>Alternaria alternata</i> (NRC17) | Quinones | Reddish-brown | Wool |
| <i>Alternaria</i> sp. (NRC 97) | Quinones | Brown | Wool |
| <i>Aspergillus niger</i> (NRC 95) | Quinones | Brown | Wool |
| <i>Bisporomyces</i> sp. (NRC 63) | Quinones | Deep brown | Wool |
| <i>Penicillium murcianum</i> | Cartenoids | Yellow | Wool |
| <i>Talaromyces australis</i> | 2,4-Di-tert-butylphenol | Red | Cotton fabric |
| <i>Phoma harbarum</i> | Magenta pigment | Magenta | Nylon |
| <i>Talaromyces verruculosus</i> | Polyketide | Red | Cotton fabric |
| <i>Monascus purpureus</i> | Rubropunctamine | Red | Wool |

Note. Fungal pigments and their application in the textile industry. Adapted from Elkhateeb W, Elnahas M, Daba G [49].







Fig. 9: The 2016 winner of the American Society of Microbiology's Agar Contest, titled *The First Race*, depicted fertilization. Graduate student Md Zohorul Islam of the University of Copenhagen "painted" with four bacteria on a selective agar canvas. The red was *Staphylococcus aureus*, a pathogen in both humans and animals. *Staphylococcus xylosum*, a commensal organism in human skin, generated green. The white was *Staphylococcus hyicus*, an animal pathogen responsible for grassy pig disease. And yellow came from *Corynebacterium glutamicum*, a bacterium used to produce amino acids, such as L-glutamate and L-lysine. Other colors came from mixing two or more of these microbes. Artwork by Md Zohorul Islam (University of Copenhagen, Copenhagen). Image courtesy of American Society for Microbiology. Reproduced from Madhusoodanan Y [62].




Xylindein (quinone pigment, dimeric naphthoquinone derivative, produced by fungi from genus *Chlorociboria*)


Aspergellone (pigment produced by fungi from genus *Aspergillus*, e.g. *A. awamori*, *A. niger*)



6-[(Z)-2-Carboxyvinyl]-N-gamma Aminobutyric Acid-PP-V (azaphilone pigment, produced by the fungus *Talaromyces albobiverticillius*)



Sclerotiorin (orange azaphilone pigment produced by fungi such as *Penicillium sclerotiorum*)



Torulene (A) and β -carotene (B) (carotenoid pigments, produced by the yeast *Sporidiobolus pararoseus*)




Fig. 10: Chemical structures of fungal pigments with potential coloring properties that could be used in textile dyeing. Reproduced from Venil CK, Velmurugan P, Dufossé L, Renuka Devi P, Veera Ravi A [45].




Fig. 11: A sankey diagram visualising the key medical astro-microbiology considerations relating to spaceflight and celestial body habitation. Reproduced from McDonagh et al. [64].




Fig. 12: EXPOSE-E payload, fully integrated and accommodated onto EuTEF at Kennedy Space Center, USA. Arranged vertically from left to right are tray 1 and tray 2—experiments in the four compartments of both trays from bottom compartment to top are ADAPT, PROTECT, $\frac{1}{2}$ ADAPT, and $\frac{1}{2}$ PROTECT sharing the third compartment and LIFE in the top compartment; on the right, separated from tray 2 by the three open lids and their motor drives, is tray 3 with R3DE in the bottom compartment, two compartments with PROCESS, and the top compartment with SEEDS. On the left side of EXPOSE-E, the experiment MEDET (wrapped in golden multilayer insulation) is located. Reproduced from Rabbow E et al. [58].

References

- [1] Black Death | Plague, Death Toll, Definition, Cause, Symptoms, Effects, & Facts | Britannica, <https://www.britannica.com/event/Black-Death> (Nov. 2025).
- [2] M. A. Spyrou, L. Musralina, G. A. Gnechi Ruscone, A. Kocher, P.-G. Borbone, V. I. Khartanovich, A. Buzhilova, L. Djansugurova, K. I. Bos, D. Kühnert, W. Haak, P. Slavin, J. Krause, The source of the Black Death in fourteenth-century central Eurasia, *Nature* 606 (7915) (2022) 718–724. doi:10.1038/s41586-022-04800-3.
- [3] The Black Death and European Expansion, <https://www.history.ox.ac.uk/black-death-and-european-expansion>.
- [4] J. Aberth, *From the Brink of the Apocalypse: Confronting Famine, War, Plague and Death in the Later Middle Ages*, 2nd Edition, Routledge, London, 2013. doi:10.4324/9781315881133.
- [5] Decameron Web | Plague, <https://webhelper.brown.edu/decameron/plague/effects/social.php>.
- [6] Labor Shortages Alter Europe's Social Structure | Research Starters | EBSCO Research, <https://www.ebsco.com/research-starters/history/labor-shortages-alter-europes-social-structure>.
- [7] J. J. Mark, Religious Responses to the Black Death, *World History Encyclopedia* (Apr. 2020).
- [8] M. H. Zentner, The Black Death and Its Impact on the Church and Popular Religion.
- [9] R. J. Eisen, A. P. Wilder, S. W. Bearden, J. A. Monteneri, K. L. Gage, Early-Phase Transmission of *Yersinia pestis* by Unblocked *Xenopsylla cheopis* (Siphonaptera: Pulicidae) Is as Efficient as Transmission by Blocked Fleas, *J Med Entomol* 44 (4) (2007) 678–682. doi:10.1093/jmedent/44.4.678.
- [10] B. J. Hinnebusch, D. M. Bland, C. F. Bosio, C. O. Jarrett, Comparative Ability of *Oropsylla montana* and *Xenopsylla cheopis* Fleas to Transmit *Yersinia pestis* by Two Different Mechanisms, *PLOS Neglected Tropical Diseases* 11 (1) (2017) e0005276. doi:10.1371/journal.pntd.0005276.
- [11] D. J. D. Earn, J. Ma, H. Poinar, J. Dushoff, B. M. Bolker, Acceleration of plague outbreaks in the second pandemic, *Proceedings of the National Academy of Sciences* 117 (44) (2020) 27703–27711. doi:10.1073/pnas.2004904117.
- [12] Plague, <https://www.who.int/news-room/fact-sheets/detail/plague>.
- [13] J. Mackay-Alderson, M. Quastel, E. Wilson, D. Belamy, Factors influencing the re-emergence of plague in Madagascar, *Emerging Topics in Life Sciences* 4 (2020) 423–433. doi:10.1042/ETLS20200334.
- [14] K. B. Patterson, T. Runge, Smallpox and the Native American, *The American Journal of the Medical Sciences* 323 (4) (2002) 216–222. doi:10.1097/00000441-200204000-00009.
- [15] J. S. Marr, J. T. Cathey, New Hypothesis for Cause of Epidemic among Native Americans, *New England, 1616–1619, Emerg Infect Dis* 16 (2) (2010) 281–286. doi:10.3201/edi1602.090276.
- [16] G. Legnaghi de Almeida, C. Santos, The microscopic settler: Smallpox dissemination in the context of european New World exploration *El colono microscópico: Diseminación de la viruela en el contexto de la exploración europea del Nuevo Mundo*, *Esbocos: histórias em contextos globais* 32 (2025) 1–22. doi:10.5007/2175-7976.2025.e101292.
- [17] S. Riedel, Smallpox and biological warfare: A disease revisited, *Proc (Bayl Univ Med Cent)* 18 (1) (2005) 13–20. doi:10.1080/08998280.2005.11928026.
- [18] N. Barquet, P. Domingo, Smallpox: The triumph over the most terrible of the ministers of death, *Ann Intern Med* 127 (8 Pt 1) (1997) 635–642. doi:10.7326/0003-4819-127-8_part_1-199710150-00010.
- [19] P. Vaishali, N. Boopathy, Edward Jenner: The Pioneer of Vaccination and His Enduring Legacy in Modern Medicine, *Cureus* 16 (Sep. 2024). doi:10.7759/cureus.68805.
- [20] Smallpox, <https://www.who.int/health-topics/smallpox>.
- [21] K. A. Simonsen, J. Snowden, Smallpox, in: *StatPearls*, StatPearls Publishing, Treasure Island (FL), 2025.
- [22] Louis Pasteur - Microbiology, Germ Theory, Pasteurization | Britannica, <https://www.britannica.com/biography/Louis-Pasteur/Spontaneous-generation> (Nov. 2025).
- [23] Louis Pasteur and germ theory - Medicine in 18th- and 19th-century Britain, c.1700-c.1900 - Edexcel - GCSE History Revision - Edexcel, <https://www.bbc.co.uk/bitesize/guides/zxbqjsg/revision/2>.
- [24] Pasteurization | Definition, Process, Inventor, & Facts | Britannica, <https://www.britannica.com/technology/pasteurization> (Nov. 2025).
- [25] K. A. Smith, Louis Pasteur, the Father of Immunology?, *Front Immunol* 3 (2012) 68. doi:10.3389/fimmu.2012.00068.

- [26] B. -. A. du texte Repiquet, Méthode de M. Pasteur pour prévenir la rage après morsure, par R. Repiquet, ..., 1891.
- [27] R. Rappuoli, 1885, the first rabies vaccination in humans, *Proceedings of the National Academy of Sciences* 111 (34) (2014) 12273–12273. doi:10.1073/pnas.1414226111.
- [28] R. Gaynes, The Discovery of Penicillin—New Insights After More Than 75 Years of Clinical Use, *Emerg Infect Dis* 23 (5) (2017) 849–853. doi:10.3201/eid2305.161556.
- [29] The discovery of antibiotics.
- [30] S. Y. Tan, Y. Tatsumura, Alexander Fleming (1881–1955): Discoverer of penicillin, *Singapore Med J* 56 (7) (2015) 366–367. doi:10.11622/smedj.2015105.
- [31] Sir Alexander Fleming – Nobel Lecture, <https://www.nobelprize.org/prizes/medicine/1945/fleming/lecture/>.
- [32] J. A. Segre, What does it take to satisfy Koch’s postulates two centuries later? Microbial genomics and Propionibacteria acnes, *J Invest Dermatol* 133 (9) (2013) 2141–2142. doi:10.1038/jid.2013.260.
- [33] L. Walker, H. LeVine, M. Jucker, Koch’s Postulates and Infectious Proteins, *Acta Neuropathol* 112 (1) (2006) 1–4. doi:10.1007/s00401-006-0072-x.
- [34] R. Koch, Die Ätiologie der Milzbrand-Krankheit, begründet auf die Entwicklungsgeschichte des Bacillus Anthracis (Mar. 2010). doi:10.25646/5064.
- [35] Ueber den augenblicklichen Stand der bakteriologischen Choleradiagnose, <https://zenodo.org/records/2077681>.
- [36] G. K. Gesundheitsamt, Mittheilungen aus dem Kaiserlichen Gesundheitsamte: Bd. Die Aetiologie der Tuberkulose, Norddeutsche Buchdruckerei und Verlagsanstalt ; zu beziehen durch Louis Gerschel Verlagsbuchhandlung (G. Gossmann), 1884.
- [37] J. S. Tyagi, The timeless legacy of Robert Koch: Founder of medical microbiology, *Reson* 11 (9) (2006) 20–28. doi:10.1007/BF02834330.
- [38] L. K. Charkoudian, J. T. Fitzgerald, C. Khosla, A. Champlin, In Living Color: Bacterial Pigments as an Untapped Resource in the Classroom and Beyond, *PLoS Biol* 8 (10) (2010) e1000510. doi:10.1371/journal.pbio.1000510.
- [39] S. Y. Choi, S. Lim, K.-h. Yoon, J. I. Lee, R. J. Mitchell, Biotechnological Activities and Applications of Bacterial Pigments Violacein and Prodigiosin, *J Biol Eng* 15 (2021) 10. doi:10.1186/s13036-021-00262-9.
- [40] H. S. Tuli, P. Chaudhary, V. Beniwal, A. K. Sharma, Microbial pigments as natural color sources: Current trends and future perspectives, *Journal of Food Science and Technology* 52 (8) (2014) 4669. doi:10.1007/s13197-014-1601-6.
- [41] T. Keerthana, S. S. Subhashini, Microbial Pigments: Types, Production Methods, and Diverse Applications in Various Industries: A Review, *IJSAT* 16 (1) (2025) 1593. doi:10.71097/IJSAT.v16.i1.1593.
- [42] M. Gomes, G. Soares, A Review of Bacterial Pigments: Harnessing Nature’s Colors for Functional Materials and Dyeing Processes, *Textile & Leather Review* 6 (2023) 167–190. doi:10.31881/TLR.2023.016.
- [43] L. J. Rather, S. Jameel, S. A. Ganie, K. A. Bhat, Lichen Derived Natural Colorants: History, Extraction, and Applications, in: *Handbook of Renewable Materials for Coloration and Finishing*, John Wiley & Sons, Ltd, 2018, Ch. 6, pp. 103–114. doi:10.1002/9781119407850.ch6.
- [44] M. E. Paillière-Jiménez, P. Stincone, A. Brandelli, Natural Pigments of Microbial Origin, *Front. Sustain. Food Syst.* 4 (Sep. 2020). doi:10.3389/fsufs.2020.590439.
- [45] C. K. Venil, P. Velmurugan, L. Dufossé, P. Renuka Devi, A. Veera Ravi, Fungal Pigments: Potential Coloring Compounds for Wide Ranging Applications in Textile Dyeing, *J Fungi (Basel)* 6 (2) (2020) 68. doi:10.3390/jof6020068.
- [46] L. Lin, J. Xu, Production of Fungal Pigments: Molecular Processes and Their Applications, *J Fungi (Basel)* 9 (1) (2022) 44. doi:10.3390/jof9010044.
- [47] B. Basto, N. R. da Silva, J. A. Teixeira, S. C. Silvério, B. Basto, N. R. da Silva, J. A. Teixeira, S. C. Silvério, Production of Natural Pigments by *Penicillium brevicompactum* Using Agro-Industrial Byproducts, *Fermentation* 8 (10) (Oct. 2022). doi:10.3390/fermentation8100536.
- [48] A. Kramar, M. M. Kostic, A. Kramar, M. M. Kostic, Bacterial Secondary Metabolites as Biopigments for Textile Dyeing, *Textiles* 2 (2) (2022) 252–264. doi:10.3390/textiles2020013.
- [49] W. Elkhateeb, M. O. Elnahas, G. Daba, Wide Range Applications of Fungal Pigments in Textile Dyeing, in: T. Satyanarayana, S. K. Deshmukh (Eds.), *Fungi and Fungal Products in Human Welfare and Biotechnology*, Springer Nature, Singapore, 2023, pp. 289–304. doi:10.1007/978-981-19-8853-0_10.
- [50] M. Yokoyama, What is Astro Microbiology?, <https://www.news-medical.net/life-sciences/What-is-Astro-Microbiology.aspx> (Oct. 2018).

- [51] G. Horneck, D. M. Klaus, R. L. Mancinelli, Space Microbiology, *Microbiology and Molecular Biology Reviews* 74 (1) (2010) 121–156. doi:10.1128/mmbr.00016-09.
- [52] R. Moeller, P. Setlow, G. Reitz, W. L. Nicholson, Roles of small, acid-soluble spore proteins and core water content in survival of *Bacillus subtilis* spores exposed to environmental solar UV radiation, *Appl Environ Microbiol* 75 (16) (2009) 5202–5208. doi:10.1128/AEM.00789-09.
- [53] G. Gallo, M. Aulitto, Advances in Extremophile Research: Biotechnological Applications through Isolation and Identification Techniques, *Life (Basel)* 14 (9) (2024) 1205. doi:10.3390/life14091205.
- [54] N. Merino, H. S. Aronson, D. P. Bojanova, J. Feyhl-Buska, M. L. Wong, S. Zhang, D. Giovannelli, Living at the Extremes: Extremophiles and the Limits of Life in a Planetary Context, *Front. Microbiol.* 10 (Apr. 2019). doi:10.3389/fmicb.2019.00780.
- [55] B. N. Rekadwad, W.-J. Li, J. M. Gonzalez, R. Punnappady Devasya, A. Ananthapadmanabha Bhagwath, R. Urana, K. Parwez, Extremophiles: The species that evolve and survive under hostile conditions, *3 Biotech* 13 (9) (2023) 316. doi:10.1007/s13205-023-03733-6.
- [56] T. Watson, 86 Percent of Earth's Species Still Unknown?, <https://www.nationalgeographic.com/science/article/110824-earths-species-8-7-million-biology-planet-animals-science> (Oct. 2025).
- [57] Microbiology | Definition, History, & Microorganisms | Britannica, <https://www.britannica.com/science/microbiology> (Dec. 2025).
- [58] E. Rabbow, P. Rettberg, S. Barczyk, M. Bohmeier, A. Parpart, C. Panitz, G. Horneck, R. von Heise-Rotenburg, T. Hoppenbrouwers, R. Willnecker, P. Baglioni, R. Demets, J. Dettmann, G. Reitz, EXPOSE-E: An ESA Astrobiology Mission 1.5 Years in Space, *Astrobiology* 12 (5) (2012) 374–386. doi:10.1089/ast.2011.0760.
- [59] M. Wassmann, R. Moeller, E. Rabbow, C. Panitz, G. Horneck, G. Reitz, T. Douki, J. Cadet, H. Stan-Lotter, C. S. Cockell, P. Rettberg, Survival of Spores of the UV-Resistant *Bacillus subtilis* Strain MW01 After Exposure to Low-Earth Orbit and Simulated Martian Conditions: Data from the Space Experiment ADAPT on EXPOSE-E, *Astrobiology* 12 (5) (2012) 498–507. doi:10.1089/ast.2011.0772.
- [60] Pasteur-Theorie für Lebensbeginn: Stock-Vektorgrafik (Lizenzfrei) 1182778669, <https://www.shutterstock.com/de/image-vector/pasteur-theory-life-beginning-1182778669>.
- [61] E. Frankel, J. Temple, E. Dikener, M. Berkmen, Bridging the gap with bacterial art, *FEMS Microbiol Lett* 370 (2023) fnad025. doi:10.1093/femsle/fnad025.
- [62] J. Madhusoodanan, Science and Culture: Petri palettes create microbial masterpieces, *Proc Natl Acad Sci U S A* 113 (40) (2016) 11056–11058. doi:10.1073/pnas.1610867113.
- [63] Y. Caro, M. Venkatachalam, J. Lebeau, M. Fouillaud, L. Dufossé, Pigments and Colorants from Filamentous Fungi, in: *Fungal Metabolites*, Springer, Cham, 2015, pp. 1–70. doi:10.1007/978-3-319-19456-1_26-1.
- [64] F. McDonagh, M. Cormican, D. Morris, L. Burke, N. K. Singh, K. Venkateswaran, G. Miliotis, Medical Astro-Microbiology: Current Role and Future Challenges, *J Indian Inst Sci* 103 (3) (2023) 771–796. doi:10.1007/s41745-023-00360-1.