

Assessment	Test	Domain	Skill	Difficulty
SAT	Reading and Writing	Information and Ideas	Command of Evidence	Medium

Question

Mean Ratings for Patients after 21 Days

Measure	Mean rating for participants aware of taking a placebo	Mean rating for participants in the control group
Global improvement	5.0	3.9
Symptom severity reduction	92.00	46.00
Quality of life improvement	11.4	5.4

To test whether a medication is effective, scientists compare outcomes for patients taking it and patients taking a placebo (a medically inactive substance). Patients normally aren’t told they’re receiving a placebo, but a research team conducted a study to see if there might be a medical benefit to telling them. The team used various measures to evaluate participants, with higher ratings indicating greater well-being in each measure. Compared to the mean ratings after 21 days for participants in the control group, the mean ratings for participants who were aware of taking a placebo _____

Which choice most effectively uses data from the table to complete the statement?

Answer

- A. ranged from 5.0 to 92.00, indicating that well-being varied widely from participant to participant.
- B. were lower for two measures, with the rating for only one measure indicating greater well-being for these participants.
- C. ranged from 3.9 to 46.00, with no rating indicating greater well-being in any measure for these participants.
- D. were higher for all three measures, indicating greater overall well-being for these participants.

Correct Answer: D

Rationale

Choice D is the best answer because it most effectively uses data from the table to complete the statement comparing the mean ratings for two different groups of participants in a study. The text explains that a research team evaluated the study’s participants using various measures in order to learn whether there might be a medical benefit to telling patients they’re receiving a placebo. The table shows the team’s mean ratings for participants after 21 days for three of the measures: global improvement, symptom severity reduction, and quality of life improvement. According to the table, the mean ratings were higher for all three measures for participants aware of taking a placebo than for participants in the control group. Given that higher ratings indicate greater well-being, as the text states, the mean ratings in the table indicate greater overall well-being for participants aware of taking a placebo than for participants in the control group.

Choice A is incorrect because the table doesn’t include data about individual participants; rather, it presents means, or mathematical averages, of ratings. For this reason, no conclusions can be drawn from data in the table about the extent to which well-being may have varied from participant to participant. Choice B is incorrect because according to the table, the mean ratings for participants aware of taking a placebo were higher for all three measures than for participants in the control group, not lower for two of the measures. Choice C is incorrect because it cites data from the table related to participants in the control group, not to participants aware of taking a placebo. Additionally, the mean ratings in the table for participants aware of taking a placebo are higher for all three measures than for participants in the control group. Given that higher ratings indicate greater well-being, as the text states, the ratings in the table for participants aware of taking a placebo indicate greater well-being for these participants in all three measures.