


Assessment	Test	Domain	Skill	Difficulty
SAT	Reading and Writing	Information and Ideas	Command of Evidence	Medium

Question



In high concentrations, hydrogen sulfide (H₂S) is typically toxic to many plants. Frederick D. Dooley and colleagues wanted to understand what effects low doses of H₂S might have on plant growth. They treated bean, corn, wheat, and pea seeds with various concentrations (measured in micromoles per liter) of H₂S and tracked the germination of those seeds along with the germination of untreated seeds. Treatment with particular concentrations of H₂S was associated with accelerated germination: for example, _____

Which choice most effectively uses data from the graph to complete the statement?

Answer

- A. at 24 hours, less than 10% of seeds treated with H₂S at a concentration of 10 micromoles per liter had germinated, whereas more than 90% of those seeds had germinated at 168 hours.
- B. at 48 hours, more than 70% of seeds treated with H₂S at a concentration of 10 micromoles per liter had germinated, whereas only approximately 50% of untreated seeds had germinated.
- C. at 168 hours, more than 90% of seeds treated with H₂S at concentrations of 10 or 500 micromoles per liter had germinated, whereas less than 70% of untreated seeds had germinated.
- D. at 48 hours, approximately 50% of seeds treated with H₂S at a concentration of 10 micromoles per liter had germinated, whereas only approximately 30% of untreated seeds had germinated.