

Rationale

Choice C is the best answer because it most logically completes the text based on supporting data in the graph. The text indicates that in the fly *D. melanogaster*, *DptA* and *DptB* are genes that encode peptides that both fight pathogens and promote beneficial microbes. Researchers tested *D. melanogaster*'s resistance to *P. rettgeri* and *A. sicerae* bacteria based on which variation of the peptide-encoding gene the flies exhibit: *DptA* silenced (referred to as type A), *DptB* silenced (type B), or both silenced (type AB). The text also indicates that resistance to *P. rettgeri* correlates with *DptA* activity but not with *DptB* activity (which would manifest as type B flies surviving at a higher rate than other fly types when exposed to *P. rettgeri*). The graph shows the post-*A. sicerae* infection results, which indicate that *DptB* activity was most strongly associated with survival, whereas *DptA* activity was not (manifesting in the graph as the type A flies having greater survival rates than the other fly types). In other words, when *DptA* activity was silenced, the flies showed relatively high survival rates, but when *DptB* activity was silenced, whether on its own or in conjunction with *DptA* activity being silenced, survival rates were low, suggesting that *DptB* may have developed as a specific defense against *A. sicerae*.

Choice A is incorrect. The graph suggests that *DptA* activity is associated with a low rate of survival, not a high one. Furthermore, the graph shows results for flies where *DptA* alone was silenced, *DptB* alone was silenced, and both were silenced and thus does not show any flies with activity in both *DptA* and *DptB*, which would be necessary to determine whether *DptA* conferred defense against *A. sicerae* in the presence of *DptB*. Choice B is incorrect. Only two bacteria species were considered in the text: *P. rettgeri* and *A. sicerae*. The text and graph taken together suggest that activity in *DptA* is associated with resistance to *P. rettgeri* while *DptB* activity is not, and that *DptB* activity is associated with resistance to *A. sicerae* while *DptA* is not. There is no further information to suggest one genetic type confers resistance to a greater number of pathogens than the other. Choice D is incorrect. The graph does not address flies with activity in both *DptA* and *DptB*. All flies represented in the graph had one or both of *DptA* and *DptB* silenced, or inactive.