# Unit 02 - Propositional Logic - Basics

Peter Bauer

### Outline

Propositions and Propositional Variable

Compositions of Propositions

Propositional Formulas

Properties of Compositions

Propositions are done in everyday life.

- Propositions are done in everyday life.
- ► For propositions it makes sense to ask whether they are "true" or "false".

- Propositions are done in everyday life.
- For propositions it makes sense to ask whether they are "true" or "false".
- ► The answer to this question is either "true" or "false" (nothing in between).

- Propositions are done in everyday life.
- For propositions it makes sense to ask whether they are "true" or "false".
- ► The answer to this question is either "true" or "false" (nothing in between).
- Example: "The circuit is electrically conducting."

► Instead of propositions we are often talking of *propositional* variables

- Instead of propositions we are often talking of propositional variables
- Expresses that we are not interested in a specific proposition but a variable that can hold arbitrary propositions

- Instead of propositions we are often talking of propositional variables
- Expresses that we are not interested in a specific proposition but a variable that can hold arbitrary propositions
- ► Simplified we can understand propositional variables as variables that hold either the value *t* or *f*

- Instead of propositions we are often talking of propositional variables
- Expresses that we are not interested in a specific proposition but a variable that can hold arbitrary propositions
- ► Simplified we can understand propositional variables as variables that hold either the value *t* or *f*
- ► Instead of "Let A be an arbitrary proposition" we can say "Let A be a propositional variable"

### Definition

A proposition is a "linguistic item" for which it makes sense to ask whether it is "true" or "false". The terms "true" and "false" are called the *truth values* of a proposition.

### **Definition**

A proposition is a "linguistic item" for which it makes sense to ask whether it is "true" or "false". The terms "true" and "false" are called the *truth values* of a proposition.

#### Remark

Propositions are mainly denoted by upper case letters A, B, C, ...

### Definition

A proposition is a "linguistic item" for which it makes sense to ask whether it is "true" or "false". The terms "true" and "false" are called the *truth values* of a proposition.

#### Remark

- Propositions are mainly denoted by upper case letters A, B, C, ...
- ▶ "true" is often abbreviated by t and "false" by f

### **Definition**

A proposition is a "linguistic item" for which it makes sense to ask whether it is "true" or "false". The terms "true" and "false" are called the *truth values* of a proposition.

#### Remark

- Propositions are mainly denoted by upper case letters A, B, C, ...
- ▶ "true" is often abbreviated by t and "false" by f
- We are only interested in the structure of propositions and not in its content.

# Examples

**▶** *x* ≤ 3

### **Examples**

- > x ≤ 3
- ▶ In German: Wenn Anastasia den Most holt, wird es bald etwas zu trinken geben, und das Abendessen wird beginnen, vorausgesetzt, dass Bartholomäus das Brot schon gebacken hat.

#### Definition

$$\begin{array}{c|c|c|c}
A & B & A \land B & A \lor B & \neg A \\
\hline
t & t & & \\
\end{array}$$

#### Definition

Α	В	$A \wedge B$	$A \vee B$	$\neg A$
t	t	t	t	f

#### Definition

A	В	$A \wedge B$	$A \vee B$	$\neg A$
t	t	t	t	f
t	f		'	

#### Definition

Α	В	$A \wedge B$	$A \vee B$	$\neg A$
t	t	t	t	f
t	f	f	t	f

#### Definition

À	В	$A \wedge B$	$A \vee B$	$\neg A$
t	t	t	t	f
t	f	f	t	f
f	t			•

#### Definition

A	В	$A \wedge B$	$A \vee B$	$\neg A$
t	t	t	t	f
t	f	f	t	f
f	t	f	t	t

#### Definition

Ă	В	$A \wedge B$	$A \vee B$	$\neg A$
t	t	t	t	f
t	f	f	t	f
f	t	f	t	t
f	f			

#### Definition

A	В	$A \wedge B$	$A \vee B$	$\neg A$
t	t	t	t	f
t	f	f	t	f
f	t	f	t	t
f	f	f	f	t

#### Definition

Let A and B be two propositions. Then the truth values of the propositions  $A \wedge B$  ("A and B"),  $A \vee B$  ("A or B), and  $\neg A$  ("not A") are given by the following truth table:

_ <i>A</i>	В	$A \wedge B$	$A \vee B$	$\neg A$
t	t	t	t	f
t	f	f	t	f
f	t	f	t	t
f	f	f	f	t

#### Remark

 $\wedge$  is called *conjunction*,  $\vee$  is called *disjunction* and  $\neg$  is called *negation*.

#### Definition

### Definition

$$\begin{array}{c|c}
A & B & A \Rightarrow B \\
\hline
t & t
\end{array}$$

### Definition

#### Definition

A	В	$A \Rightarrow B$
t	t	t
t	f	f
f	l t '	

### Definition

`A	В	$A \Rightarrow B$
t	t	t
t	f	f
f	t	t
f	f '	1

### Definition

A	В	$A \Rightarrow B$
t	t	t
t	f	f
f	t	t
f	f	t

### Definition

### Definition

$$\begin{array}{c|c}
A & B & A \Leftrightarrow B \\
\hline
t & t
\end{array}$$

#### Definition

`A	В	$A \Leftrightarrow B$
t	t	t
t	f	

### Definition

A	В	$A \Leftrightarrow B$
t	t	t
t	f	f
f	t '	

### Definition

A	В	$A \Leftrightarrow B$
t	t	t
t	f	f
f	t	f
f	f	1

### Equivalence

### Definition

Let A and B be two propositions. Then the equivalence  $A \Leftrightarrow B$  ("A if and only if B") is defined by the following truth table:

A	В	$A \Leftrightarrow B$
t	t	t
t	f	f
f	t	f
f	f	t

▶ The composition of two propositions is again a proposition

- ▶ The composition of two propositions is again a proposition
- ▶ The compositions are binary, i.e., they have two "operands"

- ▶ The composition of two propositions is again a proposition
- ▶ The compositions are binary, i.e., they have two "operands"
- By nesting compositions and using parentheses we can construct more complex propositions (more specific propositional formulas)

- ▶ The composition of two propositions is again a proposition
- ▶ The compositions are binary, i.e., they have two "operands"
- By nesting compositions and using parentheses we can construct more complex propositions (more specific propositional formulas)
- ▶ To reduce the number of necessary parentheses we use the precedence rules ¬ has higher precedence than ∧ higher than ∨ higher than ⇒ higher than ⇔

- The composition of two propositions is again a proposition
- The compositions are binary, i.e., they have two "operands"
- By nesting compositions and using parentheses we can construct more complex propositions (more specific propositional formulas)
- To reduce the number of necessary parentheses we use the precedence rules ¬ has higher precedence than ∧ higher than ∨ higher than ⇒ higher than ⇔
- ► The syntax of the compositions has to be considered, e.g.,  $\neg \land AB$  is NO proposition

## Propositional Formula

### Definition

Given a set of propositional variables or propositions  $A, B, C, \ldots$  If we combine these propositions by means of the compositions given above, the result of this compositions is called a *propositional* formula. If we assign true or false propositions to each of the propositional variables in a propositional formula we get a concrete proposition.

## Equivalence and Implication of Propositional Formulas

### Definition

Let P = P(A, B, C, ...) and Q = Q(A, B, C, ...) be propositional formulas then

1. P and Q are called equivalent ( $P \equiv Q$ ), if P and Q have the same truth value for each possible assignment of their propositional variables

## Equivalence and Implication of Propositional Formulas

### Definition

Let P = P(A, B, C, ...) and Q = Q(A, B, C, ...) be propositional formulas then

- 1. P and Q are called equivalent ( $P \equiv Q$ ), if P and Q have the same truth value for each possible assignment of their propositional variables
- 2. P implies Q ( $P \vdash Q$ ) if for each assignment to the propositional variables holds that if P is true then also Q is true.

### Equivalence and Implication of Propositional Formulas

#### Definition

Let P = P(A, B, C, ...) and Q = Q(A, B, C, ...) be propositional formulas then

- 1. P and Q are called equivalent ( $P \equiv Q$ ), if P and Q have the same truth value for each possible assignment of their propositional variables
- 2. P implies Q ( $P \vdash Q$ ) if for each assignment to the propositional variables holds that if P is true then also Q is true.

### Remark

Very often we loosely interchange the symbols  $\equiv$  and  $\Leftrightarrow$  as well as  $\vdash$  and  $\Rightarrow$ 

# Commutativity, Associativity

### Theorem

1. Commutativity

$$A \wedge B \equiv B \wedge A$$
  
 $A \vee B \equiv B \vee A$ 

# Commutativity, Associativity

### **Theorem**

1. Commutativity

$$A \wedge B \equiv B \wedge A$$
  
 $A \vee B \equiv B \vee A$ 

2. Associativity

$$A \wedge (B \wedge C) \equiv (A \wedge B) \wedge C \equiv A \wedge B \wedge C$$
  
 $A \vee (B \vee C) \equiv (A \vee B) \vee C \equiv A \vee B \vee C$ 

## Distributivity, De-Morgan

#### Theorem

1. Distributivity

$$A \wedge (B \vee C) \equiv (A \wedge B) \vee (A \wedge C)$$
  
 $A \vee (B \wedge C) \equiv (A \vee B) \wedge (A \vee C)$ 

# Distributivity, De-Morgan

#### **Theorem**

1. Distributivity

$$A \wedge (B \vee C) \equiv (A \wedge B) \vee (A \wedge C)$$
  
 $A \vee (B \wedge C) \equiv (A \vee B) \wedge (A \vee C)$ 

2. De-Morgan

$$\neg (A \land B) \equiv \neg A \lor \neg B$$
$$\neg (A \lor B) \equiv \neg A \land \neg B$$

# Absorption, Idempotence, and Double Negation

#### **Theorem**

1. Absorption

$$A \wedge (A \vee B) \equiv A$$
  
 $A \vee (A \wedge B) \equiv A$ 

# Absorption, Idempotence, and Double Negation

### Theorem

1. Absorption

$$A \wedge (A \vee B) \equiv A$$
  
 $A \vee (A \wedge B) \equiv A$ 

2. Idempotence

$$A \wedge A \equiv A$$
  
 $A \vee A \equiv A$ 

# Absorption, Idempotence, and Double Negation

### Theorem

1. Absorption

$$A \wedge (A \vee B) \equiv A$$
  
 $A \vee (A \wedge B) \equiv A$ 

2. Idempotence

$$A \wedge A \equiv A$$
  
 $A \vee A \equiv A$ 

3. Double Negation

$$\neg(\neg A) \equiv A$$

## Tautology and Contradiction

### Definition

A formula which is always "true", no matter which truth values are assigned to its variables is called a tautology (sign T). A formula which is always "false", no matter which truth values are assigned to its variables is called a contradiction (sign F).

## Tautology and Contradiction

### Definition

A formula which is always "true", no matter which truth values are assigned to its variables is called a tautology (sign T). A formula which is always "false", no matter which truth values are assigned to its variables is called a contradiction (sign F).

### Example

It is easy to show that  $(A \Rightarrow B) \Leftrightarrow (\neg A \lor B)$  is a tautology.

# Laws with Contradictions and Tautologies

### Theorem

For a proposition A holds:

$$A \wedge T \equiv A, \quad A \vee T \equiv T$$
  
 $A \wedge F \equiv F, \quad A \vee F \equiv A$   
 $A \wedge \neg A \equiv F, \quad A \vee \neg A \equiv T$