

# Unit 05 – Proofs

Peter Bauer

# Outline

Axioms

Direct Proof

Indirect Proof

Proof by Contradiction

# Axioms

## Definition

*Axioms* are basic propositions or propositional formulas which are true (hold) from the beginning. They are normally accepted without any proof.

## Example

- ▶ An equivalence relation  $R$  is reflexive, symmetric, and transitive.
- ▶ The composition  $\wedge$  is associative, commutative, and distributive.

# Proof Rules

Proof rules determine how you can make new true formulas from already existing true formulas.

# Direct Proof

- ▶ Step-by-step conclusion
- ▶  $(A \Rightarrow C) \wedge (C \Rightarrow B) \vdash (A \Rightarrow B)$

# Direct Proof

- ▶ Step-by-step conclusion
- ▶  $(A \Rightarrow C) \wedge (C \Rightarrow B) \vdash (A \Rightarrow B)$

## Example

1. If you own a longhair cat it will lose hairs

# Direct Proof

- ▶ Step-by-step conclusion
- ▶  $(A \Rightarrow C) \wedge (C \Rightarrow B) \vdash (A \Rightarrow B)$

## Example

1. If you own a longhair cat it will lose hairs
2. If somebody/something loses hairs, your couch will be (sooner or later) contaminated with hairs

# Direct Proof

- ▶ Step-by-step conclusion
- ▶  $(A \Rightarrow C) \wedge (C \Rightarrow B) \vdash (A \Rightarrow B)$

## Example

1. If you own a longhair cat it will lose hairs
2. If somebody/something loses hairs, your couch will be (sooner or later) contaminated with hairs
3. If you own a longhair cat your couch will be (sooner or later) contaminated with hairs



# Direct Proof

- ▶ Step-by-step conclusion
- ▶  $(A \Rightarrow C) \wedge (C \Rightarrow B) \vdash (A \Rightarrow B)$

## Example

1. If you own a longhair cat it will lose hairs
2. If somebody/something loses hairs, your couch will be (sooner or later) contaminated with hairs
3. If you own a longhair cat your couch will be (sooner or later) contaminated with hairs

## Remark

A simplified but famous version of the direct proof is the so-called *Modus Ponens*:  $A \wedge (A \Rightarrow B) \vdash B$

# Indirect Proof (Modus Tollens)

- ▶  $A \Rightarrow B$  has to be proven

# Indirect Proof (Modus Tollens)

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Sometimes this turns out to be hard or even impossible to be proven

# Indirect Proof (Modus Tollens)

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Sometimes this turns out to be hard or even impossible to be proven
- ▶ Therefore, we assume that  $\neg B$

# Indirect Proof (Modus Tollens)

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Sometimes this turns out to be hard or even impossible to be proven
- ▶ Therefore, we assume that  $\neg B$
- ▶ Then we try to derive  $\neg A$

# Indirect Proof (Modus Tollens)

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Sometimes this turns out to be hard or even impossible to be proven
- ▶ Therefore, we assume that  $\neg B$
- ▶ Then we try to derive  $\neg A$
- ▶  $A \Rightarrow B \equiv \neg B \Rightarrow \neg A$

# Indirect Proof (Modus Tollens)

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Sometimes this turns out to be hard or even impossible to be proven
- ▶ Therefore, we assume that  $\neg B$
- ▶ Then we try to derive  $\neg A$
- ▶  $A \Rightarrow B \equiv \neg B \Rightarrow \neg A$

## Example

- ▶ If it is the last school day of the year we get our final certificates.

# Indirect Proof (Modus Tollens)

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Sometimes this turns out to be hard or even impossible to be proven
- ▶ Therefore, we assume that  $\neg B$
- ▶ Then we try to derive  $\neg A$
- ▶  $A \Rightarrow B \equiv \neg B \Rightarrow \neg A$

## Example

- ▶ If it is the last school day of the year we get our final certificates.
- ▶ This is logically equivalent to



# Indirect Proof (Modus Tollens)

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Sometimes this turns out to be hard or even impossible to be proven
- ▶ Therefore, we assume that  $\neg B$
- ▶ Then we try to derive  $\neg A$
- ▶  $A \Rightarrow B \equiv \neg B \Rightarrow \neg A$

## Example

- ▶ If it is the last school day of the year we get our final certificates.
- ▶ This is logically equivalent to
- ▶ If we do not get our final certificates it can't be the last school day of the year.

# Indirect Proof (Modus Tollens)

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Sometimes this turns out to be hard or even impossible to be proven
- ▶ Therefore, we assume that  $\neg B$
- ▶ Then we try to derive  $\neg A$
- ▶  $A \Rightarrow B \equiv \neg B \Rightarrow \neg A$

## Example

- ▶ If it is the last school day of the year we get our final certificates.
- ▶ This is logically equivalent to
- ▶ If we do not get our final certificates it can't be the last school day of the year.

Prove the rule via truth table.

# Modus Tollens – A More Formal Example

## Example

- ▶  $a \cdot b$  is even  $\Rightarrow a$  is even or  $b$  is even.

# Modus Tollens – A More Formal Example

## Example

- ▶  $a \cdot b$  is even  $\Rightarrow a$  is even or  $b$  is even.
- ▶ Assume that  $a$  is odd and  $b$  is odd

# Modus Tollens – A More Formal Example

## Example

- ▶  $a \cdot b$  is even  $\Rightarrow a$  is even or  $b$  is even.
- ▶ Assume that  $a$  is odd and  $b$  is odd
- ▶ Show under this assumption that  $a \cdot b$  must be odd

# Proof by Contradiction

- ▶  $A \Rightarrow B$  has to be proven

# Proof by Contradiction

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Assume  $A$

# Proof by Contradiction

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Assume  $A$
- ▶ Assume  $\neg B$



# Proof by Contradiction

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Assume  $A$
- ▶ Assume  $\neg B$
- ▶ Then we try to proof a contradiction

# Proof by Contradiction

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Assume  $A$
- ▶ Assume  $\neg B$
- ▶ Then we try to proof a contradiction
- ▶  $A \Rightarrow B \equiv A \wedge \neg B \Rightarrow \mathbf{F}$

# Proof by Contradiction

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Assume  $A$
- ▶ Assume  $\neg B$
- ▶ Then we try to proof a contradiction
- ▶  $A \Rightarrow B \equiv A \wedge \neg B \Rightarrow \mathbf{F}$
- ▶ Prove this rule via a truth table

# Proof by Contradiction

- ▶  $A \Rightarrow B$  has to be proven
- ▶ Assume  $A$
- ▶ Assume  $\neg B$
- ▶ Then we try to proof a contradiction
- ▶  $A \Rightarrow B \equiv A \wedge \neg B \Rightarrow \mathbf{F}$
- ▶ Prove this rule via a truth table

## Example

- ▶ Prove that  $a^2$  is even  $\Rightarrow a$  is even
- ▶ Assume that  $a^2$  is even
- ▶ Assume for the sake of contradiction that  $a$  is odd
- ▶ Then you can conclude that  $a^2$  is odd, which is in contradiction to our first assumption

# Proof by Contradiction

## A Specific Variant

- ▶ Sometimes the proposition to be proven is **not** of the form  $A \Rightarrow B$
- ▶ Instead only a proposition  $S$  is to be proven
- ▶ This is a short form of  $K \Rightarrow S$ , where  $K$  is the already proven knowledge or the axioms about the underlying universe of discourse
- ▶ In this case it is to proof  $K \wedge \neg S \Rightarrow \mathbf{F}$
- ▶ In short: In order to proof  $S$  we assume  $\neg S$  and show that this leads to “absurdity”

# Proof by Contradiction

## A Specific Variant

### Example

- ▶  $\neg(\exists n \in \mathbb{N}, \forall i \in \mathbb{N} : i \leq n)$  aka "There is no greatest natural number"
- ▶ We assume  $\exists n \in \mathbb{N}, \forall i \in \mathbb{N} : i \leq n$  aka "There is a greatest natural number"
- ▶ Let's take a new natural number  $k = n + 1$  which exists since  $n \in \mathbb{N}$  and adding 1 to  $n$  is again a natural number (since  $\mathbb{N}$  is closed under  $+$ ).
- ▶ Now we have a new natural number  $k > n$  which contradicts our assumption
- ▶ Therefore the initial proposition is proven