

Predicate Logic

Peter Bauer

Outline

Terminology

Quantifiers

Motivation

- ▶ Propositional logic deals with propositions.

Motivation

- ▶ Propositional logic deals with propositions.
- ▶ Propositions were considered to be "atomic"

Motivation

- ▶ Propositional logic deals with propositions.
- ▶ Propositions were considered to be "atomic"
- ▶ To express propositions more concretely we need more sophisticated terms

Motivation

- ▶ Propositional logic deals with propositions.
- ▶ Propositions were considered to be "atomic"
- ▶ To express propositions more concretely we need more sophisticated terms
- ▶ Examples
 - ▶ "x is a prime number."
 - ▶ "4 is an even number."
 - ▶ "This shark is 20 ft. long."

Variable

Definition

A *variable* is a name for an arbitrary element out of a universe.

Variable

Definition

A *variable* is a name for an arbitrary element out of a universe.

Remark

- ▶ Variables denote places where values out of the universe can be seen

Variable

Definition

A *variable* is a name for an arbitrary element out of a universe.

Remark

- ▶ Variables denote places where values out of the universe can be seen
- ▶ Variables are assigned specific values out of the universe

Variable

Definition

A *variable* is a name for an arbitrary element out of a universe.

Remark

- ▶ Variables denote places where values out of the universe can be seen
- ▶ Variables are assigned specific values out of the universe
- ▶ Names of variables are combinations of characters, numbers and other symbols.

Constant

If we want to consider concrete elements of a universe we use constants.

Constant

If we want to consider concrete elements of a universe we use constants.

Definition

Constants are names of concrete objects. These can be concrete values (e.g., π , 17, 42), function constants (e.g., $+$, \cos , $\sqrt{}$) or predicate constants (e.g., \geq , \in).

Term

Definition

Constants and variables are *terms*. Syntactic structures of the form $f(t_1, t_2, \dots, t_n)$ are *terms*, if f is a function constant with arity n and t_1, t_2, \dots, t_n are terms.

Term

Definition

Constants and variables are *terms*. Syntactic structures of the form $f(t_1, t_2, \dots, t_n)$ are *terms*, if f is a function constant with arity n and t_1, t_2, \dots, t_n are terms.

Example

- ▶ $\sin(x)$, where \sin is the function constant with arity 1

Term

Definition

Constants and variables are *terms*. Syntactic structures of the form $f(t_1, t_2, \dots, t_n)$ are *terms*, if f is a function constant with arity n and t_1, t_2, \dots, t_n are terms.

Example

- ▶ $\sin(x)$, where \sin is the function constant with arity 1
- ▶ $x + y$, where $+$ is the function constant with arity 2

Term

Definition

Constants and variables are *terms*. Syntactic structures of the form $f(t_1, t_2, \dots, t_n)$ are *terms*, if f is a function constant with arity n and t_1, t_2, \dots, t_n are terms.

Example

- ▶ $\sin(x)$, where \sin is the function constant with arity 1
- ▶ $x + y$, where $+$ is the function constant with arity 2
- ▶ \sqrt{y} , where $\sqrt{}$ is the function constant with arity 1

Term

Definition

Constants and variables are *terms*. Syntactic structures of the form $f(t_1, t_2, \dots, t_n)$ are *terms*, if f is a function constant with arity n and t_1, t_2, \dots, t_n are terms.

Example

- ▶ $\sin(x)$, where \sin is the function constant with arity 1
- ▶ $x + y$, where $+$ is the function constant with arity 2
- ▶ \sqrt{y} , where $\sqrt{}$ is the function constant with arity 1

Remark

Constant values like π , 17, or “Vergissmeinnicht” can be interpreted as terms. These are terms with a functions constant of arity 0.

Predicate and Formula

Definition

Phrases which describe properties of objects or relations between objects are called *predicate constants* or sometimes *predicates*.

Predicate and Formula

Definition

Phrases which describe properties of objects or relations between objects are called *predicate constants* or sometimes *predicates*.

Example

“is prime”, “isValidInput”, \geq , $=$ are predicate constants.

Predicate and Formula

Definition

Phrases which describe properties of objects or relations between objects are called *predicate constants* or sometimes *predicates*.

Example

“is prime”, “isValidInput”, \geq , $=$ are predicate constants.

Definition

Syntactic structures of the form $P(t_1, t_2, \dots, t_n)$ are *atomic formulas*, if P is a predicate constant with arity n and t_1, t_2, \dots, t_n are terms.

Some Examples and Conventions

Remark

Instead of the bloomy phrases of everyday life language, we use very slim and symbolic structures in mathematical logic

Some Examples and Conventions

Remark

Instead of the bloomy phrases of everyday life language, we use very slim and symbolic structures in mathematical logic

Example

everyday language	logic
"y is at least as great as z"	$y \geq z$
"a leads project b"	$\text{ProjectLead}(a, b)$

Some Examples and Conventions

Remark

Instead of the bloomy phrases of everyday life language, we use very slim and symbolic structures in mathematical logic

Example

everyday language	logic
"y is at least as great as z"	$y \geq z$
"a leads project b"	$\text{ProjectLead}(a, b)$

Remark

- ▶ The symbols a , b , y , and z are variables, " \geq " and "ProjectLead" are predicate constants.

Some Examples and Conventions

Remark

Instead of the bloomy phrases of everyday life language, we use very slim and symbolic structures in mathematical logic

Example

everyday language	logic
"y is at least as great as z"	$y \geq z$
"a leads project b"	$\text{ProjectLead}(a, b)$

Remark

- ▶ The symbols a, b, y , and z are variables, " \geq " and "ProjectLead" are predicate constants.
- ▶ In the sequel, we will use upper case letters like P, Q, R, \dots to denote predicate constants.

Some Examples and Conventions

Remark

Instead of the bloomy phrases of everyday life language, we use very slim and symbolic structures in mathematical logic

Example

everyday language	logic
"y is at least as great as z"	$y \geq z$
"a leads project b"	$\text{ProjectLead}(a, b)$

Remark

- ▶ The symbols a, b, y , and z are variables, " \geq " and "ProjectLead" are predicate constants.
- ▶ In the sequel, we will use upper case letters like P, Q, R, \dots to denote predicate constants.
- ▶ P, Q, R, \dots are always predicate constants and are *not* predicate variables (first order predicate logic)

Another Example

Combining predicate formulas by means of compositions as given in propositional logic lead again to formulas of predicate logic.

Another Example

Combining predicate formulas by means of compositions as given in propositional logic lead again to formulas of predicate logic.

Example

- Let x be a student of the HTL Leonding.

Another Example

Combining predicate formulas by means of compositions as given in propositional logic lead again to formulas of predicate logic.

Example

- ▶ Let x be a student of the HTL Leonding.
- ▶ Let $P(x) = \text{"}x \text{ studies LOAL"}$ and $Q(x) = \text{"}x \text{ is a first grader"}$.

Another Example

Combining predicate formulas by means of compositions as given in propositional logic lead again to formulas of predicate logic.

Example

- ▶ Let x be a student of the HTL Leonding.
- ▶ Let $P(x) = \text{"}x \text{ studies LOAL"}$ and $Q(x) = \text{"}x \text{ is a first grader"}$.
- ▶ Then the conjunction $P(x) \wedge Q(x)$ is again a formula of predicate logic.

Another Example

Combining predicate formulas by means of compositions as given in propositional logic lead again to formulas of predicate logic.

Example

- ▶ Let x be a student of the HTL Leonding.
- ▶ Let $P(x) = \text{"}x \text{ studies LOAL"}$ and $Q(x) = \text{"}x \text{ is a first grader"}$.
- ▶ Then the conjunction $P(x) \wedge Q(x)$ is again a formula of predicate logic.
- ▶ $P(x) \wedge Q(x) = \text{"}x \text{ studies LOAL and is a first grader"}$.

Quantifier — Motivation

- ▶ The truth value of atomic formulas can only be determined if we assign constants to each variable.
- ▶ It can't be determined in general, in particular, if the formula contains free variables.

Quantifier — Motivation

- ▶ The truth value of atomic formulas can only be determined if we assign constants to each variable.
- ▶ It can't be determined in general, in particular, if the formula contains free variables.

Example

- ▶ Consider the binary predicate constant \geq over \mathbb{N} .

Quantifier — Motivation

- ▶ The truth value of atomic formulas can only be determined if we assign constants to each variable.
- ▶ It can't be determined in general, in particular, if the formula contains free variables.

Example

- ▶ Consider the binary predicate constant \geq over \mathbb{N} .
- ▶ Consider further the formula $a \geq 3$.

Quantifier — Motivation

- ▶ The truth value of atomic formulas can only be determined if we assign constants to each variable.
- ▶ It can't be determined in general, in particular, if the formula contains free variables.

Example

- ▶ Consider the binary predicate constant \geq over \mathbb{N} .
- ▶ Consider further the formula $a \geq 3$.
- ▶ a is a free variable and 3 is a constant.

Quantifier — Motivation

- ▶ The truth value of atomic formulas can only be determined if we assign constants to each variable.
- ▶ It can't be determined in general, in particular, if the formula contains free variables.

Example

- ▶ Consider the binary predicate constant \geq over \mathbb{N} .
- ▶ Consider further the formula $a \geq 3$.
- ▶ a is a free variable and 3 is a constant.
- ▶ Now consider the formula $7 \geq 3$.

Quantifier — Motivation

- ▶ The truth value of atomic formulas can only be determined if we assign constants to each variable.
- ▶ It can't be determined in general, in particular, if the formula contains free variables.

Example

- ▶ Consider the binary predicate constant \geq over \mathbb{N} .
- ▶ Consider further the formula $a \geq 3$.
- ▶ a is a free variable and 3 is a constant.
- ▶ Now consider the formula $7 \geq 3$.
- ▶ Furthermore, consider the formula “for all $a \in \mathbb{N}$ holds: $a \geq 3$ ”

Quantifier

- ▶ The last example has shown that we need phrases like “for all variables ...” or “there exists ...”.

Quantifier

- ▶ The last example has shown that we need phrases like “for all variables ...” or “there exists ...”.
- ▶ For this purpose we have *Quantifiers* in mathematical logic.

Quantifier

- ▶ The last example has shown that we need phrases like “for all variables ...” or “there exists ...”.
- ▶ For this purpose we have *Quantifiers* in mathematical logic.
- ▶ Quantifiers bind variables in formulas of predicate logic.

Quantifier — Definition

Definition

The most important quantifiers are the *universal quantifier* and the *existential quantifier* which are written and spoken as follows:

1. Universal Quantifier \forall

► $\forall x : P(x)$

Quantifier — Definition

Definition

The most important quantifiers are the *universal quantifier* and the *existential quantifier* which are written and spoken as follows:

1. Universal Quantifier \forall

- ▶ $\forall x : P(x)$
- ▶ for all x holds P

Quantifier — Definition

Definition

The most important quantifiers are the *universal quantifier* and the *existential quantifier* which are written and spoken as follows:

1. Universal Quantifier \forall

- ▶ $\forall x : P(x)$
- ▶ for all x holds P

2. Existential Quantifier \exists

- ▶ $\exists x : P(x)$

Quantifier — Definition

Definition

The most important quantifiers are the *universal quantifier* and the *existential quantifier* which are written and spoken as follows:

1. Universal Quantifier \forall

- ▶ $\forall x : P(x)$
- ▶ for all x holds P

2. Existential Quantifier \exists

- ▶ $\exists x : P(x)$
- ▶ There exists (at least one) x for which holds P

Quantifier — Definition

Definition

The most important quantifiers are the *universal quantifier* and the *existential quantifier* which are written and spoken as follows:

1. Universal Quantifier \forall

- ▶ $\forall x : P(x)$
- ▶ for all x holds P

2. Existential Quantifier \exists

- ▶ $\exists x : P(x)$
- ▶ There exists (at least one) x for which holds P

Remark

In the sequel we will use the notations $\forall x : P(x)$ and $\exists x : P(x)$.

Free and Bound Variables

- ▶ Variables which are referred to by a quantifier are called *bound variables*.
- ▶ All other variables are *free variables* unless they are bound by another quantifier.

Free and Bound Variables

- ▶ Variables which are referred to by a quantifier are called *bound variables*.
- ▶ All other variables are *free variables* unless they are bound by another quantifier.

Example

- ▶ Consider the formula “ x has birthday in the same month as y ”.
- ▶ Let x and y denote students of the HTL Leonding.
- ▶ x and y are both free variables.

Free and Bound Variables

- ▶ Variables which are referred to by a quantifier are called *bound variables*.
- ▶ All other variables are *free variables* unless they are bound by another quantifier.

Example

- ▶ Consider the formula “ x has birthday in the same month as y ”.
- ▶ Let x and y denote students of the HTL Leonding.
- ▶ x and y are both free variables.
- ▶ $\exists x : x$ has birthday in the same month as y .
- ▶ x is a bound variable and y is a free variable.

Quantified Formulas

1. The formula $\forall x : P(x)$ is true if and only if P is true for every value of the universe that is assigned to x .

Quantified Formulas

1. The formula $\forall x : P(x)$ is true if and only if P is true for every value of the universe that is assigned to x .
2. The formula $\exists x : P(x)$ is true if and only if P is true for at least one value of the universe that is assigned to x .

Quantified Formulas

1. The formula $\forall x : P(x)$ is true if and only if P is true for every value of the universe that is assigned to x .
2. The formula $\exists x : P(x)$ is true if and only if P is true for at least one value of the universe that is assigned to x .
3. The formulas $\forall x : P(x, y, z)$ and $\exists x : P(x, y, z)$ are formulas with free variables y and z . Therefore, no truth value can be determined for these formulas.

Quantified Formulas

1. The formula $\forall x : P(x)$ is true if and only if P is true for every value of the universe that is assigned to x .
2. The formula $\exists x : P(x)$ is true if and only if P is true for at least one value of the universe that is assigned to x .
3. The formulas $\forall x : P(x, y, z)$ and $\exists x : P(x, y, z)$ are formulas with free variables y and z . Therefore, no truth value can be determined for these formulas.

Remark

- Since we can determine a truth value for the first and the second formula, we can call these formulas propositions.

Quantified Formulas

1. The formula $\forall x : P(x)$ is true if and only if P is true for every value of the universe that is assigned to x .
2. The formula $\exists x : P(x)$ is true if and only if P is true for at least one value of the universe that is assigned to x .
3. The formulas $\forall x : P(x, y, z)$ and $\exists x : P(x, y, z)$ are formulas with free variables y and z . Therefore, no truth value can be determined for these formulas.

Remark

- ▶ Since we can determine a truth value for the first and the second formula, we can call these formulas propositions.
- ▶ The third remains a formula of predicate logic.

Example

Example

- ▶ Let's consider the universe of all students of the HTL Leonding.
- ▶ Further consider the formula “ x attends the 3rd grade”.

Example

Example

- ▶ Let's consider the universe of all students of the HTL Leonding.
- ▶ Further consider the formula “ x attends the 3rd grade”.
- ▶ This formula has one free variable x .

Example

Example

- ▶ Let's consider the universe of all students of the HTL Leonding.
- ▶ Further consider the formula “ x attends the 3rd grade”.
- ▶ This formula has one free variable x .
- ▶ “ $\forall x : x$ attends the 3rd grade” would only be true if all students of this school would attend the 3rd grade.
- ▶ “ $\exists x : x$ attends the 3rd grade” is true, since there exists at least one student at this school who attends the 3rd grade.

Summary

We know two possibilities to turn a formula of predicate logic into a proposition

1. Assignment of constants out of the universe to each free variable.
2. Binding the free variables with quantifiers.

Summary

We know two possibilities to turn a formula of predicate logic into a proposition

1. Assignment of constants out of the universe to each free variable.
2. Binding the free variables with quantifiers.
3. Formulas with no free variables are propositions and are sometimes called *closed formulas*.
4. Only these have a truth value.

Relationships Between Negation and Quantifiers

Theorem

Let P be a unary predicate constant and $x \in D$ a variable. Then the following relations hold:

$$\neg \exists x : P(x) \equiv \forall x : \neg P(x)$$

$$\neg \forall x : P(x) \equiv \exists x : \neg P(x)$$

Relationships Between Negation and Quantifiers

Theorem

Let P be a unary predicate constant and $x \in D$ a variable. Then the following relations hold:

$$\neg \exists x : P(x) \equiv \forall x : \neg P(x)$$

$$\neg \forall x : P(x) \equiv \exists x : \neg P(x)$$

Example

$$\neg \forall x \in \mathbb{N} : \text{isPrime}(x) \Rightarrow \text{odd}(x)$$

$$\exists x \in \mathbb{N} : \text{isPrime}(x) \wedge \neg \text{odd}(x)$$