Forecast Cosmological Constraints with the 1D Wavelet Scattering Transform and the Lyman-alpha forest

Hurum Maksora Tohfa^{1,2},* Simeon Bird¹,† Ming-Feng Ho¹, Mahdi Qezlou¹, and Martin Fernandez^{1,3}

¹Department of Physics & Astronomy, University of California Riverside, Riverside, CA 92521, United States

²Department of Astronomy, University of Washington, Seattle, WA 98195, United States

³Department of Atmospheric Science, Colorado State University, Fort Collins, CO, United States

(Dated: June 5, 2024)

We make forecasts for the constraining power of the 1D Wavelet Scattering Transform (WST) when used with a Lyman- α forest cosmology survey. Using mock simulations and a Fisher matrix, we show that there is considerable cosmological information in the scattering transform coefficients not captured by the flux power spectrum. We estimate mock covariance matrices assuming uncorrelated Gaussian pixel noise for each quasar, at a level drawn from a simple lognormal model. The extra information comes from a smaller estimated covariance in the first-order wavelet power, and from second-order wavelet coefficients which probe non-Gaussian information in the forest. Forecast constraints on cosmological parameters from the WST are more than an order of magnitude tighter than for the power spectrum, shrinking a 4D parameter space by a factor of 10^6 . Should these improvements be realised with DESI, inflationary running would be constrained to test common inflationary models predicting $\alpha_s = -6 \times 10^{-4}$ and neutrino mass constraints would be improved enough for a $5-\sigma$ detection of the minimal neutrino mass.

INTRODUCTION

An important probe of small-scale structures is the Lyman- α forest, absorption features of neutral hydrogen. The Lyman- α forest allows us to explore fundamental questions about the Universe, such as the nature of dark matter, the total neutrino mass, and the physics of the inflationary era [1-4], as well as measure the thermal history of the intergalactic medium (IGM) [5]. However, as the Lyman- α forest probes the nonlinear regime, hydrodynamic simulations are needed to accurately predict its behavior under different cosmological parameters. There are a variety of Lyman- α forest surveys [e.g. 6–9], with the largest statistical sample being from the extended Baryon Oscillation Spectroscopic Survey (eBOSS) [10]. The ongoing Dark Energy Spectroscopic Instrument (DESI)[11, 12], and WEAVE-QSO [13] will vastly increase the number of spectra. However, SDSS and DESI will be systematics dominated in the flux power spectrum, suggesting a need for new summary statistics, especially those which can extract non-Gaussian information from the non-linear density field.

Higher order n-point functions such as the bispectrum have been calculated for galaxy and weak lensing surveys [14–19]. However, they suffer from increased variance and reduced robustness to outliers [20]. In highly non-Gaussian distributions with heavy tails, there can be significant fluctuations or noise in the data that make it difficult to extract meaningful features related to interactions/correlations. These fluctuations become more pronounced as we move toward higher orders and dilute important features [21].

Here we investigate using the wavelet scattering transform for Lyman- α forest cosmology. The wavelet scattering transform (WST) was initially proposed by Mallat for

signal processing [22]. In recent years, it has been used successfully in several signal processing fields [23, 24]. Each order of the WST is fully deterministic, mathematically robust, has clearly interpretable structures, and does not require any training of the model [25, 26]. The WST convolves signals with a complex-valued wavelet at different scales, selecting Fourier modes around a central frequency and thus separating information from different scales. This first order WST is conceptually similar to the power spectrum, but may be applied repeatedly to extract higher order summary statistics.

The WST has been applied to weak lensing shear maps, where it has been shown to outperform the power spectrum and achieve similar constraining power to neural network techniques [27]. While forecasts suggest that Convolutional Neural Networks (CNNs) can improve cosmological inference over two-point statistics, particularly for weak lensing data [28], other forecasts have suggested that scattering architectures are able to outperform a CNN [29], and are more interpretable [30]. The WST has also been applied to mock 21-cm data, where it forecast tighter constraints than the 3D spherically averaged power spectrum [31]. Finally, 3D WST has been shown to preserve 50% more of the information in an idealised mock density field than the marked power spectrum [32].

The wavelet power has been used to extract thermal history information from the Lyman- α forest by Ref. [33], although constraints are similar to those from the flux power spectrum [34]. We make forecasts for an extended wavelet-based analysis which both considers cosmological parameters and uses higher order WST coefficients to extract non-Gaussian information. We use the Fisher information matrix to make forecast constraints from a mock survey for both the WST and the power spectrum, finding that the WST substantially improves constrain-

ing power. Our analysis suggests a novel approach for extracting cosmological information from the Lyman- α forest.

MOCK DATA FOR ANALYSIS

We use 10 hydrodynamic simulations, described fully in Ref. [35]. Since this paper is aimed at making forecasts for the constraining power of different summary statistics rather than making a comparison to observational data, we use relatively fast, small simulations containing 2×256^3 dark matter and Smoothed-Particle Hydrodynamics particles within a 40 Mpc/h box.

Fully hydrodynamic simulations were performed with the simulation code MP-Gadget [36]. We generate 32 000 spectra Lyman- α absorption spectra from each simulation at z=4,3,2 with "fake_spectra" [37]. Spectra are generated parallel to the x-axis of the simulation box. We approximate a realistic survey by adding Gaussian noise and smoothing with a Gaussian filter with width corresponding to the spectral resolution of eBOSS.

Our simulation suite includes 4 parameters: the spectral index, n_P , and amplitude, A_P of the primordial power spectrum. The power spectrum is given by the equation:

$$P(k) = A_{\rm P} \left(\frac{k}{k_P}\right)^{n_P - 1},\tag{1}$$

where $k_P = 2\pi/8 = 0.78 \; \mathrm{Mpc^{-1}}$. We include parameters to model uncertainty in the thermal history of the IGM [38]. We rescale the photo-heating rate by a density-dependent factor, $\tilde{\epsilon} = H_A \epsilon \Delta^{H_S}$. ϵ is the photo-heating rate, Δ is the overdensity of the IGM, H_A controls the IGM temperature at mean density, and H_S controls the slope of the temperature-density relation.

We add an extra parameter in post-processing for the observed mean flux in the forest, which is proportional to the overall ionization fraction of neutral hydrogen. We rescale our spectra to have the same mean flux by multiplying the optical depth in each spectral pixel by a constant factor. The mean optical depth follows the power law redshift evolution from Ref. [39]:

$$\tau = 2.3 \times 10^{-3} (1+z)^{3.65}.$$
 (2)

THE 1D WAVELET SCATTERING TRANSFORM

We represent a spectrum by its input flux field $\mathcal{F}(x)$, a 1D array with velocity coordinate x. The 1D flux power spectrum is

$$P(k) = \left\langle |\widehat{\mathcal{F}}(x)|^2 \right\rangle = \left\langle |\mathcal{F}(x) \star \psi_k(x)|^2 \right\rangle.$$
 (3)

where $\psi_k = e^{-ikx}$ and \star is a convolution. We average over all spectra in the box, denoted by $\langle \rangle$.

We define the WST coefficients via recursive convolution of the field $\mathcal{F}(x)$ with a series of wavelets with different numbers of octaves (denoted by Q) of different scales (j). Each wavelet in the wavelet family maintains an identical shape but varies in scale and orientation. After convolution, we take the modulus of the generated fields and use low pass filters to smooth out the high-frequency components of the signal. We set Q=1 to employ a dilation factor of 2, making the wavelet's real-space size roughly equivalent to 2^J pixels [26], where J denotes the largest physical scale possible. By averaging the resultant fields, we obtain scattering coefficients that describe the statistical characteristics of the input flux. The zeroth, first, and second order scattering coefficients are defined as:

$$S_0 = \langle \mathcal{F}(x) \star \phi[0] \rangle \tag{4}$$

$$S_1(j_1) = \langle |\mathcal{F}(x) \star \psi^{j_1}| \star \phi[1] \rangle \tag{5}$$

$$S_2(j_1, j_2) = \langle || \mathcal{F}(x) \star \psi^{j_1} | \star \psi^{j_2} | \star \phi[2] \rangle \tag{6}$$

Here ψ^j are Morlet wavelet filters and $\phi[n]$ are low-pass filters. We compute the coefficients for $0 \leq j \leq J$, take their modulus, and average over all sightlines, as denoted by $\langle \rangle$. Lower 'j' values correspond to smaller scales, meaning they oscillate more slowly and capture more detailed small-scale structures. Higher-order scattering coefficients are computationally expensive [40], so we limit ourselves to scattering coefficients up to the second order. We set J=5, as we found that this extracted most of the information from our simulations, .

The zeroth-order coefficient is acquired by applying a low-pass filter, $\phi[J]$ to the input field, $\mathcal{F}(x)$. It thus represents the local average of the field, which, for the Lyman- α spectra, corresponds to the local intensity of the transmitted flux. We expect this to be highly degenerate with the mean flux and continuum fitting and so to be conservative we focus on the first and second order coefficients.

We use the publicly available KYMATIO package for generating 1D scattering coefficients [41].

The WST coefficients are Lipschitz continuous to deformation, meaning that similar fields differing by small deformations are also similar in terms of their scattering coefficient representation [26]. This makes the scattering characterization stable even when there are slight variations or noise in the data.

FISHER MATRIX FORMALISM

We use the Fisher information approach to forecast the parameter constraints that we could achieve with a WST analysis. The Fisher matrix is defined as the second derivative of the log-likelihood function, $\ln \mathcal{L}(p|d, M)$




FIG. 1. Dependence of the scattering coefficients described in Equation. 4 on changes in the mean flux at z=2. The top panels show the first and second order scattering coefficients and the bottom panels show the fractional change in mean flux with respect to the observed model. We see about a 2.5% change in scattering coefficients as the value of the mean flux changes by 10%.




FIG. 2. Change in the wavelet scattering coefficients defined in Equation 4 as we change our two cosmological parameters: n_P , A_P at z= 2. We show the fractional change in the cosmological parameters with respect to our fiducial model, similar to Fig. 1. Our fiducial cosmology is defined as $n_P = 0.897$, $A_P = 1.9 \times 10^{-9}$, $H_S = -0.3$, $H_A = 0.9$.

around the maximum likelihood location, where p is the parameter value of a model, and d is the data. Under this definition, we define the Fisher matrix as:

$$F_{ij} = \left\langle \frac{\partial^2 \ln \mathcal{L}}{\partial p_i^2} \right\rangle = \frac{\partial \mathbf{S}}{\partial p_i} \cdot \Sigma^{-1} \cdot \frac{\partial \mathbf{S}}{\partial p_j}.$$
 (7)

Following Refs. [42, 43], we model the covariance Σ by adding Gaussian noise to the sightlines from our fiducial simulation. The continuum-to-noise ratio (CNR) for each simulated spectrum is sampled from a log-normal distribution, whose parameters are chosen to fit to the

noise in the DR16Q observed spectral sample ¹[44]:

$$\log(\text{CNR}) \sim \mathcal{N}(\mu = 0.53, \sigma = 0.36) \tag{8}$$

Each spectrum, i has a CNR value, CNR_i, drawn from the distribution in Eq. 8. Gaussian noise, ϵ_i , realising CNR_i is then added to each pixel:

$$\mathcal{F}(x)_{j}' = \mathcal{F}(x)_{j} + \epsilon_{i}, \tag{9}$$

$$\epsilon_i \sim \mathcal{N}(\mu = 0, \sigma^2 = \text{CNR}_i). \tag{10}$$

In this work, we do not consider the errors from incomplete QSO continuum fitting as they are subdominant to

https://live-sdss4org-dr16.pantheonsite.io/algorithms/ qso_catalog

the CNR [42]. A comprehensive treatment of continuum errors is deferred to future work.

We generate the covariance matrix Σ by taking the variance of the scattering coefficients (or power spectrum), defined as:

$$\Sigma_{jk} = \frac{1}{N-1} \sum_{i=1}^{N} (X_{ij} - \bar{X}_j) (X_{ik} - \bar{X}_k) . \tag{11}$$

Here X is the summary statistic, either the WST coefficients or the 1D flux power spectrum. We use 2000 random noise realizations.

RESULTS

In this Section, we describe the sensitivity of the WST and the 1D flux power spectrum to the parameters A_P , n_P , H_S and H_A , as well as the mean flux $\tau_{\rm eff}$. The scattering transform with J=5 generates a total of 19 scattering coefficients (1 zeroth order, 6 first order, and 12 second-order coefficients). We do not include the zeroth order coefficient in our analysis.

Figure 1 shows the sensitivity of WST to changes in the mean flux. We computed the first and second order scattering transforms with a mean flux changed by 10% in post-processing. The top panel shows the scattering coefficients and the bottom panel shows fractional change in the WST coefficients. A 10% change in $\tau_{\rm eff}$ changes the first and second-order WST coefficients by $\sim 2.5\%$. For comparison, the 1D flux power spectrum is more sensitive to the mean flux, changing by about 7%.

Figure 2 show how the WST coefficients are affected as we change cosmological parameters. As in Figure 1, the top panel shows the scattering coefficients and the bottom panel shows the fractional change in the WST coefficients with respect to the fiducial model. We see that changes in cosmological parameters indeed affect the scattering coefficients, signifying the presence of cosmological information. Next, to compare the performance of WST with the power spectrum, we calculated the Fisher matrix. Figure 3 shows the Fisher information in the WST coefficients and the power spectrum, for our four parameters n_P , A_P , H_A , H_S . The second-order fields contribute about 50% extra information to the Fisher matrix over the first order fields. However, even the first order coefficients outperform the flux power spectrum.

We can invert the Fisher matrix to estimate posterior uncertainties. For a parameter X, $\sigma_X = 1/\sqrt{F_{ij}}$. Table I shows σ_X for the power spectrum and combined WST coefficients. The WST improves σ_X by a factor of 30-60.

These constraints are generated using a Fisher matrix rather than a full likelihood calculation. On the other hand, our Fisher matrix estimates neglect parameter degeneracies. The pivot scale k_P is chosen to minimize

Parameter	Power Spectrum	WST
Δn_P	0.178	0.0059
$\Delta A_P/10^{-9}$	0.503	0.0079
ΔH_S	0.207	0.0049
ΔH_A	0.258	0.0062

TABLE I. Estimated posterior uncertainties in parameters constraints obtained from both the power spectrum and the combined WST coefficients.

correlation between A_P and n_P , but there is some correlation between the cosmological and thermal parameters, visible in Figure 8 of Ref. [45].

Figure 3 also shows the redshift dependence of both summary statistics. The constraining power of both the WST coefficients and the flux power spectrum varies similarly with redshift. We find that redshift 3 contains the most information in both WST and the flux power spectrum for all parameters except H_A , which controls the mean IGM temperature and is most constrained by z=2.

CONCLUSION

Our Fisher matrix formalism does not account for degeneracies between the cosmological parameters. Ref. [45] discusses the parameter degeneracies for the 1D flux power spectrum, in a model closely related to ours. The largest correlation was between the mean optical depth $\tau_{\rm eff}$ and the primordial power spectrum amplitude A_P . The WST is less sensitive to τ_{eff} and so it is likely this correlation will be reduced. Most other correlations were relatively small, with a correlation coefficient < 0.4. Our purpose here is to compare the WST as a summary statistic to the flux power spectrum; it is likely that the parameter correlations will be similar between the two statistics, and so our main conclusion that the 1D WST provides information not captured by the 1D flux power spectrum should be robust. Our small simulations also do not include the largest scales accessible to the survey. This particularly impacts our forecasts for n_P , which were substantially too pessimistic for the flux power spectrum, and are likely also too pessimistic for the WST.

There is no a priori reason to suppose that these and other systematics should affect the WST more than the flux power spectrum, but a robust analysis of systematic errors will be needed when using the WST to obtain parameter constraints from DESI. There are many avenues for future work. A maximum likelihood estimator for the WST coefficients from noisy and sparse data will need to be constructed. Our Fisher matrix forecast is simplis-




FIG. 3. (Left) Fisher information from the power spectrum and wavelet scattering transform at z=2 for our four parameters. For the WST coefficients, we divide them into the contribution from first and second order coefficients, with second order coefficients generally being more constraining. (Right) The Fisher information for the combined WST coefficients and flux power spectrum for different redshift bins at z=2,3,4.

tic and we will need to build a full likelihood function using some form of emulator of simulation based inference. However, despite these limitations, our work opens up the possibility to improve parameter constraints from the Lyman- α forest by multiple orders of magnitude.

SB was supported by NASA-80NSSC21K1840. MQ was supported by NSF grant AST-2107821 and the UCOP Dissertation Year Program. MFH is supported by a NASA FINESST grant No. ASTRO20-0022. Computing resources were provided by Frontera LRAC AST21005. The authors acknowledge the Frontera computing project at the Texas Advanced Computing Center (TACC) for providing HPC and storage resources that have contributed to the research results reported within this paper. Frontera is made possible by National Science Foundation award OAC-1818253. Analysis computations were performed using the resources of the UCR HPCC, which were funded by grants from NSF (MRI-2215705, MRI-1429826) and NIH (1S10OD016290-01A1).

- * htohfa@uw.edu
- † sbird@ucr.edu
- U. Seljak, A. Makarov, P. McDonald, S. F. Anderson, N. A. Bahcall, J. Brinkmann, S. Burles, R. Cen, M. Doi, J. E. Gunn, Ž. Ivezić, S. Kent, J. Loveday, R. H. Lupton, J. A. Munn, R. C. Nichol, J. P. Ostriker, D. J. Schlegel, D. P. Schneider, M. Tegmark, D. E. Berk, D. H. Weinberg, and D. G. York, Phys. Rev. D 71, 103515 (2005), arXiv:astro-ph/0407372 [astro-ph].
- [2] N. Palanque-Delabrouille, C. Yèche, J. Baur, C. Magneville, G. Rossi, J. Lesgourgues, A. Borde, E. Burtin, J.-M. LeGoff, J. Rich, M. Viel, and D. Weinberg, J. Cosmology Astropart. Phys. 2015, 011 (2015), arXiv:1506.05976 [astro-ph.CO].
- [3] V. Iršič, M. Viel, M. G. Haehnelt, J. S. Bolton, S. Cris-

- tiani, G. D. Becker, V. D'Odorico, G. Cupani, T.-S. Kim, T. A. M. Berg, S. López, S. Ellison, L. Christensen, K. D. Denney, and G. Worseck, Phys. Rev. D **96**, 023522 (2017), arXiv:1702.01764 [astro-ph.CO].
- [4] C. Yèche, N. Palanque-Delabrouille, J. Baur, and H. du Mas des Bourboux, J. Cosmology Astropart. Phys. 2017, 047 (2017), arXiv:1702.03314 [astro-ph.CO].
- [5] T. Theuns, M. Viel, S. Kay, J. Schaye, R. F. Carswell, and P. Tzanavaris, ApJ 578, L5 (2002), arXiv:astroph/0208418 [astro-ph].
- [6] N. Palanque-Delabrouille, C. Yèche, A. Borde, J.-M. Le Goff, G. Rossi, M. Viel, É. Aubourg, S. Bailey, J. Bautista, M. Blomqvist, A. Bolton, J. S. Bolton, N. G. Busca, B. Carithers, R. A. C. Croft, K. S. Dawson, T. Delubac, A. Font-Ribera, S. Ho, D. Kirkby, K.-G. Lee, D. Margala, J. Miralda-Escudé, D. Muna, A. D. Myers, P. Noterdaeme, I. Pâris, P. Petitjean, M. M. Pieri, J. Rich, E. Rollinde, N. P. Ross, D. J. Schlegel, D. P. Schneider, A. Slosar, and D. H. Weinberg, Astron. Astrophys. 559, A85 (2013), arXiv:1306.5896 [astro-ph.CO].
- [7] V. Iršič, M. Viel, T. A. M. Berg, V. D'Odorico, M. G. Haehnelt, S. Cristiani, G. Cupani, T.-S. Kim, S. López, S. Ellison, G. D. Becker, L. Christensen, K. D. Denney, G. Worseck, and J. S. Bolton, MNRAS 466, 4332 (2017), arXiv:1702.01761 [astro-ph.CO].
- [8] S. Chabanier, N. Palanque-Delabrouille, C. Yèche, J.-M. Le Goff, E. Armengaud, J. Bautista, M. Blomqvist, N. Busca, K. Dawson, T. Etourneau, A. Font-Ribera, Y. Lee, H. du Mas des Bourboux, M. Pieri, J. Rich, G. Rossi, D. Schneider, and A. Slosar, JCAP 2019 (7), 017, arXiv:1812.03554 [astro-ph.CO].
- [9] N. G. Karaçaylı, N. Padmanabhan, A. Font-Ribera, V. Iršič, M. Walther, D. Brooks, E. Gaztañaga, R. Kehoe, M. Levi, P. Ntelis, N. Palanque-Delabrouille, and G. Tarlé, MNRAS 509, 2842 (2022), arXiv:2108.10870 [astro-ph.CO].
- [10] A. Day, D. Tytler, and B. Kambalur, MNRAS 489, 2536 (2019).
- [11] DESI Collaboration and others, arXiv e-prints , arXiv:1611.00036 (2016), arXiv:1611.00036 [astro-ph.IM].

- [12] DESI Collaboration and others, arXiv e-prints arXiv:2306.06308 (2023), arXiv:2306.06308 [astroph.CO].
- [13] M. M. Pieri, S. Bonoli, J. Chaves-Montero, I. Pâris, M. Fumagalli, J. S. Bolton, M. Viel, P. Noterdaeme, J. Miralda-Escudé, N. G. Busca, H. Rahmani, C. Peroux, A. Font-Ribera, and S. C. Trager, in SF2A-2016: Proceedings of the Annual meeting of the French Society of Astronomy and Astrophysics, edited by C. Reylé, J. Richard, L. Cambrésy, M. Deleuil, E. Pécontal, L. Tresse, and I. Vauglin (2016) pp. 259– 266, arXiv:1611.09388 [astro-ph.CO].
- [14] F. Bernardeau, Y. Mellier, and L. van Waerbeke, A&A 389, L28 (2002), arXiv:astro-ph/0201032 [astro-ph].
- [15] M. Takada and B. Jain, MNRAS 344, 857 (2003), arXiv:astro-ph/0304034 [astro-ph].
- [16] E. Semboloni, T. Schrabback, L. van Waerbeke, S. Vafaei, J. Hartlap, and S. Hilbert, MNRAS 410, 143 (2011), arXiv:1005.4941 [astro-ph.CO].
- [17] L. Fu, M. Kilbinger, T. Erben, C. Heymans, H. Hildebrandt, H. Hoekstra, T. D. Kitching, Y. Mellier, L. Miller, E. Semboloni, P. Simon, L. Van Waerbeke, J. Coupon, J. Harnois-Déraps, M. J. Hudson, K. Kuijken, B. Rowe, T. Schrabback, S. Vafaei, and M. Velander, MNRAS 441, 2725 (2014), arXiv:1404.5469 [astro-ph.CO].
- [18] V. Ajani, J. Harnois-Déraps, V. Pettorino, and J.-L. Starck, A&A 672, L10 (2023), arXiv:2211.10519 [astro-ph.CO].
- [19] E. Paillas, C. Cuesta-Lazaro, P. Zarrouk, Y.-C. Cai, W. J. Percival, S. Nadathur, M. Pinon, A. de Mattia, and F. Beutler, MNRAS 522, 606 (2023), arXiv:2209.04310 [astro-ph.CO].
- [20] M. Welling, in International Conference on Artificial Intelligence and Statistics (2005).
- [21] S. Cheng, Astrophysics and cosmology with the scattering transform, Ph.D. thesis, Johns Hopkins University, Maryland (2021).
- [22] J. Anden and S. Mallat, IEEE Transactions on Signal Processing 62, 4114 (2014), arXiv:1304.6763 [cs.SD].
- [23] J. Andén and S. Mallat, in International Society for Music Information Retrieval Conference (2011).
- [24] L. Sifre and S. Mallat, in 2013 IEEE Conference on Computer Vision and Pattern Recognition (2013) pp. 1233–1240.
- [25] S. Mallat, arXiv e-prints , arXiv:1101.2286 (2011), arXiv:1101.2286 [math.FA].
- [26] S. Cheng, Y.-S. Ting, B. Ménard, and J. Bruna, MNRAS 499, 5902 (2020), arXiv:2006.08561 [astro-ph.CO].
- [27] S. Cheng and B. Ménard, MNRAS 507, 1012 (2021), arXiv:2103.09247 [astro-ph.CO].
- [28] D. Ribli, B. Á. Pataki, J. M. Zorrilla Matilla, D. Hsu, Z. Haiman, and I. Csabai, MNRAS 490, 1843 (2019), arXiv:1902.03663 [astro-ph.CO].
- [29] C. Pedersen, S. Ho, and M. Eickenberg, in *Machine Learning for Astrophysics* (2022) p. 40, arXiv:2307.14362

- [astro-ph.IM].
- [30] J. Bruna and S. Mallat, arXiv e-prints, arXiv:1203.1513 (2012), arXiv:1203.1513 [cs.CV].
- [31] B. Greig, Y.-S. Ting, and A. A. Kaurov, MNRAS 513, 1719 (2022), arXiv:2204.02544 [astro-ph.CO].
- [32] G. Valogiannis and C. Dvorkin, Phys. Rev. D 105, 103534 (2022), arXiv:2108.07821 [astro-ph.CO].
- [33] A. Lidz, C.-A. Faucher-Giguère, A. Dall'Aglio, M. Mc-Quinn, C. Fechner, M. Zaldarriaga, L. Hernquist, and S. Dutta, ApJ 718, 199 (2010), arXiv:0909.5210 [astro-ph.CO].
- [34] P. Gaikwad, R. Srianand, M. G. Haehnelt, and T. R. Choudhury, MNRAS 506, 4389 (2021), arXiv:2009.00016 [astro-ph.CO].
- [35] S. Bird, K. K. Rogers, H. V. Peiris, L. Verde, A. Font-Ribera, and A. Pontzen, J. Cosmology Astropart. Phys. 2019, 050 (2019), arXiv:1812.04654 [astro-ph.CO].
- [36] Y. Feng, S. Bird, L. Anderson, A. Font-Ribera, and C. Pedersen, Mp-gadget/mp-gadget: A tag for getting a doi (2018).
- [37] S. Bird, FSFE: Fake Spectra Flux Extractor, Astrophysics Source Code Library, record ascl:1710.012 (2017), ascl:1710.012.
- [38] M. Viel, M. G. Haehnelt, and V. Springel, MNRAS 354, 684 (2004), arXiv:astro-ph/0404600 [astro-ph].
- [39] T. S. Kim, J. S. Bolton, M. Viel, M. G. Haehnelt, and R. F. Carswell, MNRAS 382, 1657 (2007), arXiv:0711.1862 [astro-ph].
- [40] J. Bruna, S. Mallat, E. Bacry, and J.-F. Muzy, arXiv e-prints, arXiv:1311.4104 (2013), arXiv:1311.4104 [stat.ME].
- [41] M. Andreux, T. Angles, G. Exarchakis, R. Leonarduzzi, G. Rochette, L. Thiry, J. Zarka, S. Mallat, J. andén, E. Belilovsky, J. Bruna, V. Lostanlen, M. Chaudhary, M. J. Hirn, E. Oyallon, S. Zhang, C. Cella, and M. Eickenberg, arXiv e-prints, arXiv:1812.11214 (2018), arXiv:1812.11214 [cs.LG].
- [42] K.-G. Lee, N. Suzuki, and D. N. Spergel, AJ 143, 51 (2012), arXiv:1108.6080 [astro-ph.CO].
- [43] M. Qezlou, A. B. Newman, G. C. Rudie, and S. Bird, ApJ 930, 109 (2022), arXiv:2112.03930 [astro-ph.GA].
- [44] B. W. Lyke, A. N. Higley, J. N. McLane, D. P. Schurhammer, A. D. Myers, A. J. Ross, K. Dawson, S. Chabanier, P. Martini, N. G. Busca, H. d. Mas des Bourboux, M. Salvato, A. Streblyanska, P. Zarrouk, E. Burtin, S. F. Anderson, J. Bautista, D. Bizyaev, W. N. Brandt, J. Brinkmann, J. R. Brownstein, J. Comparat, P. Green, A. de la Macorra, A. Muñoz Gutiérrez, J. Hou, J. A. Newman, N. Palanque-Delabrouille, I. Pâris, W. J. Percival, P. Petitjean, J. Rich, G. Rossi, D. P. Schneider, A. Smith, M. Vivek, and B. A. Weaver, ApJS 250, 8 (2020), arXiv:2007.09001 [astro-ph.GA].
- [45] M. A. Fernandez, S. Bird, and M.-F. Ho, arXiv e-prints , arXiv:2309.03943 (2023), arXiv:2309.03943 [astro-ph.CO].