



Fisher's Exact Test Calculator — 2×2, 2×3, 2×4, 2×5 Tables | MedCalc

Use the Fisher's Exact Test to determine whether or not there is a significant association between two categorical variables. It is typically used as an alternative to the Chi-Squared Test (2x2 table) when cell counts are small.

	Category 1	Category 2
Group 1	11	13
Group 2	12	12

Results

Exact probability:	1.00000000
--------------------	------------

Interpretation

When the (two-sided) P-value (the probability of obtaining the observed result or a more extreme result) is less than the conventional 0.05, the conclusion is that there is a significant relationship between the two classification factors Group and Category.

Start using MedCalc on your desktop today

Join thousands of researchers who trust MedCalc for precision and speed. Get started for as little as **179 EUR** per year, or choose our popular lifetime license for **549 EUR** and never worry about renewals again.

[Order information](#)

Literature

- Altman DG (1991) Practical statistics for medical research. London: Chapman and Hall.

How to cite this page

- MedCalc Software Ltd. Fisher's Exact Test Calculator — 2×2, 2×3, 2×4, 2×5 Tables | MedCalc.
<https://www.medcalc.org/en/calc/fisher.php> (Version 23.7.1; accessed September 23, 2026)

Other free statistical calculators

- [Comparison of proportions](#)
- [McNemar test on paired proportions](#)
- [Two-way Chi-squared test](#)
- [Other statistical tests calculators](#)
- [Sample size calculation](#)
- [Post-hoc power analysis](#)

Trusted by researchers worldwide — [download the full MedCalc statistical software](#) and perform advanced statistical analysis in seconds. No registration required.

<https://www.medcalc.org/en/calc/fisher.php>

© 2026 MedCalc Software Ltd