



Fisher's Exact Test Calculator — 2×2, 2×3, 2×4, 2×5 Tables | MedCalc

Use the Fisher's Exact Test to determine whether or not there is a significant association between two categorical variables. It is typically used as an alternative to the Chi-Squared Test (2x2 table) when cell counts are small.

	Category 1	Category 2
Group 1	2	22
Group 2	17	7

OK

Results

Exact probability:	0.00001712
--------------------	------------

Interpretation

When the (two-sided) P-value (the probability of obtaining the observed result or a more extreme result) is less than the conventional 0.05, the conclusion is that there is a significant relationship between the two classification factors Group and Category.

Trusted by researchers worldwide — [download the full MedCalc statistical software](#) and perform advanced statistical analysis in seconds. No registration required.

Literature

- Altman DG (1991) Practical statistics for medical research. London: Chapman and Hall.

How to cite this page

- MedCalc Software Ltd. Fisher's Exact Test Calculator — 2×2, 2×3, 2×4, 2×5 Tables | MedCalc.
<https://www.medcalc.org/en/calc/fisher.php> (Version 23.7.1; accessed September 23, 2026)

Other free statistical calculators

- [Comparison of proportions](#)

- [McNemar test on paired proportions](#)
- [Two-way Chi-squared test](#)
- [Other statistical tests calculators](#)
- [Sample size calculation](#)
- [Post-hoc power analysis](#)

Trusted by researchers worldwide — [download the full MedCalc statistical software](#) and perform advanced statistical analysis in seconds. No registration required.

<https://www.medcalc.org/en/calc/fisher.php>

© 2026 MedCalc Software Ltd