




Contents lists available at ScienceDirect

NeuroImage

journal homepage: www.elsevier.com/locate/neuroimage

Patterns of thought: Population variation in the associations between large-scale network organisation and self-reported experiences at rest

Hao-Ting Wang^{a,*}, Danilo Bzdok^{b,c,d}, Daniel Margulies^e, Cameron Craddock^f, Michael Milham^f, Elizabeth Jefferies^a, Jonathan Smallwood^a

^a Department of Psychology, University of York, Heslington, York, England, UK

^b Department of Psychiatry, Psychotherapy and Psychosomatics, RWTH Aachen University, Germany

^c JARA-BRAIN, Jülich-Aachen Research Alliance, Germany

^d Parietal Team, INRIA, Neurospin, Bat 145, CEA Saclay, 91191, Gif-sur-Yvette, France

^e Neuroanatomy and Connectivity Group, Max Planck Institute for Cognition and Brain Sciences, Leipzig, Germany

^f Center for Biomedical Imaging and Neuromodulation, Nathan S. Kline Institute for Psychiatric Research, New York State Office of Mental Health, USA

ABSTRACT

Contemporary cognitive neuroscience recognises unconstrained processing varies across individuals, describing variation in meaningful attributes, such as intelligence. It may also have links to patterns of on-going experience. This study examined whether dimensions of population variation in different modes of unconstrained processing can be described by the associations between patterns of neural activity and self-reports of experience during the same period. We selected 258 individuals from a publicly available data set who had measures of resting-state functional magnetic resonance imaging, and self-reports of experience during the scan. We used machine learning to determine patterns of association between the neural and self-reported data, finding variation along four dimensions. ‘Purposeful’ experiences were associated with lower connectivity - in particular default mode and limbic networks were less correlated with attention and sensorimotor networks. ‘Emotional’ experiences were associated with higher connectivity, especially between limbic and ventral attention networks. Experiences focused on themes of ‘personal importance’ were associated with reduced functional connectivity within attention and control systems. Finally, visual experiences were associated with stronger connectivity between visual and other networks, in particular the limbic system. Some of these patterns had contrasting links with cognitive function as assessed in a separate laboratory session - purposeful thinking was linked to greater intelligence and better abstract reasoning, while a focus on personal importance had the opposite relationship. Together these findings are consistent with an emerging literature on unconstrained states and also underlines that these states are heterogeneous, with distinct modes of population variation reflecting the interplay of different large-scale networks.

Introduction

Unconstrained processing reflects important population level variation in measures of cognition, affect, and demographic/lifestyle factors. Psychological studies show that almost a third of on-going thought is unconstrained by events in the here-and-now (Killingsworth and Gilbert, 2010) with important links to cognitive and affective processing (Mooneyham and Schooler, 2013). In neuroscience, metrics defined from the brain during wakeful rest, describe the organisation of neural function at both the micro and macro scale (Glasser et al., 2016; Margulies et al., 2016). They also reflect individual differences in cognitive function (Finn et al., 2015), psychiatric conditions (Nooner et al., 2012) and demographic/lifestyle factors (Smith et al., 2015). These findings establish unconstrained neuro-cognitive processing as a core element of human cognition, highlighting the need to formally understand the underlying neural architecture, and the associated patterns of experience.

One perspective on unconstrained processing emphasises the role of memory, with contributions of conceptual and episodic representations to on-going thought (Binder et al., 2009; Gusnard et al., 2001). Psychological studies have shown patterns of unconstrained processing have links with memory retrieval, creativity and planning (Baird et al., 2012; Leszczynski et al., 2017; Medea et al., 2016; Poerio et al., 2017). Such evidence raises the possibility that episodic representations anchored in the medial temporal lobe (Moscovitch et al., 2016) or conceptual representation anchored in anterior temporal lobe (Ralph et al., 2017) contribute to on-going thought (Smallwood et al., 2016). It is hypothesised that these systems contribution to unconstrained states may be linked to the ability for these regions to become functionally decoupled from systems more directly involved in action and perception, allowing them to operate in an offline manner (Smallwood, 2013). This process of decoupling may also be important in neural systems closely allied to those involved in memory – the default mode network (Raichle et al.,

* Corresponding author. Department of Psychology, The University of York, Heslington, York, England, UK.

E-mail address: haoting.wang@york.ac.uk (H.-T. Wang).

<https://doi.org/10.1016/j.neuroimage.2018.04.064>

Received 27 March 2018; Received in revised form 25 April 2018; Accepted 28 April 2018

Available online xxx

1053-8119/© 2018 Elsevier Inc. All rights reserved.

2001). These regions of transmodal cortex are relatively distant in functional and structural space from systems involved in perception and action, potentially facilitating their role in stimulus independent aspects of cognition (Buckner and Krienen, 2013; Margulies et al., 2016; Mesulam, 1998). Together these ‘representational’ accounts of unconstrained processing highlight default mode and limbic networks as important candidate neural systems, especially when decoupled from systems directly involved in perception and action.

Alternative perspectives on unconstrained thought emerge from links between types of on-going experience and problems maintaining a task relevant goal in mind. This “executive-failure” view (Kane and McVay, 2012; McVay and Kane, 2010) takes as a starting point evidence that patterns of on-going thought, such as the experience of mind-wandering, are linked to problems on tasks including sustained attention (McVay and Kane, 2009) and measures of general aptitude and executive control (Mrazek et al., 2012). Task-based neuroimaging investigations highlight a network of regions that increase their activity across many different task situations - so called multiple demand regions (Duncan, 2010). These regions broadly correspond to three well described intrinsic networks: ventral attention, dorsal attention, and frontal-parietal networks. Since these systems are important for the effective performance of many different tasks then dysregulation within these systems could reflect the hypothesised ‘executive-failure’ contribution to aspects of on-going thought (McVay and Kane, 2010; Weissman et al., 2006).

Other aspects of unconstrained processing could reflect the importance of affective processes, or different modalities of processing. On-going thought is linked to mood state: Experimental inductions of mood (Smallwood et al., 2009; Smallwood and O’Connor, 2011), as well as natural fluctuations (Poerio et al., 2013; Ruby et al., 2013) impact on on-going thought. Contemporary accounts of emotional processing emphasise the role of limbic regions including the amygdala (Bzdok et al., 2013; Lindquist et al., 2012) and anterior aspects of the insula (Touroutoglou et al., 2012), suggesting these regions may be important in determining affective aspects of on-going thought. Psychological studies of on-going thought also suggest that another important dimension of unconstrained processing may reflect the different modalities of processing (Konishi et al., 2017; Smallwood et al., 2016). It has been shown, for example, that the visual system plays an important role in the expression of visual imagery (Ganis et al., 2004; Kosslyn et al., 2001). Recent work has extended this evidence to shown patterns of activity with visual regions are linked to the emergence of visual, non-verbal, elements of on-going thought (Raij and Riekkki, 2017). It is also possible that sensorimotor processes may be implicated in language processing during unconstrained processing, given that a role for these regions in language processing extends beyond production (Bzdok et al., 2016; Pulvermuller, 2010; Pulvermuller and Fadiga, 2010).

Current study

Our study aimed to identify patterns of intrinsic connectivity associated with different patterns of unconstrained states and examines their neuro-cognitive features from the perspectives outlined above. We used a large publicly available dataset, containing measures of resting-state functional magnetic resonance imaging (fMRI), and an accompanying self-report instrument describing cognition experienced during the resting-state (Gorgolewski et al., 2014; Nooner et al., 2012). We previously explored the relationships between patterns of on-going thought and measures of neural activity, such as the fractional amplitude of low frequency oscillations, as well as the regional homogeneity of neural activity, in a sub sample of this data set (Gorgolewski et al., 2014). In this study we focused on connectivity, we applied sparse canonical correlation analysis (SCCA) to obtain a conjoined decomposition of self-reports of experience with matrices of whole brain connectivity data. This analysis produces multivariate patterns that reflect dimensions of variation that are mutually constrained by both brain and experience. In this way we capitalize on the fact that self-reports of experience during

scanning and descriptions of on-going neural processing provide complementary descriptions of unconstrained cognition. Our analysis, therefore, helps define, at a population level, the shared links between brain patterns and different types of experience. It is important to note that this approach necessarily conflates state and trait related aspects of any brain-experience associations that are identified in this manner, and this aspect of our design should be borne in mind when interpreting our results. As our analytic approach respects the multivariate nature of brain and behaviour space, it can accommodate complex many-to-many relationships between patterns of connectivity and self-reports, and therefore is sensitive to the possibility of complex relationships in the underlying data. We took two steps to explore the robustness of the components that our study identifies. First, we use permutation testing to examine the extent to which our components are different from those that would be achieved based on a null distribution. Second we established whether these neuro-cognitive dimensions are associated with performance on a battery of available cognitive tasks, including measures of executive control and intelligence. When interpreting the results produced through our analysis it is important to give greater weight to components that show evidence of robustness in both comparisons.

We use the dimensions our analysis produces, and their links with cognitive function to evaluate the perspectives on unconstrained thought outlined earlier. ‘Representational’ accounts emphasise links with neural systems involved in memory, such as the limbic system, and regions of transmodal cortex, such as the default mode network. They highlight states with lower levels of functional communication between these regions and those more directly involved in external action. In contrast, ‘executive-failure’ accounts emphasise dysregulation in attention and control networks as contributing to patterns of on-going thought that are linked to problems in domain general task performance. Affective accounts highlight limbic regions as important hubs in aspects of on-going thoughts linked to emotion. Finally, modality specific influences on unconstrained thought may depend on information codes represented in regions that specialise in that particular types of information, such as a role of visual cortex in experiences dominated by images. Notably, some views lead to dissociable predictions with respect to cognitive performance. For example, executive-failure accounts predict patterns of thoughts linked to worse performance on measures of cognitive function, while representational accounts makes the alternative prediction.

Materials and methods

Participant

We analysed 258 participants (females = 162; age range 18–55, $M = 34.97$, $SD = 12.24$) obtained from the enhanced Nathan Kline Institute-Rockland sample (NKI-RS; http://fcon_1000.projects.nitrc.org/indi/enhanced/). Full details of the acquisition of this sample can be found in Nooner et al. (2012). We selected participants between 18 and 55 years old as our sample, this choice allowed us to maximise the cohesive nature of our sample. All the participants have the MRI data and less than 5 missing data points among the selected assessments.

Cognitive measures and questionnaires

Based on prior studies examining the links between spontaneous thought and cognitive performance (see Mooneyham and Schooler, 2013) we selected established neuropsychological measures linked to executive control, abstract reasoning and intelligence. The measures included the Delis-Kaplan Executive Function System (D-KEFS; Swanson, 2005), Wechsler Abbreviated Scale of Intelligence (WASI-II; Wechsler, 1999), and Wechsler Individual Achievement Test – Second Edition Abbreviated (WIAT-IIA; Wechsler, 2005). In D-KEFS we selected the tower test (move accuracy ratio), colour-word interference test (errors inhibition/switching), verbal fluency test (letter fluency - category fluency), design fluency test (design accuracy), trail making test

(sequencing errors score + set-loss errors score + time-discontinue errors score), and the proverb test (a measure of abstract semantic reasoning). We used the rescaled score ($M = 10$, $SD = 3$) in our analysis. Tasks measures that reflected error rates (i.e. the colour-word interference test and trail making test) were reversed, so that high rescaled scores indicated better task performance. All the scores were transformed to z-scores.

On-going cognition measure

The New York Cognition Questionnaire (NYC-Q) is a self-report tool used to assess the thoughts experienced at rest (Gorgolewski et al., 2014; Sanders et al., 2017). It assesses thoughts and feelings experienced during the resting-state period. The first section contains 23 questions about the content of thought. These questions covers the temporal, social, emotional aspects of spontaneous thoughts that have been shown to be important by prior studies (e.g. Ruby et al., 2013). Participants rated each question on a scale of 1 (Completely did not describe my thoughts) to 9 (Completely did describe my thoughts). The second section contains 8 questions about the forms thoughts take, capturing aspects of experience such as modality and detail associated with experience that prior studies suggest as important for spontaneous thoughts (Smallwood et al., 2016). Participants rated each question on a scale of 1 (Completely did not characterise my experience) to 9 (Completely did characterise my experience). In the current study we analysed the two sections together to provide single solutions that combined information on both the content form of experience. The full list of questions and the corresponding labels are presented in Table 1. The questionnaire was administrated once after the resting-state scan in order to assess experiences during the scanning session. For the full details of the NYC-Q, please refer to Gorgolewski et al., 2014. We have placed the questionnaire measure used in this study along with an example self-report collection task on GitHub at the following address: https://github.com/htwangtw/restingstate_thoughtreports.

MR data processing

Resting-state fMRI

We used resting-state fMRI to describe the general functional organisation of the brain. We selected resting-state multiband functional magnetic resonance imaging (R-mfMRI; TR = 1400msec; voxel size = 2 mm isotropic; duration = 10 min) for our analysis. Functional and structural data were pre-processed using Configurable Pipeline for the Analysis of Connectomes (C-PAC; <https://fcp-indi.github.io/>) to interface with FMRIB's Software Library (FSL version 5.0, www.fmrib.ox.ac.uk/fsl). Individual FLAIR and T1 weighted structural brain images were extracted using Brain Extraction Tool (BET). Structural images were linearly registered to the MNI-152 template using FMRIB's Linear Image Registration Tool (FLIRT). The resting-state functional data were pre-processed and analysed using the FMRI Expert Analysis Tool (FEAT). X, Y, Z displacement and the three axis rotations were used to calculate the mean frame displacement (FD), characterising movement of each participant during the scanning session (Power et al., 2014). Mean of the absolute values for FD were later used to account for subject specific head motion. No global signal regression was applied. The individual subject analysis involved: motion correction using MCFLIRT; slice-timing correction using Fourier space time series phase-shifting; spatial smoothing using a Gaussian kernel of FWHM 6 mm; bandpass filtering ($0.1 \text{ Hz} < f < 0.009 \text{ Hz}$); six motion parameters (as estimated by MCFLIRT) regressed out; cerebrospinal fluid and white matter signal regressed out (top five PCA components, CompCor method).

Connectivity matrices

To describe the functional architecture of the whole brain, we transformed the resting-state BOLD time series into connection strength values of the different networks for each participant. The whole brain parcellation was obtained from connectivity-based functional parcellation created by

Table 1

The new York cognition questionnaire (NYC-Q).

#	Questions	Labels
Q01	I thought about things I am currently worried about	Concerns
Q02	I thought about people I have just recently met	People
Q03	I thought of people I have known for a long time (friends)	Friend
Q04	I thought about members of my family	Family
Q05	I thought about an event that took place earlier today	Today - Past
Q06	I thought about an interaction I may possibly have in the future	Social - Future
Q07	I thought about an interaction with somebody that took place in the past	Social - Past
Q08	I thought about something that happened at a place very close to me	Near Location
Q09	I thought about something that made me feel guilty	Guilt
Q10	I thought about an event that may take place later today	Today - Plan
Q11	I thought about something that happened in the recent past (last couple of days but not today)	Recent Past
Q12	I thought about something that happened a long time ago in the past	Distant Past
Q13	I thought about something that made me angry	Anger
Q14	I thought about something that made me happy	Happiness
Q15	I thought about something that made me cheerful	Cheerfulness
Q16	I thought about something that made me calm	Calm
Q17	I thought about something that made me sad	Sadness
Q18	I thought about something that is important to me	Importance
Q19	I thought about something that could still happen today	Today - Future
Q20	I thought about something that may take place in the distant future	Distant Future
Q21	I thought about something that could take place in the near future (days or weeks but not today)	Near Future
Q22	I thought about personal worries	Worries
Q23	I thought about something that happened in a place far away from where I am now	Distant Location
Q24	In the form of images:	Image
Q25	In the form of words:	Words
Q26	Like an inner monologue or audiobook:	Monologue
Q27	Like a television program or film:	Film
Q28	Had a strong and consistent personal narrative:	Narrative
Q29	Had a clear sense of purpose:	Purpose
Q30	Vague and non-specific:	Vague
Q31	Fragmented and disjointed:	Fragment

Yeo and colleagues (Yeo et al., 2011). The 7 network parcellation was used in the current study. We split the networks into two hemispheres and extracted clusters. Two voxels are considered connected only if they are adjacent within the same x, y, or z direction. This yielded 57 clusters from the Yeo 7 networks parcellation. The implementation of spatial clusters extraction was retrieved from python library Nilearn (Abraham et al., 2014; <http://nilearn.github.io/>, version 0.3.1). Next, we extracted and then averaged the time series of all voxels within each cluster to create a cluster specific time series. We used these time series to create region-to-region symmetrical correlation matrices representing the correlations of the network signal that was computed for all the individual subjects. The off-diagonal of each correlation matrix contained 1596 unique region-region connection strengths (i.e., the upper or lower triangle of the network covariance matrix). This approach provided a measure of connection strength of the whole brain for each participant. Finally, Fisher's r-to-z transformation was applied to each network covariance matrix.

Conjoint decomposition of functional connectomes and mind-wandering measures

Decomposition method

We performed a sparse canonical correlation analysis (SCCA; see Hastie et al., 2015) on the functional connectomes and the NYC-Q reports, to yield latent components that reflect multivariate patterns across neural organisation and experience (For similar application, see Wang et al., 2017). SCCA maximised the linear correlation between the low-rank projections of two sets of multivariate data sets with sparse model to regularise the decomposition solutions a process that helps

Nested K-fold Cross-Validation with Model Selection




Fig. 1. A diagram of the nested k-fold cross-validation with model selection.

maximise the interpretability of the results. The regularisation function of choice is L_1 penalty, which produces 'sparse' coefficients, meaning that the canonical vectors (i.e., translating from full variables to a data matrix's low-rank components of variation) will contain a number of exactly zero elements. L_1 regularisation conducted (i) feature selection (i.e., select only relevant components) and (ii) model estimation (i.e., determine what combination of components best disentangles the neuro-cognitive relationship) in an identical process. This way we handle adverse behaviours of classical linear models in high-dimensional data. A reliable and robust open-source implementation of the SCCA method was retrieved as R package from CRAN (PMA, penalized multivariate analysis, version 1.0.9, Witten et al., 2009). The amount of L_1 penalty for the functional connectomes and the NYC-Q reports were chosen by cross-validation. The procedure is described below.

Model selection

We employed cross-validation (CV) to select the most useful model across population samples and avoid overfitting (Bzdok and Yeo, 2017). The amount of the two L_1 penalty terms for the functional connectomes and the NYC-Q reports, respectively, were chosen by a nested K-fold CV, where the coefficient for the penalty were chosen using a grid search to maximise the quality of CV objective metric. The objective metric of choice cumulative explained variances. The explained variance of each latent component was calculated using the squared canonical correlation. High explained variance suggests a high pattern recovery rate between the two data set. The sparse assumption is fundamentally in conflict with the statistical goal of finding components with high explained variance. Therefore we decided the number of components in the model before searching for the best parameter.

We performed confound removal on functional connectomes and the NYC-Q reports as suggested by prior studies (Smith et al., 2015). We removed the effects of nuisance variables from the dataset. These confound variables were sex, age, and head motion indicated by Jenkinson's mean FD (Jenkinson et al., 2002). The removal steps was performed on the training set in each CV fold. We standardized the confound by calculating the z-score, and also squared the three confound measures to account for potentially nonlinear effects of these confounds. The 6 resulting confounds were regressed out of both data matrices. The implementation of the confound removal method (Friston et al., 1994) was retrieved from python library Nilearn (Abraham et al., 2014; <http://nilearn.github.io/>, version 0.3.1).

The number of latent components was determined by a preliminary analysis with no sparsity and calculated the explained variances for the two datasets (i.e., brain network correlations and questionnaire ratings). The explained variance increased with the number of components and growth stabilised at 10 components. We selected the number of components based on the point where the tangent stabilised. This led to a model of 4 components, and it accounted for a total of 78% of the variance in connection strength and 29% of the variance in the self-report data. Next, we determined the two coefficients for the L_1 penalty terms that was associated with the best model performance with 4 latent components.

We searched for the best L_1 penalty values between 0.1 and 0.9 in 0.1 increments, which resulted in 81 set of parameters. For the nested K-Fold CV, we first separate the data into 5 consecutive folds after shuffling the data and retained one fold as the evaluation set ($N \sim 50$); the other four folds were used as the development set. The development set was further separated into 5 folds for parameter selection and each fold ($N \sim 40$) was used as the validation set once. The model was estimated on the training folds with all parameter sets, and after completion, we trained the model with the winning parameter on the whole development set and the finally tested the performance on the independent, unseen evaluation set. We selected the final parameters according to the best performance on the evaluation set across all folds of the outer CV loop (Fig. 1). This parameter set is used to train on the full development set and tested on the evaluation set. The parameter grid search and k-fold CV was conducted by the implementation in a Python library scikit-learn (Pedregosa et al., 2011; <http://scikit-learn.org/stable/>, version 0.18.2). The detailed algorithm for selecting the penalty values are presented in Appendix: Nested K-Fold CV.

The model with the best test performance was selected as the final model. The final model's sparsity coefficient are 0.8 (functional connectivity) and 0.5 (self-reports), and the out-of-sample explained variance was 48%. We used the ensuing canonical vectors of the winning SCCA model to compute the latent component scores. There are two sets of canonical scores in a latent component, a weighted sum of variables forms the canonical vectors. For each latent component, we averaged the z-score of the canonical scores of the connection strength and NYC-Q as the combined scores. These scores described the summary of the experience with both the neural basis and the content reports.

Test of component robustness

After identifying the well performed components in compressing the brain-experience data, we examined the robustness of the four components in two different ways. The permutation test is a purely data-driven strategy that access the chance of discovering components in null samples. We also leveraged the brain-experience components to explain the cognitive functions, so that we can identify meaningful patterns by well-established cognitive measurements.

Permutation test

We used permutation testing to assess the robustness of the components identified through our analysis. We constructed a null distribution for each canonical component by holding the functional connectivity data in place and randomising the row order of self-reports data. This permutation scheme broke the link of individual differences in the dataset, therefore testing the robustness of the components in the hypothetical population. By calculating the false-discovery rate in the null distribution, we can conclude the possibility of discovering our components by chance with the given penalty coefficients. Hypotheses that are accepted with a 5% level of significance. In the current analyses we adopt the permutation test with the FWE-corrected p-value by Smith et al. (2015) with data

argumentation to increase the size of the resampling datasets to 1000. The four components were compared to the first sparse canonical correlation of the permuted sample. The low-rank components are more relevant than the rest, therefore we yield more conservative p-value by comparing to the first canonical correlation only. We performed 5000 permutation tests to get enough estimates for 4 decimal places.

Group analysis

To determine how patterns of unconstrained neuro-cognitive activity related to performance on the battery of cognitive tests, we conducted an independent statistical analysis on the identical subjects. A Type III multivariate multiple regression with Pillai's trace test was applied to 4 individual scores for each of the latent components describing experience from the SCCA were the independent variables, and the original 8 measures of cognitive performance were the dependent variables that we hoped to be described by the linear combination of the latent components. Pillai's trace test is considered to be the most powerful and robust statistic for general use (Huberty and Olejnik, 2006). The p-values reported were based on Bonferroni correction. We also performed a principal components analysis (PCA) to identify the patterns of covariance among the 8 measures of cognitive performance and compressed the data. The relation between the principle score and the 4 brain-experience dimensions identified through SCCA was examined in a linear regression model with Pillai's trace test. The analysis was conducted in R (version 3.3.1). The multivariate multiple regression was conducted in R (version 3.3.1) using function 'Manova' in R package 'car' (companion to applied regression, version 2.1–5).

Code availability

The full analysis pipeline is freely available at <https://github.com/htwangtw/patterns-of-thought>.

Results

Determining constituent categories of experience

We used Sparse Canonical Correlation Analysis (SCCA) to determine connectome-wide dimensions that describe common variance shared by

descriptions of brain and experience. This took as input individual scores for the connections between each of the regions extracted from Yeo's 7 networks parcellation and the scores of each item of the New York Cognition Questionnaire (NYC-Q).

We applied SCCA with nested 5-fold CV as the model selection strategy. We obtained a model of 4 canonical components with penalty levels of 0.8 on the functional connectivity and 0.5 on the NYC-Q that indicated the best out-of-sample prediction on our data (see 3.5.2 Model Selection). The canonical correlations of the 4 latent components were 0.28, 0.19, 0.16, and 0.07. The latent components yielded by the best model are presented in Fig. 2. For the ease of presentation and interpretation, we summarized the components as network-network connectivity instead of 57-by-57 connectivity matrices. The heat maps describe the network-to-network correlations while the word clouds describe the loadings on the self-report items. The components in full and the heat map for the self-report items can be found in **Supplementary Materials**.

Component 1, describes patterns of reduced within network connectivity within all of the networks studied, with this pattern most prominent in the dorsal attention network. Between network connections are generally reduced, with the exception of visual to limbic. Sensorimotor was decoupled from all the other systems, and, in addition, the default and limbic were most decoupled from the attention networks. Experiential themes in Component 1 are dominated by themes related to deliberate planning with a verbal component (high loadings on "words", "monologue", "today-plan", "social-future", "purpose" and "deliberate"). We refer to this pattern of reports as reflecting thoughts with "purpose".

Component 2 is dominated by relatively higher within and between network connections. Connectivity within each network was strong with the exception of the limbic network. Between network connections were stronger, with this pattern most apparent in the connections between limbic and ventral attention. In addition, the visual network was strongly correlated with the other networks. This component is dominated by emotional responses (high loadings on "anger", "guilt", "cheerfulness" and "happiness") and social content ("friends" and "people"). We refer to this pattern of reports as reflecting "emotional" experience.

Component 3 emphasises reduced connections both between and within networks. Within network connectivity is weakest for the dorsal and ventral attention networks. Edge-to-edge connections are low, with




Fig. 2. Unique neuro-cognitive dimensions of population variation revealed by sparse canonical correlation analysis of measures of whole brain connectivity and self-reported descriptions of on-going experience. The heat map describes the canonical variate of the network-to-network connectivity between different Yeo networks. The connectivity matrices describes the coefficients from the model, separated into within and between network relationships. The word clouds reflect the coefficients on the relevant self-report items. In both cases the colour bars indicate the magnitudes of the coefficients. A detailed version of the canonical variates and alternative presentation of the self-report coefficients can be found in **Supplementary Material Figure S1–S5**.

the ventral and dorsal attention and fronto-parietal networks showing reduced correlations with each other as well as the visual and sensori-motor systems. This component was characterised by themes linked to personal “importance” with social temporal contents (“distant future”, “near future”, “social past”, “family” and “recent past”). We refer to this pattern of reports as reflecting “personal importance”.

Component 4 has the most heterogeneous pattern of within and between network connectivity. It is associated with stronger connections within networks with the exception of the limbic system. In addition, the visual system was strongly connected to all other networks, with this pattern most apparent for the limbic network. In contrast, lower network-to-network connectivity was observed between the default mode and sensori-motor and attention networks. This component is characterised by experiential patterns reflecting a modality difference in experience, with the highest loadings on “images” and lowest on “inner monologue”. We refer to this pattern of reports as describing “modality”.

The relationship between neuro-cognitive components and cognitive function assessed in the laboratory

Having documented four neuro-cognitive dimensions, we next examined the robustness of the components using two complementary approaches. We first used a permutation test to identify the chance of discovering components in a null samples as employed by Smith et al. (2015). The top three components passed the permutation test and the 4th component showed variance that was similar to that produced in a null sample (Component 1 $p = 0.0002$; Component 2 $p = 0.0010$; Component 3 $p = 0.0204$, Component 4 $p = 0.998$, $\alpha = 0.05$). This analysis suggests that Components 1–3 are unlikely to have occurred by chance. Component 4 may be a Type II error and so we discuss this component in only a limited manner moving forward.

Our next test of the robustness of our components is whether they explained unique patterns of expertise in our battery of cognitive tasks. We used multiple multivariate regression model in which performance on the battery of selected tasks was the dependent variables and the individual scores for each of the canonical components describing experience from the SCCA were the independent variables. In this analysis two of the four canonical components described significant variance in our

battery of tasks at multivariate level: Component 1 ($F(8, 246) = 2.21$, $p = .027$, $\eta^2_p = .067$) and Component 3 ($F(8, 246) = 2.56$, $p = .024$, $\eta^2_p = .068$).

In the univariate results of the significant component, Component 1 was linked to good performance in proverb test ($\beta = 0.48$, $t(251) = 3.27$, $p = .006$, 95% CI [0.191 0.766]) and both fluid intelligent tests WASI ($\beta = 0.39$, $t(251) = 2.74$, $p = .033$, 95% CI [0.111 0.677]) and WIAT ($\beta = 0.45$, $t(251) = 3.15$, $p = .009$, 95% CI [0.167 0.724]). Component 3 showed a reversed pattern of the cognitive functions related to Component 1: proverb test ($\beta = -0.45$, $t(251) = -0.14$, $p = .007$, 95% CI [-0.176 -0.727]); WASI ($\beta = -0.42$, $t(251) = -3.10$, $p = .012$, 95% CI [-0.151 -0.693]) and WIAT ($\beta = -0.41$, $t(251) = -3.06$, $p = .012$, 95% CI [-0.148 -0.682]). The relationships between the neuro-cognitive dimensions and the pattern of relationships on the full cognitive battery and the adjusted variable scatter plots of the significant results are summarized in the form of a heat map in Fig. 3.

Finally, we performed a simple principle component analysis on the eight task measures to explore the associations between experience and the structure of the laboratory data. The aim of this analysis was to see if the pattern retrieved from the univariate level in the previous multiple multivariate regression was related to the internal structure of the data. Component selection was determined based on the scree plot, and we accepted one component explaining 39% of the variance. The principle component loaded on the intelligence measures and the proverb test. We fitted a linear model to this data to understand the relationship to the four canonical components. The results are reported in Fig. 4. The overall linear model was significant ($F(4, 253) = 5.43$, $p = .0003$). In the linear regression model, Component 1 ($\beta = 0.82$, $t(253) = 3.5$, $p = .001$, 95% CI [0.36 1.29]) showed significant contribution to explaining the task principle component. Component 3 showed a negative correlation to the task components ($\beta = -0.69$, $t(253) = -3.04$, $p = .003$, 95% CI [-1.13 -0.24]). The relationships between tasks and the neuro-cognitive components here were similar to the ones uncovered by the multiple multivariate regression. In this analysis Component 4 ($\beta = 0.442$, $t(253) = 3.09$, $p = .002$, 95% CI [0.16 0.72]) showed a significant contribution in the regression model, but it did not pass the permutation test of robustness ($p = 0.998$). The related results should be treated cautiously. Together with our prior analysis, these results suggest that




Fig. 3. The relationship between the different neural-cognitive components and the measures assessed in the cognitive battery. The components 1 and 3 were significant at the multivariate level determined by multiple multivariate regression, indicated by the asterisk outside of the heat map. The cells with asterisk(s) indicates the significant results from the univariate test (bonferroni corrected) and the parameter estimates for each variable. CWI – Colour-word interference, DF – Design fluency, Pro- Proverbs, TOW – Tower of London, TMT – Trail making task, VF- Verbal Fluency, WASI – Wechsler Adult Intelligence Test, WIAT – Wechsler Individual Attainment Test. P-value significant codes: 0 **** 0.001 **** 0.01 ***.




Fig. 4. The principle component and its relationship to the different neural-cognitive components. The heat map describes the principle component of the task battery, and the scatter plots describe the association with the components identified in our study. Component 1 and 3 passed the permutation test for component robustness significantly contributed in explaining the principle component of the task. Component 4 showed a significant contribution in the regression model, but it did not pass the permutation test. The related results should be treated cautiously.

Components 1 and 3 are the most robust components identified in our study.

Discussion

We set out to describe different modes of neuro-cognitive patterns derived through the simultaneous decomposition of whole brain connectivity data with self-reports of on-going experience. We used a whole brain parcellation that describes cortical function in seven independent networks (Yeo et al., 2011). We combined this data with self-reports of the experience of our participants at rest, using a multivariate approach that allows for the possibility of many-to-many mappings between neural patterns and on-going cognition. Our analyses identified four stable canonical components, describing unique dimensions of neural-experiential variation. Permutation testing demonstrated the statistical robustness of Components 1–3. Furthermore, two components (1 and 3) described independent patterns of performance in a battery of commonly used cognitive measures. This association with cognitive performance that establishes a source of independent validity for these neuro-cognitive components since they are related to independent measures of cognitive performance. We next consider the fit between the dimensions produced by our analysis and theoretical views of unconstrained neuro-cognitive processing.

We found evidence broadly consistent with contemporary representational accounts of unconstrained processing. The neural patterns described by Component One reflect a pattern of reduced correlation between regions with links to memory and representation (e.g. limbic, default mode) from those with links to external behaviour (e.g. visual and sensorimotor cortex and attention networks). This pattern was associated with experiences characterised by a sense of purposefulness, and with verbally mediated content that was social and temporal in nature. Participants high on this dimension were proficient at generating abstract semantic links and performed well on measures of reasoning and intelligence. Together the features of Component One support the hypothesis that the functional decoupling of systems important for memory and representation are important for aspects of unconstrained cognition (Smallwood, 2013). This capacity may arise from the topographical organisation of the cortex, in which neural systems that can take on more transmodal properties tend to be located in regions that are more distant in functional and structural terms (Buckner and Krienen, 2013; Margulies et al., 2016; Mesulam, 1998). This spatial location may allow neural signals in these regions to take on properties that are discrepant from the neural signal more closely tethered to inputs describing the external world (Buckner and Krienen, 2013; Friston, 2013). The pattern identified by Component One, therefore, may reflect a pattern of population variation describing the hypothesised role of functional decoupling of memory and representational systems plays in the generation of more abstract aspects of human cognition (Margulies et al., 2016; Mesulam, 1998). Importantly, in our prior work, limbic and default mode networks were the most distant in functional connectivity terms from unimodal systems (Margulies et al., 2016).

Our data also highlights neural patterns that capture the hypothesised influence of attention and control on on-going thought (McVay and Kane,

2010). Component 3 highlights links between reduced connectivity within attention and control systems and patterns of thought that emphasise personal importance. This is associated with worse performance on measures of intelligence and reasoning. The combination of a focus on personally important themes linked to poor performance on measures of general aptitude, captures the hallmark psychological features of the current concerns X executive-failure accounts of on-going thought (McVay and Kane, 2010). This view suggests that failures in attentional control lead to highly personally relevant cognition to intrude into ongoing thought, leading to lapses in task performance. Importantly, the neural pattern described by this component emphasises dysregulated connectivity both within and between networks implicated in attention and control by task-based studies (Duncan, 2010). Our prior work established that spontaneous mind-wandering is linked to cortical thinning within regions linked to attention and control, such as the intra-parietal sulcus (Golchert et al., 2017). Spontaneous mind-wandering has been linked to worse cognitive control (Robison and Unsworth, 2018), as well as showing stronger links with attention related problems, including ADHD (Seli et al., 2015). Together with these prior studies, our data suggests that population variation in the intrinsic neural functioning within networks with an established role in external task performance captures the hypothesised contribution of executive-failure to patterns of on-going thought.

The method of decomposition used in the current study also highlighted patterns related to affective processing and the modality of the experience that are similar to those seen in our prior work that applied principal components analysis (PCA) to self-reported data only. Component Four places experiences with visual features (“images”) in opposition to experiences with verbal features (“monologue”), capturing dissociations between visual and verbal thinking observed in our prior studies (Konishi et al., 2017; Medea et al., 2016; Smallwood et al., 2016). The accompanying neural pattern were associated with higher connectivity between the visual network with other networks, in particular the limbic system. It is important to note that our permutation analysis failed to validate this component, so despite its association with task performance using the PCA analysis it should be treated with relative caution. Component Two loads on emotional experiences (“cheerfulness”, “anger”, “guilt” and “happiness”) with the exception of those that are unhappy (“sad”). In neural terms this component was characterised by high levels of connectivity, however, unlike Component Four, this was highest between limbic and ventral attention networks. This pattern of coupling is consistent with accounts that emphasise interactions between saliency and limbic systems in affective processing (Touroutoglou et al., 2012). In the case of Component Two permutation testing indicated this component was likely to be robust in statistical terms, however, we did not observe associations with task performance. As with Component Four, interpretations of Component Two should be made with caution in lieu of more empirical work.

Before closing it is worth considering several important limiting factors of our study. We focused on patterns of population variance in unconstrained neuro-cognitive processing that were measured once in each individual. Our study, therefore, cannot separate the influences state and traits on our observed components. Treating patterns of

unconstrained processing as a trait is common in both the psychological (McVay and Kane, 2009; Smallwood et al., 2013) and neural domains (Smith et al., 2015). Nonetheless, it remains an open question how consistent these components will be across individuals over time, as well as which aspects may be better described as traits. Importantly, by its very nature there are dimensions of experience that our study cannot adequately address. We cannot, for example, identify brain-experience associations that are highly dynamic in nature and in particular those that change rapidly within an individual. Insight into this issue could be achieved by a focus on dynamic rather than static connectivity (Kucyi, 2017). For example, the application of techniques such as sliding window analysis (Chang and Glover, 2010) or Hidden Markov models (Vidaurre et al., 2017) to fMRI could provide information that would complement our analyses. However, it may also be more important to examine these across multiple sessions within the same individuals, as this would also make it most possible to dissociate state from trait related influences on neural activity (Mueller et al., 2013). There are also types of experience that may be difficult to assess using the measure of retrospective experience sampling we have employed (Smallwood and Schooler, 2015). For important features of experience, such as whether it has evolving features (Mills et al., 2017), or when the participant is unaware of the content of their experience (Schooler, 2002), these experiential features may be best assessed using experience sampling techniques that capture momentary elements of experience (Smallwood, 2013).

There are a number of methodological improvements that could enhance future studies of brain-experience association. A recent benchmark study by Ciric and colleagues (Ciric et al., 2017) shows that scrubbing can improve the performance of resting state analyses. Regarding to the analysis pipeline, we gained hyper-parameters and best model with nested-CV an approach that can help prevent overfitting (Bzdok and Yeo, 2017). There are also alternative ways that could provide better tests of the robustness of the components we identified. We assessed the validity of the components in three different ways; 1) with a data-driven, non-parametric permutation test (Smith et al., 2015) that establishes the statistical validity of the identified components and 2) by establishing the relationship between the laboratory cognitive measures and 3) by consideration of their links with contemporary theoretical accounts of ongoing cognition. In our study, Components 1 and 3 were statistically significant in both cases and fitted well with contemporary accounts of ongoing cognition. Accordingly we place encourage readers to focus on these patterns from our data. There are alternative strategies that could help validate the robustness of patterns of brain-experience association. One approach could be to compare the relationship between multiple sessions within the same individual (Poldrack et al., 2015) and to have a larger sample that would allow the reproducibility of these results through a formal split-half validation procedure. To achieve this latter aim for future studies, we have placed the questionnaire measure used in this study along with an example self-report collection task on GitHub at the following address: https://github.com/htwangtw/restingstate_thoughtreports. We encourage interested investigators to apply these measures in their resting-state investigation and to also upload the resultant data onto open fMRI. These studies could be used in conjunction with the openly access data used in this study to enable future investigations the opportunity to cross validate experiential analyses in a more sophisticated manner than we have been able to achieve in this study. The analysis pipeline of the current study can be further unified into one frame work that benefits from both validation strategies. We can include the number of components along with penalty coefficients in the hyper-parameters determined in the CV process, or determine the best penalty terms with the first component. The permutation test will then identify the reliable components occurring above chance level. After all the data-driven component selection, we can examine the survived components through their relations with well-documented cognitive measures and conclude the meaningful patterns. Finally, it is likely that our measure of on-going thought lacks important questions regarding the content of experience. It will be

important, therefore, in the future to examine the relationships of the type described in this study with a more exhaustive description of on-going experience. We hope that by publishing our questionnaire collection task in a GitHub repository we will be able to harness the power of the broader community to help generate and test plausible questions for use in future studies.

Funding

EJ was supported by ERC (SEMBIND - 283530). JS was supported by ERC (WANDERINGMINDS - 646927), Volkswagen Foundation (Wandering Minds - 89440 and 89439) and the John Templeton Foundation, "Prospective Psychology Stage 2: A Research Competition". DB was supported by the German Research Foundation (DFG, BZ2/2-1, BZ2/3-1, and BZ2/4-1), International Research Training Group (IRTG2150), Amazon AWS Research Grant (2016 and 2017), the German National Academic Foundation, and the START-Program of the Faculty of Medicine and an Exploratory Research Space Seed Fund, RWTH Aachen University.

Conflicts of interest

The authors have declared that no competing interests exist.

Acknowledgements

The authors thank the Charlotte Murphy, Mladen Sormaz, Giulia Poerio, Irene De Caso and the rest of the semantics and mind-wandering lab members for informative discussions; Theodoros Karapanagiotidis and Deniz Vatansever for their help on fMRI data management and preprocessing.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.neuroimage.2018.04.064>.

Appendix. Nested K-Fold CV

- 1 Separate the model into 5 folds. In every iteration, 1 fold is the test set and the rest are the development set.
- 2 For each outer fold:
 - a For each parameter set to be considered:
 - i Separate the development set into 5 folds. In every iteration, 1 fold is the validation set and the rest are the training set.
 - ii For each inner fold:
 - 1 Train the model on the training set
 - 2 Calculate test error in the validation set
 - iii Compute the average inner CV test error.
 - b Choose the best parameter set with minimum average test error.
 - c Use this parameter set to train on the development set.
 - d Calculate test error in the test set
- 3 Determine the optimal model based on the outer fold test error
- 4 Train the full dataset on the optimal model

References

- Abraham, A., Pedregosa, F., Eickenberg, M., Gervais, P., Mueller, A., Kossaifi, J., et al., 2014. Machine learning for neuroimaging with scikit-learn. *Front. Neuroinf.* 8.
- Baird, B., Smallwood, J., Mrazek, M.D., Kam, J.W., Franklin, M.S., Schooler, J.W., 2012. Inspired by distraction mind wandering facilitates creative incubation. *Psychol. Sci.* 0956797612446024.
- Binder, J.R., Desai, R.H., Graves, W.W., Conant, L.L., 2009. Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cereb. Cortex* 19 (12), 2767–2796. <https://doi.org/10.1093/cercor/bhp055>.
- Buckner, R.L., Krienen, F.M., 2013. The evolution of distributed association networks in the human brain. *Trends Cognit. Sci.* 17 (12), 648–665.

- Bzdok, D., Hartwigsen, G., Reid, A., Laird, A.R., Fox, P.T., Eickhoff, S.B., 2016. Left inferior parietal lobe engagement in social cognition and language. *Neurosci. Biobehav. Rev.* 68, 319–334.
- Bzdok, D., Laird, A.R., Zilles, K., Fox, P.T., Eickhoff, S.B., 2013. An investigation of the structural, connectional, and functional subspecialization in the human amygdala. *Hum. Brain Mapp.* 34 (12), 3247–3266.
- Bzdok, D., Yeo, B.T., 2017. Inference in the age of big data: future perspectives on neuroscience. *Neuroimage*.
- Chang, C., Glover, G.H., 2010. Time-frequency dynamics of resting-state brain connectivity measured with fMRI. *Neuroimage* 50 (1), 81–98. <https://doi.org/10.1016/j.neuroimage.2009.12.011>.
- Ciric, R., Wolf, D.H., Power, J.D., Roalf, D.R., Baum, G.L., Ruparel, K., et al., 2017. Benchmarking of participant-level confound regression strategies for the control of motion artifact in studies of functional connectivity. *Neuroimage* 154, 174–187.
- Duncan, J., 2010. The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. *Trends Cognit. Sci.* 14 (4), 172–179.
- Finn, E.S., Shen, X., Scheinost, D., Rosenberg, M.D., Huang, J., Chun, M.M., et al., 2015. Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. *Nat. Neurosci.* 18 (11), 1664–1671. <https://doi.org/10.1038/nn.4135>.
- Friston, K., 2013. Life as we know it. *J. R. Soc. Interface* 10 (86), 20130475. <https://doi.org/10.1098/rsif.2013.0475>.
- Friston, K., Holmes, A.P., Worsley, K.J., Poline, J.P., Frith, C.D., Frackowiak, R.S., 1994. Statistical parametric maps in functional imaging: a general linear approach. *Hum. Brain Mapp.* 2 (4), 189–210.
- Ganis, G., Thompson, W.L., Kosslyn, S.M., 2004. Brain areas underlying visual mental imagery and visual perception: an fMRI study. *Brain Res. Cognit. Brain Res.* 20 (2), 226–241. <https://doi.org/10.1016/j.cogbrainres.2004.02.012>.
- Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub, E., et al., 2016. A multi-modal parcellation of human cerebral cortex. *Nature* 536 (7615), 171–178. <https://doi.org/10.1038/nature18933>.
- Golchert, J., Smallwood, J., Jefferies, E., Seli, P., Huntenburg, J.M., Liem, F., et al., 2017. Individual variation in intentionality in the mind-wandering state is reflected in the integration of the default-mode, fronto-parietal, and limbic networks. *Neuroimage* 146, 226–235. <https://doi.org/10.1016/j.neuroimage.2016.11.025>.
- Gorgolewski, K.J., Lurie, D., Urchs, S., Kipping, J.A., Craddock, R.C., Milham, M.P., et al., 2014. A correspondence between individual differences in the brain's intrinsic functional architecture and the content and form of self-generated thoughts. *PLoS One* 9 (5), e97176.
- Gusnard, D.A., Raichle, M.E., Raichle, M.E., 2001. Searching for a baseline: functional imaging and the resting human brain. *Nat. Rev. Neurosci.* 2 (10), 685–694. <https://doi.org/10.1038/35094500>.
- Hastie, T., Tibshirani, R., Wainwright, M., 2015. *Statistical Learning with Sparsity: the Lasso and Generalizations*. CRC press.
- Huberty, C.J., Olejnik, S., 2006. *Applied MANOVA and Discriminant Analysis*, vol. 498. John Wiley & Sons.
- Jenkinson, M., Bannister, P., Brady, M., Smith, S., 2002. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* 17 (2), 825–841.
- Kane, M.J., McVay, J.C., 2012. What mind wandering reveals about executive-control abilities and failures. *Curr. Dir. Psychol. Sci.* 21 (5), 348–354. <https://doi.org/10.1177/0963721412454875>.
- Killingsworth, M.A., Gilbert, D.T., 2010. A wandering mind is an unhappy mind. *Science* 330 (6006), 932–932.
- Konishi, M., Brown, K., Battaglini, L., Smallwood, J., 2017. When attention wanders: pupillometric signatures of fluctuations in external attention. *Cognition* 168, 16–26. <https://doi.org/10.1016/j.cognition.2017.06.006>.
- Kosslyn, S.M., Ganis, G., Thompson, W.L., 2001. Neural foundations of imagery. *Nat. Rev. Neurosci.* 2 (9), 635–642. <https://doi.org/10.1038/35090055>.
- Kucyi, A., 2017. Just a thought: how mind-wandering is represented in dynamic brain connectivity. *Neuroimage*. Published online July 3, 2017.
- Leszczynski, M., Chaieb, L., Reber, T.P., Derner, M., Axmacher, N., Fell, J., 2017. Mind wandering simultaneously prolongs reactions and promotes creative incubation. *Sci. Rep.* 7 (1), 10197.
- Lindquist, K.A., Wager, T.D., Kober, H., Bliss-Moreau, E., Barrett, L.F., 2012. The brain basis of emotion: a meta-analytic review. *Behav. Brain Sci.* 35 (3), 121–143. <https://doi.org/10.1017/S0140525X11000446>.
- Margulies, D.S., Ghosh, S.S., Goulas, A., Falkiewicz, M., Huntenburg, J.M., Langs, G., et al., 2016. Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proc. Natl. Acad. Sci. U. S. A.* 113 (44), 12574–12579. <https://doi.org/10.1073/pnas.1608282113>.
- McVay, J.C., Kane, M.J., 2009. Conducting the train of thought: working memory capacity, goal neglect, and mind wandering in an executive-control task. *J. Exp. Psychol. Learn. Mem. Cognit.* 35 (1), 196–204. <https://doi.org/10.1037/a0014104>.
- McVay, J.C., Kane, M.J., 2010. Does mind wandering reflect executive function or executive failure? Comment on Smallwood and Schooler (2006) and Watkins (2008). *Psychol. Bull.* 136 (2), 188–197. <https://doi.org/10.1037/a0018298> discussion 198–207.
- Medea, B., Karapanagiotidis, T., Konishi, M., Ottaviani, C., Margulies, D., Bernasconi, A., et al., 2016. How do we decide what to do? Resting-state connectivity patterns and components of self-generated thought linked to the development of more concrete personal goals. *Exp. Brain Res.* <https://doi.org/10.1007/s00221-016-4729-y>.
- Mesulam, M.-M., 1998. From sensation to cognition. *Brain a J. neurology* 121 (6), 1013–1052.
- Mills, C., Raffaelli, Q., Irving, Z.C., Stan, D., Christoff, K., 2017. Is an off-task mind a freely-moving mind? Examining the relationship between different dimensions of thought. *Conscious. Cognit.* <https://doi.org/10.1016/j.concog.2017.10.003>.
- Mooneyham, B.W., Schooler, J.W., 2013. The costs and benefits of mind-wandering: a review. *Can. J. Exp. Psychology/Revue Can. de Psychol. expérimentale* 67 (1), 11.
- Moscovitch, M., Cabeza, R., Winocur, G., Nadel, L., 2016. Episodic memory and beyond: the Hippocampus and neocortex in transformation. *Annu. Rev. Psychol.* 67 (67) <https://doi.org/10.1146/annurev-psych-113011-143733>, 105–+.
- Mrazek, M.D., Smallwood, J., Franklin, M.S., Chin, J.M., Baird, B., Schooler, J.W., 2012. The role of mind-wandering in measurements of general aptitude. *J. Exp. Psychol. General* 141 (4), 788.
- Mueller, S., Wang, D., Fox, M.D., Yeo, B.T., Sepulcre, J., Sabuncu, M.R., et al., 2013. Individual variability in functional connectivity architecture of the human brain. *Neuron* 77 (3), 586–595.
- Nooner, K.B., Colcombe, S.J., Tobe, R.H., Mennes, M., Benedict, M.M., Moreno, A.L., et al., 2012. The NKI-Rockland sample: a model for accelerating the pace of discovery science in psychiatry. *Front. Neurosci.* 6.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., et al., 2011. Scikit-learn: machine learning in Python. *J. Mach. Learn. Res.* 12 (Oct), 2825–2830.
- Poerio, G.L., Sormaz, M., Wang, H.T., Margulies, D., Jefferies, E., Smallwood, J., 2017. The role of the default mode network in component processes underlying the wandering mind. *Soc. Cognit. Affect. Neurosci.* <https://doi.org/10.1093/scan/nrx041>.
- Poerio, G.L., Totterdell, P., Miles, E., 2013. Mind-wandering and negative mood: does one thing really lead to another? *Conscious. Cognit.* 22 (4), 1412–1421.
- Poldrack, R.A., Laumann, T.O., Koyejo, O., Gregory, B., Hover, A., Chen, M.-Y., et al., 2015. Long-term neural and physiological phenotyping of a single human. *Nat. Commun.* 6, 8885.
- Power, J.D., Mitra, A., Laumann, T.O., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2014. Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage* 84, 320–341. <https://doi.org/10.1016/j.neuroimage.2013.08.048>.
- Pulvermüller, F., 2010. Brain embodiment of syntax and grammar: discrete combinatorial mechanisms spelt out in neuronal circuits. *Brain Lang.* 112 (3), 167–179. <https://doi.org/10.1016/j.bandl.2009.08.002>.
- Pulvermüller, F., Fadiga, L., 2010. Active perception: sensorimotor circuits as a cortical basis for language. *Nat. Rev. Neurosci.* 11 (5), 351–360. <https://doi.org/10.1038/nrn2811>.
- Raichle, M.E., MacLeod, A.M., Snyder, A.Z., Powers, W.J., Gusnard, D.A., Shulman, G.L., 2001. A default mode of brain function. *Proc. Natl. Acad. Sci. U. S. A.* 98 (2), 676–682. <https://doi.org/10.1073/pnas.98.2.676>.
- Raij, T.T., Riekk, T.J.J., 2017. Dorsomedial prefrontal cortex supports spontaneous thinking per se. *Hum. Brain Mapp.* 38 (6), 3277–3288. <https://doi.org/10.1002/hbm.23589>.
- Ralph, M.A.L., Jefferies, E., Patterson, K., Rogers, T.T., 2017. The neural and computational bases of semantic cognition. *Nat. Rev. Neurosci.* 18 (1), 42–55. <https://doi.org/10.1038/nrn.2016.150>.
- Robison, M.K., Unsworth, N., 2018. Cognitive and contextual correlates of spontaneous and deliberate mind-wandering. *J. Exp. Psychol. Learn. Mem. Cognit.* 44 (1), 85–98. <https://doi.org/10.1037/xlm0000444>.
- Ruby, F.J., Smallwood, J., Engen, H., Singer, T., 2013. How self-generated thought shapes mood—the relation between mind-wandering and mood depends on the socio-temporal content of thoughts. *PLoS One* 8 (10), e77554.
- Sanders, J.G., Wang, H.-T., Schooler, J., Smallwood, J., 2017. Can I get me out of my Head? Exploring strategies for controlling the self-referential aspects of the mind-wandering state during reading. *Q. J. Exp. Psychol.* 70 (6), 1053–1062. <https://doi.org/10.1080/17470218.2016.1216573>.
- Schooler, J.W., 2002. Re-presenting consciousness: dissociations between experience and meta-consciousness. *Trends Cognit. Sci.* 6 (8), 339–344.
- Seli, P., Smallwood, J., Cheyne, J.A., Smilek, D., 2015. On the relation of mind wandering and ADHD symptomatology. *Psychonomic Bull. Rev.* 22 (3), 629–636.
- Smallwood, J., 2013. Distinguishing how from why the mind wanders: a process-occurrence framework for self generated thought. *Psychol. Bull.*
- Smallwood, J., Fitzgerald, A., Miles, L.K., Phillips, L.H., 2009. Shifting moods, wandering minds: negative moods lead the mind to wander. *Emotion* 9 (2), 271.
- Smallwood, J., Karapanagiotidis, T., Ruby, F., Medea, B., de Caso, I., Konishi, M., et al., 2016. Representing representation: integration between the temporal lobe and the posterior cingulate influences the content and form of spontaneous thought. *PLoS One* 11 (4). <https://doi.org/10.1371/journal.pone.0152272> e0152272.
- Smallwood, J., O'Connor, R.C., 2011. Imprisoned by the past: unhappy moods lead to a retrospective bias to mind wandering. *Cognit. Emot.* 1–10. <https://doi.org/10.1080/02699931.2010.545263>.
- Smallwood, J., Ruby, F.J., Singer, T., 2013. Letting go of the present: mind-wandering is associated with reduced delay discounting. *Conscious. Cognit.* 22 (1), 1–7.
- Smallwood, J., Schooler, J.W., 2015. The science of mind wandering: empirically navigating the stream of consciousness. *Annu. Rev. Psychol.* 66, 487–518.
- Smith, S.M., Nichols, T.E., Vidaurre, D., Winkler, A.M., Behrens, T.E.J., Glasser, M.F., et al., 2015. A positive-negative mode of population covariation links brain connectivity, demographics and behavior. *Nat. Neurosci.* 18 (11), 1565–1567. <http://doi.org/10.1038/nn.4125>.
- Swanson, J., 2005. The delis-kaplan executive function system: a review. *Can. J. Sch. Psychol.* 20 (1–2), 117–128.
- Touroutoglou, A., Hollenbeck, M., Dickerson, B.C., Barrett, L.F., 2012. Dissociable large-scale networks anchored in the right anterior insula subserve affective experience and attention. *Neuroimage* 60 (4), 1947–1958.
- Vidaurre, D., Smith, S.M., Woolrich, M.W., 2017. Brain network dynamics are hierarchically organized in time. *Proc. Natl. Acad. Sci. U. S. A.* 114 (48), 12827–12832. <https://doi.org/10.1073/pnas.1705120114>.

- Wang, H.-T., Poerio, G., Murphy, C., Bzdok, D., Jefferies, E., Smallwood, J., 2017. Dimensions of experience: exploring the heterogeneity of the wandering mind. *Psychol. Sci.* 0956797617728727.
- Wechsler, D., 1999. Wechsler Abbreviated Scale of Intelligence. The Psychological Corporation, New York, NY.
- Wechsler, D., 2005. Wechsler Individual Achievement Test 2nd Edition (WIAT II). The Psychological Corporation, New York, NY.
- Weissman, D.H., Roberts, K., Visscher, K., Woldorff, M., 2006. The neural bases of momentary lapses in attention. *Nat. Neurosci.* 9 (7), 971.
- Witten, D.M., Tibshirani, R., Hastie, T., 2009. A penalized matrix decomposition, with applications to sparse principal components and canonical correlation analysis. *Biostatistics* 10 (3), 515–534.
- Yeo, B.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M., et al., 2011. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J. Neurophysiol.* 106 (3), 1125–1165.