
A
Nine pre-specified signalling axes
Periostin-integrin
Fibronectin-integrin
Inflammatory
Notch
Angiogenic
Collagen-integrin/DDR
TGF-beta
PDGF
Chemokine
d = +0.90, p = 0.153
d = +1.99, p = 0.014
d = +0.60, p = 0.333
d = +0.32, p = 0.599
d = +0.37, p = 0.534
d = +0.03, p = 0.964
d = -0.01, p = 0.983
d = -0.44, p = 0.469
d = -0.30, p = 0.613
−0.5
0.0
0.5
1.0
1.5
log2 ratio of axis score (failed / matured)
B
Twelve most significant individual interactions (exploratory)
T cell → Pericyte TNF–TNFRSF1A
Endothelial → Myofibroblast FN1–ITGAV
AF stromal → Myeloid FN1–ITGB1
AF resident → Myeloid FN1–ITGAV
AF stromal → Myeloid FN1–ITGAV
AF resident → AF resident FN1–ITGB1
AF resident → AF stromal FN1–ITGB1
SMC contractile → SMC contractile TGFB3–TGFBR2
AF resident → Endothelial FN1–ITGAV
SMC contractile → Myeloid FN1–ITGB1
Myeloid → Myofibroblast FN1–ITGAV
AF resident → Myeloid FN1–ITGB1
−2.0
−1.5
−1.0
−0.5
0.0
0.5
1.0
1.5
log2 ratio of interaction score (failed / matured)