
A
Genes per compartment, before and after cell-of-origin screening
Myeloid
Endothelial
Mast
T cell
Myofibroblast
SMC contractile
Pericyte
16
7
7
5
2
2
2
2
1
1
1
1
1
1
Before screening
After screening
0.0
2.5
5.0
7.5
10.0
12.5
15.0
17.5
Differentially expressed genes
B
Matrisome programmes and computed zonation
ECM1 interstitial
ECM2 basement membrane
ECM3 microfibril
ECM4 fibrillar collagen
ECM5 fibronectin/ versican
Zonation index
p = 0.29
p = 0.99
p = 0.76
p = 0.73
p = 0.41
p = 0.98
−0.02
0.00
0.02
0.04
0.06
0.08
0.10
Difference in programme usage (failed - matured)
C
Whole-tissue bulk, GSE220796 (11 failed vs 9 matured)
COL8A1
LOXL1
FN1
MFAP5
COL1A1
ACTA2
MYH11
DCN
COL3A1
POSTN
TIMP1
PI16
FDR = 0.02*
FDR = 0.12
FDR = 0.23
FDR = 0.52
FDR = 0.57
FDR = 0.84
FDR = 0.93
FDR = 0.99
FDR = 0.81
FDR = 0.71
FDR = 0.05*
FDR = 0.54
−1.0
−0.5
0.0
0.5
1.0
1.5
2.0
log2 fold change (failed vs matured)