

Lecture 6: Kinematics: Velocity Kinematics - the Jacobian

- Skew Symmetric Matrices

Lecture 6: Kinematics: Velocity Kinematics - the Jacobian

- Skew Symmetric Matrices
- Linear and Angular Velocities of a Moving Frame

Angular Velocity for Describing Rotation around Fixed Axis

When a rigid body rotates around a fixed axis

- Every point of the body moves in a circle

Angular Velocity for Describing Rotation around Fixed Axis

When a rigid body rotates around a fixed axis

- Every point of the body moves in a circle
- As the body rotates a perpendicular from any point of the body to the axis sweeps the same an angle θ

Angular Velocity for Describing Rotation around Fixed Axis

When a rigid body rotates around a fixed axis

- Every point of the body moves in a circle
- As the body rotates a perpendicular from any point of the body to the axis sweeps the same an angle θ
- If \vec{k} is a unit vector in the direction of the axis of rotation, then the angular velocity is given by

$$\vec{\omega} = \frac{d}{dt}\theta \cdot \vec{k}$$

Angular Velocity for Describing Rotation around Fixed Axis

When a rigid body rotates around a fixed axis

- Every point of the body moves in a circle
- As the body rotates a perpendicular from any point of the body to the axis sweeps the same an angle θ
- If \vec{k} is a unit vector in the direction of the axis of rotation, then the angular velocity is given by

$$\vec{\omega} = \frac{d}{dt}\theta \cdot \vec{k}$$

- The linear velocity of any point of the body is then

$$\vec{v} = \vec{\omega} \times \vec{r}$$

where \vec{r} is the vector from the origin, which is assumed to lie on the axis.

Skew Symmetric Matrices

An $n \times n$ matrix S is said to be **skew symmetric** and denoted $S \in so(n)$ if

$$S + S^T = 0$$

Skew Symmetric Matrices

An $n \times n$ matrix S is said to be **skew symmetric** and denoted $S \in so(n)$ if

$$S + S^T = 0$$

The matrix equation written for each item of $S \in so(n)$ is

$$s_{ij} + s_{ji} = 0, \quad i = 1, \dots, n, \quad j = 1, \dots, n$$

Skew Symmetric Matrices

An $n \times n$ matrix S is said to be **skew symmetric** and denoted $S \in so(n)$ if

$$S + S^T = 0$$

The matrix equation written for each item of $S \in so(n)$ is

$$s_{ij} + s_{ji} = 0, \quad i = 1, \dots, n, \quad j = 1, \dots, n$$

$$\Rightarrow \left\{ s_{ii} = 0 \quad \text{and} \quad s_{ij} = -s_{ji} \quad \forall i \neq j \right\}$$

Skew Symmetric Matrices

An $n \times n$ matrix S is said to be **skew symmetric** and denoted $S \in so(n)$ if

$$S + S^T = 0$$

The matrix equation written for each item of $S \in so(n)$ is

$$s_{ij} + s_{ji} = 0, \quad i = 1, \dots, n, \quad j = 1, \dots, n$$

$$\Rightarrow \left\{ s_{ii} = 0 \quad \text{and} \quad s_{ij} = -s_{ji} \quad \forall i \neq j \right\}$$

For example, if $n = 3$, then any $S \in so(3)$ has the form

$$S = \begin{bmatrix} 0 & * & * \\ * & 0 & * \\ * & * & 0 \end{bmatrix}$$

Skew Symmetric Matrices

An $n \times n$ matrix S is said to be **skew symmetric** and denoted $S \in so(n)$ if

$$S + S^T = 0$$

The matrix equation written for each item of $S \in so(n)$ is

$$s_{ij} + s_{ji} = 0, \quad i = 1, \dots, n, \quad j = 1, \dots, n$$

$$\Rightarrow \left\{ s_{ii} = 0 \quad \text{and} \quad s_{ij} = -s_{ji} \quad \forall i \neq j \right\}$$

For example, if $n = 3$, then any $S \in so(3)$ has the form

$$S = \begin{bmatrix} 0 & * & * \\ * & 0 & * \\ * & * & 0 \end{bmatrix} = \begin{bmatrix} 0 & -s_3 & s_2 \\ s_3 & 0 & -s_1 \\ -s_2 & s_1 & 0 \end{bmatrix}$$

Skew Symmetric Matrices (Cont'd)

Given $\vec{a} = [a_x, a_y, a_z]^T$, define

$$S(\vec{a}) = \begin{bmatrix} 0 & -a_z & a_y \\ a_z & 0 & -a_x \\ -a_y & a_x & 0 \end{bmatrix}$$

Skew Symmetric Matrices (Cont'd)

Given $\vec{a} = [a_x, a_y, a_z]^T$, define

$$S(\vec{a}) = \begin{bmatrix} 0 & -a_z & a_y \\ a_z & 0 & -a_x \\ -a_y & a_x & 0 \end{bmatrix}$$

Property:

$$S(\alpha\vec{a} + \beta\vec{c}) = \alpha S(\vec{a}) + \beta S(\vec{c})$$

Skew Symmetric Matrices (Cont'd)

Given $\vec{a} = [a_x, a_y, a_z]^T$, define

$$S(\vec{a}) = \begin{bmatrix} 0 & -a_z & a_y \\ a_z & 0 & -a_x \\ -a_y & a_x & 0 \end{bmatrix}$$

Property:

$$S(\alpha\vec{a} + \beta\vec{c}) = \alpha S(\vec{a}) + \beta S(\vec{c})$$

Property:

$$S(\vec{a})\vec{p} = \vec{a} \times \vec{p} \quad \text{for any } \vec{a}, \vec{p} \in \mathbb{R}^3$$

Skew Symmetric Matrices (Cont'd)

Given $\vec{a} = [a_x, a_y, a_z]^T$, define

$$S(\vec{a}) = \begin{bmatrix} 0 & -a_z & a_y \\ a_z & 0 & -a_x \\ -a_y & a_x & 0 \end{bmatrix}$$

Property:

$$S(\alpha\vec{a} + \beta\vec{c}) = \alpha S(\vec{a}) + \beta S(\vec{c})$$

Property:

$$S(\vec{a})\vec{p} = \vec{a} \times \vec{p} \quad \text{for any } \vec{a}, \vec{p} \in \mathbb{R}^3$$

$$S(\vec{a})\vec{p} = \begin{bmatrix} 0 & -a_z & a_y \\ a_z & 0 & -a_x \\ -a_y & a_x & 0 \end{bmatrix} \begin{bmatrix} p_x \\ p_y \\ p_z \end{bmatrix} = \begin{bmatrix} p_z a_y - p_y a_z \\ p_x a_z - p_z a_x \\ p_y a_x - p_x a_y \end{bmatrix}$$

Skew Symmetric Matrices (Cont'd)

Property:

$$RS(\vec{a})R^T = S(R\vec{a}) \quad \text{for any } R \in \mathcal{SO}(3), \vec{a} \in \mathbb{R}^3$$

Skew Symmetric Matrices (Cont'd)

Property:

$$R S(\vec{a}) R^T = S(R\vec{a}) \quad \text{for any } R \in \mathcal{SO}(3), \vec{a} \in \mathbb{R}^3$$

$$R [S(\vec{a}) R^T \vec{p}] = R [\vec{a} \times R^T \vec{p}] = R [\vec{a} \times \vec{b}]$$

Skew Symmetric Matrices (Cont'd)

Property:

$$RS(\vec{a})R^T = S(R\vec{a}) \quad \text{for any } R \in \mathcal{SO}(3), \vec{a} \in \mathbb{R}^3$$

$$\begin{aligned} R[S(\vec{a})R^T\vec{p}] &= R[\vec{a} \times R^T\vec{p}] = R[\vec{a} \times \vec{b}] \\ &= R\vec{a} \times R\vec{b} \end{aligned}$$

Skew Symmetric Matrices (Cont'd)

Property:

$$RS(\vec{a})R^T = S(R\vec{a}) \quad \text{for any } R \in \mathcal{SO}(3), \vec{a} \in \mathbb{R}^3$$

$$\begin{aligned} R[S(\vec{a})R^T\vec{p}] &= R[\vec{a} \times R^T\vec{p}] = R[\vec{a} \times \vec{b}] \\ &= R\vec{a} \times R\vec{b} = R\vec{a} \times R(R^T\vec{p}) = R\vec{a} \times \vec{p} \end{aligned}$$

Skew Symmetric Matrices (Cont'd)

Property:

$$RS(\vec{a})R^T = S(R\vec{a}) \quad \text{for any } R \in \mathcal{SO}(3), \vec{a} \in \mathbb{R}^3$$

$$\begin{aligned} R[S(\vec{a})R^T\vec{p}] &= R[\vec{a} \times R^T\vec{p}] = R[\vec{a} \times \vec{b}] \\ &= R\vec{a} \times R\vec{b} = R\vec{a} \times R(R^T\vec{p}) = R\vec{a} \times \vec{p} \\ &= S(R\vec{a})\vec{p} \end{aligned}$$

Skew Symmetric Matrices (Cont'd)

Property:

$$RS(\vec{a})R^T = S(R\vec{a}) \quad \text{for any } R \in \mathcal{SO}(3), \vec{a} \in \mathbb{R}^3$$

$$\begin{aligned} R[S(\vec{a})R^T\vec{p}] &= R[\vec{a} \times R^T\vec{p}] = R[\vec{a} \times \vec{b}] \\ &= R\vec{a} \times R\vec{b} = R\vec{a} \times R(R^T\vec{p}) = R\vec{a} \times \vec{p} \\ &= S(R\vec{a})\vec{p} \end{aligned}$$

Property:

$$x^T S x = 0 \quad \text{for any } S \in \mathfrak{so}(3), x \in \mathbb{R}^3$$

Derivative of Rotation Matrix

Consider a function of a scalar variable $\theta \rightarrow R(\theta) \in \mathcal{SO}(3)$

Derivative of Rotation Matrix

Consider a function of a scalar variable $\theta \rightarrow R(\theta) \in \mathcal{SO}(3)$

How to compute $\frac{d}{d\theta} R(\theta)$?

Derivative of Rotation Matrix

Consider a function of a scalar variable $\theta \rightarrow R(\theta) \in \mathcal{SO}(3)$

How to compute $\frac{d}{d\theta} R(\theta)$?

For any θ the matrix $R(\theta)$ is a rotation, then

$$R(\theta)R^T(\theta) = R^T(\theta)R(\theta) = I$$

Derivative of Rotation Matrix

Consider a function of a scalar variable $\theta \rightarrow R(\theta) \in \mathcal{SO}(3)$

How to compute $\frac{d}{d\theta} R(\theta)$?

For any θ the matrix $R(\theta)$ is a rotation, then

$$R(\theta)R^T(\theta) = R^T(\theta)R(\theta) = I$$

Therefore

$$\frac{d}{dt} [R(\theta)R^T(\theta)] = \frac{d}{dt} [R(\theta)] R^T(\theta) + R(\theta) \frac{d}{dt} [R^T(\theta)] = \frac{d}{dt} I = 0$$

Derivative of Rotation Matrix

Consider a function of a scalar variable $\theta \rightarrow R(\theta) \in \mathcal{SO}(3)$

How to compute $\frac{d}{d\theta} R(\theta)$?

For any θ the matrix $R(\theta)$ is a rotation, then

$$R(\theta)R^T(\theta) = R^T(\theta)R(\theta) = I$$

Therefore

$$\frac{d}{dt} [R(\theta)R^T(\theta)] = \underbrace{\frac{d}{dt} [R(\theta)] R^T(\theta)}_{= \textcolor{red}{S}} + R(\theta) \frac{d}{dt} [R^T(\theta)] = \frac{d}{dt} I = 0$$

Derivative of Rotation Matrix

Consider a function of a scalar variable $\theta \rightarrow R(\theta) \in \mathcal{SO}(3)$

How to compute $\frac{d}{d\theta} R(\theta)$?

For any θ the matrix $R(\theta)$ is a rotation, then

$$R(\theta)R^T(\theta) = R^T(\theta)R(\theta) = I$$

Therefore

$$\frac{d}{dt} [R(\theta)R^T(\theta)] = \underbrace{\frac{d}{dt} [R(\theta)] R^T(\theta)}_{= S} + \underbrace{R(\theta) \frac{d}{dt} [R^T(\theta)]}_{= S^T} = \frac{d}{dt} I = 0$$

Derivative of Rotation Matrix

Consider a function of a scalar variable $\theta \rightarrow R(\theta) \in \mathcal{SO}(3)$

How to compute $\frac{d}{d\theta} R(\theta)$?

For any θ the matrix $R(\theta)$ is a rotation, then

$$R(\theta)R^T(\theta) = R^T(\theta)R(\theta) = I$$

Therefore

$$\frac{d}{dt} [R(\theta)R^T(\theta)] = \underbrace{\frac{d}{dt} [R(\theta)] R^T(\theta)}_{= \mathbf{S}} + \underbrace{R(\theta) \frac{d}{dt} [R^T(\theta)]}_{= \mathbf{S}^T} = \frac{d}{dt} I = 0$$

$$\Rightarrow \frac{d}{dt} [R(\theta)] = \mathbf{S} R(\theta), \quad \mathbf{S} \in \mathfrak{so}(3)$$

Derivative of Rotation Matrix (Example 4.2)

If $R(\theta) = R_{z,\theta}$, that is the basic rotation around the axis z , then

$$\mathbf{S} = \frac{d}{d\theta}[R_{z,\theta}] R_{z,\theta}^T = \frac{d}{d\theta} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}^T$$

Derivative of Rotation Matrix (Example 4.2)

If $R(\theta) = R_{z,\theta}$, that is the basic rotation around the axis z , then

$$\begin{aligned} \mathbf{S} &= \frac{d}{d\theta}[R_{z,\theta}] R_{z,\theta}^T = \frac{d}{d\theta} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}^T \\ &= \begin{bmatrix} -\sin \theta & -\cos \theta & 0 \\ \cos \theta & -\sin \theta & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

Derivative of Rotation Matrix (Example 4.2)

If $R(\theta) = R_{z,\theta}$, that is the basic rotation around the axis z , then

$$\begin{aligned} \mathbf{S} &= \frac{d}{d\theta}[R_{z,\theta}] R_{z,\theta}^T = \frac{d}{d\theta} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}^T \\ &= \begin{bmatrix} -\sin \theta & -\cos \theta & 0 \\ \cos \theta & -\sin \theta & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -a_z & a_y \\ a_z & 0 & -a_x \\ -a_y & a_x & 0 \end{bmatrix} = S(\vec{a}) \end{aligned}$$

Derivative of Rotation Matrix (Example 4.2)

If $R(\theta) = R_{z,\theta}$, that is the basic rotation around the axis z , then

$$\begin{aligned} \mathbf{S} &= \frac{d}{d\theta}[R_{z,\theta}] R_{z,\theta}^T = \frac{d}{d\theta} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}^T \\ &= \begin{bmatrix} -\sin \theta & -\cos \theta & 0 \\ \cos \theta & -\sin \theta & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -a_z & a_y \\ a_z & 0 & -a_x \\ -a_y & a_x & 0 \end{bmatrix} = S(\vec{a}) \\ &= S(\vec{k}), \quad \vec{k} = [0, 0, 1]^T \end{aligned}$$

Lecture 6: Kinematics: Velocity Kinematics - the Jacobian

- Skew Symmetric Matrices
- Linear and Angular Velocities of a Moving Frame

Angular Velocity

If $R(t) \in \mathcal{SO}(3)$ is time-varying and has a time-derivative, then

$$\frac{d}{dt}R(t) = \boldsymbol{S}(t)R(t) = \boldsymbol{S}(\boldsymbol{\omega}(t))R(t), \quad \boldsymbol{S}(\cdot) \in \mathfrak{so}(3)$$

Angular Velocity

If $R(t) \in \mathcal{SO}(3)$ is time-varying and has a time-derivative, then

$$\frac{d}{dt}R(t) = S(t)R(t) = S(\omega(t))R(t), \quad S(\cdot) \in \mathfrak{so}(3)$$

The vector $\omega(t)$ will be the

angular velocity

of rotating frame wrt to the fixed frame at time t .

Angular Velocity

If $R(t) \in \mathcal{SO}(3)$ is time-varying and has a time-derivative, then

$$\frac{d}{dt}R(t) = S(t)R(t) = S(\omega(t))R(t), \quad S(\cdot) \in \mathfrak{so}(3)$$

The vector $\omega(t)$ will be the

angular velocity

of rotating frame wrt to the fixed frame at time t .

Consider a point p rigidly attached to a moving frame, then

$$p^0(t) = R_1^0(t) p^1$$

Angular Velocity

If $R(t) \in \mathcal{SO}(3)$ is time-varying and has a time-derivative, then

$$\frac{d}{dt}R(t) = S(t)R(t) = S(\omega(t))R(t), \quad S(\cdot) \in so(3)$$

The vector $\omega(t)$ will be the

angular velocity

of rotating frame wrt to the fixed frame at time t .

Consider a point p rigidly attached to a moving frame, then

$$p^0(t) = R_1^0(t) p^1$$

Differentiating this expression we obtain

$$\begin{aligned} \frac{d}{dt} [p^0(t)] &= \frac{d}{dt} [R_1^0(t)] p^1 = S(\omega(t)) R_1^0(t) p^1 = \omega(t) \times R_1^0(t) p^1 \\ &= \omega(t) \times p^0(t) \end{aligned}$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\} \omega_{i,j}^k(t)$ expressed in k -coordinate frame

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\}^{\omega_{i,j}^k(t)}$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\} \omega_{i,j}^k(t)$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Taking time derivative of the left and right sides we have

$$\frac{d}{dt} R_2^0(t) = \frac{d}{dt} [R_1^0(t)] R_2^1(t) + R_1^0(t) \frac{d}{dt} [R_2^1(t)]$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\}^{\omega_{i,j}^k(t)}$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Taking time derivative of the left and right sides we have

$$\begin{aligned} \frac{d}{dt} R_2^0(t) &= \frac{d}{dt} [R_1^0(t)] R_2^1(t) + R_1^0(t) \frac{d}{dt} [R_2^1(t)] \\ &= S(\omega_{0,2}^0(t)) R_2^0 \end{aligned}$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\}^{\omega_{i,j}^k(t)}$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Taking time derivative of the left and right sides we have

$$\begin{aligned} \frac{d}{dt} R_2^0(t) &= \frac{d}{dt} [R_1^0(t)] R_2^1(t) + R_1^0(t) \frac{d}{dt} [R_2^1(t)] \\ &= S(\omega_{0,2}^0(t)) R_2^0 \\ &= \left[S(\omega_{0,1}^0(t)) R_1^0(t) \right] R_2^1(t) + R_1^0(t) \left[S(\omega_{1,2}^1(t)) R_2^1(t) \right] \end{aligned}$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\}^{\omega_{i,j}^k(t)}$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Taking time derivative of the left and right sides we have

$$\begin{aligned} \frac{d}{dt} R_2^0(t) &= \frac{d}{dt} [R_1^0(t)] R_2^1(t) + R_1^0(t) \frac{d}{dt} [R_2^1(t)] \\ &= S(\omega_{0,2}^0(t)) R_2^0(t) \\ &= S(\omega_{0,1}^0(t)) \underbrace{R_1^0(t) R_2^1(t)}_{= R_2^0(t)} + R_1^0(t) [S(\omega_{1,2}^1(t)) R_2^1(t)] \end{aligned}$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\}^{\omega_{i,j}^k(t)}$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Taking time derivative of the left and right sides we have

$$\begin{aligned} \frac{d}{dt} R_2^0(t) &= \frac{d}{dt} [R_1^0(t)] R_2^1(t) + R_1^0(t) \frac{d}{dt} [R_2^1(t)] \\ &= S(\omega_{0,2}^0(t)) R_2^0(t) \\ &= S(\omega_{0,1}^0(t)) R_2^0(t) + R_1^0(t) S(\omega_{1,2}^1(t)) \underbrace{R_1^0(t)^T R_1^0(t)}_{= I} R_2^1(t) \end{aligned}$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\} \omega_{i,j}^k(t)$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Taking time derivative of the left and right sides we have

$$\begin{aligned} \frac{d}{dt} R_2^0(t) &= \frac{d}{dt} [R_1^0(t)] R_2^1(t) + R_1^0(t) \frac{d}{dt} [R_2^1(t)] \\ &= S(\omega_{0,2}^0(t)) R_2^0(t) \\ &= S(\omega_{0,1}^0(t)) R_2^0(t) + \underbrace{R_1^0(t) S(\omega_{1,2}^1(t)) R_1^0(t)^T}_{= S(R_1^0(t) \omega_{1,2}^1(t))} R_1^0(t) R_2^1(t) \end{aligned}$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\}^{\omega_{i,j}^k(t)}$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Taking time derivative of the left and right sides we have

$$\begin{aligned} \frac{d}{dt} R_2^0(t) &= \frac{d}{dt} [R_1^0(t)] R_2^1(t) + R_1^0(t) \frac{d}{dt} [R_2^1(t)] \\ &= S(\omega_{0,2}^0(t)) R_2^0(t) \\ &= S(\omega_{0,1}^0(t)) R_2^0(t) + S(R_1^0(t) \omega_{1,2}^1(t)) \underbrace{R_1^0(t) R_2^1(t)}_{= R_2^0(t)} \end{aligned}$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\} \omega_{i,j}^k(t)$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Taking time derivative of the left and right sides we have

$$\begin{aligned} \frac{d}{dt} R_2^0(t) &= \frac{d}{dt} [R_1^0(t)] R_2^1(t) + R_1^0(t) \frac{d}{dt} [R_2^1(t)] \\ &= S(\omega_{0,2}^0(t)) R_2^0 \\ &= \left[S(\omega_{0,1}^0(t)) + S(R_1^0(t) \omega_{1,2}^1(t)) \right] R_2^0(t) \end{aligned}$$

Addition of Angular Velocities

An angular velocity is a free vector

it corresponds to $\left\{ \frac{d}{dt} R_j^i(t) \right\} \omega_{i,j}^k(t)$ expressed in k -coordinate frame

Consider two moving frames and the fixed one all with the common origin

$$R_2^0(t) = R_1^0(t) R_2^1(t)$$

Taking time derivative of the left and right sides we have

$$\begin{aligned} \frac{d}{dt} R_2^0(t) &= \frac{d}{dt} [R_1^0(t)] R_2^1(t) + R_1^0(t) \frac{d}{dt} [R_2^1(t)] \\ &= S(\omega_{0,2}^0(t)) R_2^0 \\ &= \left[S(\omega_{0,1}^0(t)) + S(R_1^0(t) \omega_{1,2}^1(t)) \right] R_2^0(t) \end{aligned}$$

$$\Rightarrow S(\omega_{0,2}^0(t)) = S(\omega_{0,1}^0(t)) + S(R_1^0(t) \omega_{1,2}^1(t))$$

Addition of Angular Velocities (Cont'd)

The relation

$$S(\omega_{0,2}^0(t)) = S(\omega_{0,1}^0(t)) + S(R_1^0(t)\omega_{1,2}^1(t))$$

together with the property: $S(a) + S(c) = S(a + c)$, imply that the angular velocity can be computed as

$$\omega_{0,2}^0(t) = \omega_{0,1}^0(t) + R_1^0(t)\omega_{1,2}^1(t) = \omega_{0,1}^0(t) + \omega_{1,2}^0(t)$$

Addition of Angular Velocities (Cont'd)

The relation

$$S(\omega_{0,2}^0(t)) = S(\omega_{0,1}^0(t)) + S(R_1^0(t)\omega_{1,2}^1(t))$$

together with the property: $S(a) + S(c) = S(a + c)$, imply that the angular velocity can be computed as

$$\omega_{0,2}^0(t) = \omega_{0,1}^0(t) + R_1^0(t)\omega_{1,2}^1(t) = \omega_{0,1}^0(t) + \omega_{1,2}^0(t)$$

Given n -moving frames with the same origins as for fixed one

$$R_n^0(t) = R_1^0(t)R_2^1(t) \cdots R_n^{n-1}(t) \Rightarrow \frac{d}{dt}R_n^0(t) = S(\omega_{0,n}^0(t))R_n^0(t)$$

$$\begin{aligned}\omega_{0,n}^0(t) &= \omega_{0,1}^0(t) + \omega_{1,2}^0(t) + \omega_{2,3}^0(t) + \cdots + \omega_{n-1,n}^0(t) \\ &= \omega_{0,1}^0 + R_1^0\omega_{1,2}^1 + R_2^0\omega_{2,3}^2 + \cdots + R_{n-1}^0\omega_{n-1,n}^{n-1}\end{aligned}$$

Computing a Linear Velocity of a Point

Given moving and fixed frames related by a homogeneous transform

$$H_1^0(t) = \begin{bmatrix} R_1^0(t) & o_1^0(t) \\ 0 & 1 \end{bmatrix}$$

that is, coordinates of each point of moving frame are

$$p^0(t) = R_1^0(t)p^1 + o_1^0(t)$$

Computing a Linear Velocity of a Point

Given moving and fixed frames related by a homogeneous transform

$$H_1^0(t) = \begin{bmatrix} R_1^0(t) & o_1^0(t) \\ 0 & 1 \end{bmatrix}$$

that is, coordinates of each point of moving frame are

$$p^0(t) = R_1^0(t)p^1 + o_1^0(t)$$

Hence

$$\frac{d}{dt}p^0(t) = \frac{d}{dt} [R_1^0(t)] p^1 + \frac{d}{dt} [o_1^0(t)]$$

Computing a Linear Velocity of a Point

Given moving and fixed frames related by a homogeneous transform

$$H_1^0(t) = \begin{bmatrix} R_1^0(t) & o_1^0(t) \\ 0 & 1 \end{bmatrix}$$

that is, coordinates of each point of moving frame are

$$p^0(t) = R_1^0(t)p^1 + o_1^0(t)$$

Hence

$$\begin{aligned} \frac{d}{dt}p^0(t) &= \frac{d}{dt} [R_1^0(t)] p^1 + \frac{d}{dt} [o_1^0(t)] \\ &= \mathbf{S}(\omega_1^0(t)) R_1^0 p^1 + \frac{d}{dt} [o_1^0(t)] \\ &= \omega_1^0(t) \times (R_1^0 p^1) + \frac{d}{dt} [o_1^0(t)] \end{aligned}$$