

## Problem 1

- The three different functions are as below:

$$J_1(x) = x_1^2 + x_2^2 + x_1 + x_2$$

$$J_2(x) = x_1^2 + 2x_2^2 + x_1 + x_2$$

$$J_3(x) = x_1^2 + 10x_2^2 + x_1 + x_2$$

Tables 1, 2, and 3 shows the visualization of the three different functions  $J_1(x)$ ,  $J_2(x)$ , and  $J_3(X)$  respectively.

- I start from  $[5, 5]$ , then use gradient descent to find out the minimum of the three functions above. The algorithm of gradient descent is:

---

### Algorithm 1 Gradient Descent

---

**Input:** Initial point  $w_0$ , gradient norm tolerance  $\epsilon$

**Output:**  $w_t$

Set  $t = 0$

**while**  $\|\nabla J(w_t)\| \geq \epsilon$  **then do**

$w_{t+1} = w_t - \eta_t \nabla J(w_t)$

$t = t + 1$

**end while**

---

Therefore, we can change the values of  $\epsilon$  and  $\eta$  and see how the trajectory of points using gradient descent is changing. Each red line in the four panels in Tables 1, 2, and 3 is showing the trajectory of points using different  $\epsilon$  and  $\eta$ , with the starting and ending points shown as well.

- This three functions are convex functions, which means if we can find a local minimum by gradient descent algorithm, that point is also the global minimum of that function. See summary results below of the three functions with different values of  $\epsilon$  and  $\eta$  (Please note that I used  $h$  below instead of  $\eta$  in my Python code).

First of all, the variation in tolerance  $\epsilon$  will make a difference in the number of steps needed to get to the minimum. The smaller  $\epsilon$  is, the more expensive it is to get the minimum, but the result is also more accurate.

Secondly, the choice of  $\eta$  is the key in getting the minimum. The three functions differ in their variances as  $\text{var}(J_1) < \text{var}(J_2) < \text{var}(J_3)$ . In tracing the minimum in functions with a

larger variance, which have more spread-out points, thus they have a larger gradient descent at a specific point, compared to the functions with smaller variances. Given the same  $\eta$ , and start from the same point, the step size  $\eta_t \nabla J(w_t)$  is larger, and we need to be very careful about this. That's because with a large step size it is very possible that we miss the minimum by stepping over it. For function  $J_3$ , I first set  $\eta = 0.1$ , and that doesn't give me a converged result at all; instead, it keeps producing very large norm of gradients  $\|\nabla J(w_t)\|$  which never go smaller than my  $\epsilon$  (either 0.1 or 0.01). So for function  $J_3$  I ended up setting  $\eta$  less than 0.01. A smaller  $\eta$  produces smaller steps; although it prevents us from missing the minimum point, but it also takes more steps to reach that point, which is more computational expensive.

J1(x):

Converged in 33 iterations  
 from [ 5. 5.] to [-0.49651396 -0.49651396]  
 J(x) min=-0.499975695062at[-0.49651396 -0.49651396]  
 e=0.01 h=0.1

Converged in 364 iterations  
 from [ 5. 5.] to [-0.4964791 -0.4964791]  
 J(x) min=-0.499975206463at[-0.4964791 -0.4964791]  
 e=0.01 h=0.01

Converged in 23 iterations  
 from [ 5. 5.] to [-0.46753373 -0.46753373]  
 J(x) min=-0.49789188268at[-0.46753373 -0.46753373]  
 e=0.1 h=0.1

Converged in 250 iterations  
 from [ 5. 5.] to [-0.46477252 -0.46477252]  
 J(x) min=-0.497518048899at[-0.46477252 -0.46477252]  
 e=0.1 h=0.01

J2(x):

Converged in 32 iterations  
 from [ 5. 5.] to [-0.49564245 -0.24999958]  
 J(x) min=-0.374981011767at[-0.49564245 -0.24999958]  
 e=0.01 h=0.1

Converged in 347 iterations  
 from [ 5. 5.] to [-0.49503624 -0.2499963 ]  
 J(x) min=-0.374975361039at[-0.49503624 -0.2499963 ]  
 e=0.01 h=0.01

Converged in 22 iterations  
 from [ 5. 5.] to [-0.45941716 -0.2499309 ]  
 J(x) min=-0.373353023794at[-0.45941716 -0.2499309 ]

e=0.1 h=0.1

Converged in 233 iterations  
from [ 5. 5.] to [-0.45033639 -0.24961153]  
J(x) min=-0.372533223559at[-0.45033639 -0.24961153]  
e=0.1 h=0.01

J3(x):  
Converged in 347 iterations  
from [ 5. 5.] to [-0.49503624 -0.05 ]  
J(x) min=-0.274975361066at[-0.49503624 -0.05 ]  
e=0.01 h=0.01

Converged in 3499 iterations  
from [ 5. 5.] to [-0.4950097 -0.05 ]  
J(x) min=-0.274975096906at[-0.4950097 -0.05 ]  
e=0.01 h=0.001

Converged in 233 iterations  
from [ 5. 5.] to [-0.45033639 -0.05 ]  
J(x) min=-0.272533525376at[-0.45033639 -0.05 ]  
e=0.1 h=0.01

Converged in 2348 iterations  
from [ 5. 5.] to [-0.45001109 -0.05 ]  
J(x) min=-0.272501109279at[-0.45001109 -0.05 ]  
e=0.1 h=0.001

- Table 4 is showing the visualization of the objective function

$$J(w) = \frac{\lambda}{2} \|w\|^2 + \sum_i (w^T x_i - y_i)^2$$

with different values of  $\lambda$ 's. The first panel is showing  $\lambda = 0.001$ .

- See the red lines in Table 4 for different trajectories of points with  $\lambda$ 's 0.001, 1.0, 1000.0, 100000.0, 0.0001, and 0.0. Starting and ending points are marked in each panel.
- This function  $J(w)$  is a combination of quadratic functions of  $w$ , we know that quadratic function is a convex function, thus  $J(W)$  is also a convex function, which means if we can find a local minimum by gradient descent algorithm, that point is also the global minimum of that function.

First of all, the choice of  $\epsilon$  determines the accuracy of results, which is similar to the functions  $J_1$ ,  $J_2$ , and  $J_3$  above.




Table 1: Visualization of  $J_1(x)$  and Trajectory of Points Using Gradient Descent

Secondly, the choice of  $\eta$  is determined by the 2,000 data points we have. This big amount of data points makes the step size much larger than the cases in the previous functions  $J_1$ ,  $J_2$ , or  $J_3$ . I tested on several different  $\eta$ 's, and found that  $\eta = 1e^{-5}$  is the optimal value in getting the minimum point in reasonable step numbers (I set the max iteration number to be 1000).

Lastly, the choice of  $\lambda$  makes a difference in the minimum of  $J(w)$  only when  $\lambda$  is huge. From

$$J(w) = \frac{\lambda}{2} \|w\|^2 + \sum_i (w^T x_i - y_i)^2$$

we know that the gradient of this function can be vectorized and put in Python code (see my code for reference):

$$\nabla J(w) = \lambda w + 2X^T(Xw - Y)$$

Here  $X$  is a  $2000 \times 2$  matrix of the 2,000 data points;  $Y$  is a  $1000 \times 1$  vector of the labels (either +1 or -1) of the 2,000 data points. When  $\lambda$  is small, the first part  $\lambda w$  is dominated by




Table 2: Visualization of  $J_2(x)$  and Trajectory of Points Using Gradient Descent

the second part  $2X^T(Xw - Y)$ , as  $X$  and  $Y$  are huge matrices compared to the  $2-d$  vector  $w$ ; therefore in this case  $\lambda$  almost has no impact on the gradient descent. However, when  $\lambda$  gets larger,  $w$  gets significantly important in determining  $\nabla J(w_t)$ . See summary results below of changes of  $J(w)$  with different values of  $\lambda$  (Please note that I used  $h$  below instead of  $\eta$  in my Python code). When  $\lambda = 100000.0$ , the minimum moves from  $[-0.11526332, -0.07674558]$  ( $\lambda = 0.001$ ) to  $[5.94202596e + 17, 5.96931252e + 17]$ .

```

Converged in 230 iterations
from [[ 1.]
 [ 1.]] to [[-0.11526332]
 [-0.07674558]]
J(w) min=1039.21825972at[[-0.11526332]
 [-0.07674558]] e=0.01 h=1e-05 lamb=0.001

```

```

Converged in 230 iterations
from [[ 1.]

```




Table 3: Visualization of  $J_3(x)$  and Trajectory of Points Using Gradient Descent

```
[ 1.] to [[-0.11525755]
[-0.07674949]]
J(w) min=1039.22783777at[[-0.11525755]
[-0.07674949]] e=0.01 h=1e-05lamb=1.0

Converged in 184 iterations
from [[ 1.]
[ 1.]] to [[-0.11046043]
[-0.07970587]]
J(w) min=1048.64285946at[[-0.11046043]
[-0.07970587]] e=0.01 h=1e-05lamb=1000.0
from [[ 1.]
[ 1.]] to [[ 5.94202596e+17]
[ 5.96931252e+17]]
J(w) min=6.67348784196e+40at[[ 5.94202596e+17]
[ 5.96931252e+17]] e=0.01 h=1e-05lamb=100000.0
```

Converged in 230 iterations  
from [[ 1.]  
[ 1.]] to [[-0.11526333]  
[-0.07674558]]  
J(w) min=1039.21825109at[[-0.11526333]  
[-0.07674558]] e=0.01 h=1e-05 lamb=0.0001

Converged in 230 iterations  
from [[ 1.]  
[ 1.]] to [[-0.11526333]  
[-0.07674558]]  
J(w) min=1039.21825013at[[-0.11526333]  
[-0.07674558]] e=0.01 h=1e-05 lamb=0.0




Table 4: Visualization of  $J(w)$  and Trajectory of Points with  $\lambda = 0.001, 1.0, 1000.0, 100000.0, 0.0001$ , and  $0.0$