

STA302 - Lecture 4

Cedric Beaulac

May 16, 2019

Introduction

Today's plan

- ▶ Today we introduce the linear regression model
 - ▶ The classic regression model and its assumption
 - ▶ Estimation via Maximum Likelihood
 - ▶ Inference

The linear regression model

- ▶ We will make a serie of assumptions.
- ▶ These assumptions allow us to give more information about the fitted model.
- ▶ But the results are only valid if the assumptions are respected.
- ▶ Tradeoff we make as statisticians.

The linear regression model

$$y_i = \beta_0 + \beta_1 x_i + e_i$$

- Where e_i is a random variable.

The linear regression model

- ▶ Gauss-Markov assumptions (conditions) :
 - ▶ $E(e_i) = 0$
 - ▶ $Cor(e_i, e_j) = 0 \ i \neq j$
 - ▶ $Var(e_i) = \sigma^2 < \infty \ \forall i$
- ▶ These conditions are sufficient to prove the Gauss-Markov theorem :
- ▶ The best linear unbiased estimator (BLUE) for β 's are given by minimizing the mean square error (last week solution).
- ▶ Here, *best* means lowest variance.
- ▶ Proof : https://en.wikipedia.org/wiki/Gauss-Markov_theorem

The linear regression model

- ▶ Usually we also assume a distribution for e_i .
- ▶ The typical model formulation goes as follow.

The linear regression model

$$y_i = \beta_0 + \beta_1 x_i + e_i$$

- ▶ With e_i are independently distributed $\sim N(0, \sigma^2)$
- ▶ Subtely contain all the Gauss-Markov assumption and more (stronger assumptions).

Distribution implications

- ▶ The model implies y_i is normal distributed given x_i .

$$\begin{aligned}\mathbf{E}(y_i|x_i) &= \mathbf{E}(\beta_0 + \beta_1 x_i + e_i) \\ &= \beta_0 + \beta_1 x_i + \mathbf{E}(e_i) \\ &= \beta_0 + \beta_1 x_i\end{aligned}$$

$$\begin{aligned}\text{Var}(y_i|x_i) &= \text{Var}(\beta_0 + \beta_1 x_i + e_i) \\ &= \text{Var}(e_i) \\ &= \sigma^2\end{aligned}$$

- ▶ Thus we have $y_i|x_i \sim N(\beta_0 + \beta_1 x_i, \sigma^2)$.
- ▶ Or in matrix notation $\mathbf{y}|\mathbf{x} \sim N(\mathbf{X}\beta, I\sigma^2)$.
- ▶ Here we talk about $\mathbf{y}|\mathbf{x}$ because we assume \mathbf{x} to be known values and \mathbf{y} to be random. From now, to lighten the notation we will simply say \mathbf{y} .

Maximum Likelihood

- ▶ Now that we have a distribution for the response y we can estimate the parameters with another technique.
- ▶ Let's do a quick review of Maximum likelihood.

Maximum Likelihood

- ▶ Assuming x a random variable is distributed according to p_θ
- ▶ Given a data set $\{x_i | i \in (1, \dots, n)\}$
- ▶ We would like to produce an estimate $\hat{\theta}$ of the parameter θ .
- ▶ $\hat{\theta}_{MLE}$ is the parameter that maximizes the probability of the observer data set :

$$\hat{\theta}_{MLE} = \operatorname{argmax}_{\theta \in \Theta} p_\theta(x_1, \dots, x_n)$$

- ▶ Again, it is an optimization problem (main problems in modern stats/ML).
- ▶ Optimization and statistics go hand in hand.

Maximum Likelihood

- ▶ The likelihood is a function of θ given a data set \mathbf{x} .
- ▶ If x_i 's are independent, the likelihood decomposes as a product of individual probabilities
$$\mathcal{L}(\theta|\mathbf{x}) = p_{\theta}(x_1, \dots, x_n) = \prod_{i=1}^n p_{\theta}(x_i).$$
- ▶ Since it is complicated to derive a product it is common to look at the log-likelihood : $l(\theta|\mathbf{x}) = \log(p_{\theta}(x_1, \dots, x_n)) = \log(\prod_{i=1}^n p_{\theta}(x_i)) = \sum_{i=1}^n \log(p_{\theta}(x_i)).$
- ▶ Maximizing likelihood and maximizing log-likelihood are equivalent (monotonically increasing function).

Maximum Likelihood

- ▶ In our regression model, we have three parameters : β_0 , β_1 and σ .
- ▶ Let's estimate them with maximum likelihood.
- ▶ Since $y_i|x_i \sim N(\beta_0 + \beta_1 x_i, \sigma^2)$ then :

$$p_{\theta}(y_i|x_i) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y_i - (\beta_0 + \beta_1 x_i))^2}{2\sigma^2}\right)$$

Maximum Likelihood

$$\begin{aligned}l(\theta|x_i) &= \sum_{i=1}^n \log(p_\theta(y_i|x_i)) \\&= \sum_{i=1}^n \log \left(\frac{1}{\sqrt{2\pi}} \frac{1}{\sigma} \exp \left(-\frac{(y_i - (\beta_0 + \beta_1 x_i))^2}{2\sigma^2} \right) \right) \\&= \sum_{i=1}^n -\frac{1}{2} \log(2\pi) - \log(\sigma) - \frac{1}{2\sigma^2} (y_i - (\beta_0 + \beta_1 x_i))^2 \\&= -\frac{n}{2} \log(2\pi) - n \log(\sigma) - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - (\beta_0 + \beta_1 x_i))^2\end{aligned}$$

Maximum Likelihood

- ▶ With matrix notation :

$$l(\theta|x_i) = -\frac{n}{2} \log(2\pi) - n \log(\sigma) - \frac{1}{2\sigma^2} (\mathbf{y} - \mathbf{X}\beta)^T (\mathbf{y} - \mathbf{X}\beta)$$

- Maximizing this term with respect to β is equivalent to minimizing

$$(\mathbf{y} - \mathbf{X}\beta)^T (\mathbf{y} - \mathbf{X}\beta)$$

$$\frac{d}{d\beta} (\mathbf{y} - \mathbf{X}\beta)^T (\mathbf{y} - \mathbf{X}\beta) = 0$$

$$(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y} = \hat{\beta}$$

- ▶ Exactly the same optimization from last week! Incredible!
- ▶ Both minimizing the squared error and maximizing the likelihood leads to the same estimates $\hat{\beta}$!

Maximum Likelihood

$$l(\theta|x_i) = -\frac{n}{2} \log(2\pi) - n \log(\sigma) - \frac{1}{2\sigma^2} (\mathbf{y} - \mathbf{X}\hat{\beta})^T (\mathbf{y} - \mathbf{X}\hat{\beta})$$

- Maximizing this term with respect to σ :

$$\begin{aligned} 0 &= -\frac{n}{\sigma} + \frac{1}{\sigma^3} (\mathbf{y} - \mathbf{X}\hat{\beta})^T (\mathbf{y} - \mathbf{X}\hat{\beta}) \\ \Rightarrow \hat{\sigma}^2 &= \frac{(\mathbf{y} - \mathbf{X}\hat{\beta})^T (\mathbf{y} - \mathbf{X}\hat{\beta})}{n} \\ &= \frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n} \end{aligned}$$

Maximum Likelihood

$$\hat{\sigma}^2 = \frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n}$$

- ▶ This is the mean squared error!
- ▶ Reminder : It is a biased estimator of σ^2 (unbiased would be $\sum_{i=1}^n (y_i - \hat{y}_i)^2 / (n - 2)$).
- ▶ For the moment σ is some kind of nuisance parameters.
- ▶ It is not interesting to us, but must be taking care of to proceed.

Inference

We love statistical models

- ▶ I love statistical models!
- ▶ We now have so much information about $\hat{\beta}$'s and \hat{y} .

Inference

- ▶ Inference is the action of extracting information about parameters given a data set.
- ▶ Good inference procedure considers the variability about the data: how certain are we that this parameter takes this value?

Inference

- ▶ Now that we have a model with a distribution we have more information on our estimate $\hat{\beta}$.
- ▶ Since $\mathbf{y} \sim N(\mathbf{X}\beta, I\sigma^2)$. $\mathbf{E}[\mathbf{y}] = \mathbf{X}\beta$ and $\text{Var}[\mathbf{y}] = I\sigma^2$.
- ▶ Let's use lecture 3 results on \mathbf{E} and \mathbf{V} of random vectors.
- ▶ Since $\hat{\beta} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$, then :

$$\begin{aligned}\mathbf{E}[\hat{\beta}] &= \mathbf{E}[(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}] \\ &= (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{E}[\mathbf{y}] \\ &= (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X} \beta = \beta\end{aligned}$$

- ▶ $\hat{\beta}$ is an unbiased estimator!

Inference

$$\begin{aligned}\text{Var}[\hat{\beta}] &= \text{Var}[(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}] \\&= (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \text{Var}[\mathbf{y} | \mathbf{X}] ((\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T)^T \\&= (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T I \sigma^2 ((\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T)^T \\&= \sigma^2 (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T ((\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T)^T \\&= \sigma^2 (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T (\mathbf{X} ((\mathbf{X}^T \mathbf{X})^{-1})^T) \\&= \sigma^2 (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T (\mathbf{X} ((\mathbf{X}^T \mathbf{X})^T)^{-1}) \\&= \sigma^2 (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X} (\mathbf{X}^T \mathbf{X})^{-1} \\&= \sigma^2 (\mathbf{X}^T \mathbf{X})^{-1}\end{aligned}$$

Inference

IMPORTANT result from distribution theory if \mathbf{z} is multivariate normal : $\mathbf{z} \sim N(\mu, \Sigma)$ and if $\mathbf{w} = \mathbf{c} + \mathbf{B}\mathbf{z}$ then :

$$\mathbf{w} \sim N(\mathbf{c} + \mathbf{B}\mu, \mathbf{B}\Sigma\mathbf{B}^T)$$

- ▶ Remember : $\hat{\beta} = (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{y}$.
- ▶ We also know $\mathbf{y} \sim N(\mathbf{X}\beta, I\sigma^2)$.
- ▶ We can compute the distribution of $\hat{\beta}$.

Inference

$$\begin{aligned}\hat{\beta} &\sim N((\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X} \beta, (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \sigma^2 ((\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T)^T) \\ \Rightarrow \hat{\beta} &\sim N(\beta, (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T (\mathbf{X} (\mathbf{X}^T \mathbf{X})^{-1}) \sigma^2) \\ \Rightarrow \hat{\beta} &\sim N(\beta, (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X} (\mathbf{X}^T \mathbf{X})^{-1} \sigma^2) \\ \Rightarrow \hat{\beta} &\sim N(\beta, (\mathbf{X}^T \mathbf{X})^{-1} \sigma^2)\end{aligned}$$

- ▶ To begin, notice that since $\mathbf{E}(\hat{\beta}) = \beta$ we say the estimator is **unbiased**, this is an important property of this estimator (that's great!).
- ▶ Remember that by Gauss-Markov theorem it is BLUE (Best Linear UNBIASED estimator), i.e. amongs the unbiased linear estimator it has minimal variance. We just showed it unbiased!

Inference

- ▶ With a distribution we can also build confidence interval (small reminder : it is an interval of plausible values).
- ▶ We can then do inference; to determine if a parameters is statistically significant.
- ▶ Let's take a look at β_0 and β_1 separately.

Inference

$$\begin{aligned}\mathbf{X}^T \mathbf{X} &= \begin{bmatrix} 1 & \cdot & \cdot & 1 \\ x_1 & \cdot & \cdot & x_n \end{bmatrix} \begin{bmatrix} 1 & x_1 \\ \cdot & \cdot \\ \cdot & \cdot \\ 1 & x_n \end{bmatrix} \\ &= \begin{bmatrix} \sum_{i=1}^n 1^2 & \sum_{i=1}^n x_i \\ \sum_{i=1}^n x_i & \sum_{i=1}^n x_i^2 \end{bmatrix} = n \begin{bmatrix} 1 & \bar{x} \\ \bar{x} & \frac{1}{n} \sum_{i=1}^n x_i^2 \end{bmatrix}\end{aligned}$$

Inference

$$\begin{aligned}(\mathbf{X}^T \mathbf{X})^{-1} &= \frac{1}{\det \begin{bmatrix} \frac{1}{n} \sum_{i=1}^n x_i^2 & -\bar{x} \\ -\bar{x} & 1 \end{bmatrix}} \\&= \frac{1}{n(1/n \sum_{i=1}^n (x_i^2 - (\bar{x})^2))} \begin{bmatrix} \frac{1}{n} \sum_{i=1}^n x_i^2 & -\bar{x} \\ -\bar{x} & 1 \end{bmatrix} \\&= \frac{1}{\sum_{i=1}^n (x_i - \bar{x})^2} \begin{bmatrix} \frac{1}{n} \sum_{i=1}^n x_i^2 & -\bar{x} \\ -\bar{x} & 1 \end{bmatrix}\end{aligned}$$

Inference

$$\begin{aligned}\hat{\beta} &\sim N(\beta, (\mathbf{X}^T \mathbf{X})^{-1} \sigma^2) \\ \Rightarrow \hat{\beta}_1 &\sim N\left(\beta_1, \sigma^2 \frac{1}{SSX}\right)\end{aligned}$$

where $SSX = \sum_{i=1}^n (x_i - \bar{x})^2$.

► $\hat{\beta}_0$ is left as an exercise.

Inference for β_1

- ▶ Since $\hat{\beta}_1 \sim N\left(\beta_1, \sigma^2 \frac{1}{SSX}\right)$, we have

$$\frac{\hat{\beta}_1 - \beta_1}{\sigma / \sqrt{SSX}} \sim N(0, 1)$$

- ▶ Does our predictor have a significant effect ?
- ▶ Remember $E[y_i] = \beta_0 + \beta_1 x_i$.
- ▶ If the parameter is non-zero then x affect y .
- ▶ In other word, if $\hat{\beta}_1$ is non-zero, as x moves around the expectation of y changes.
- ▶ Thus knowing the value of x helps us at better predicting y .

Inference for β_1

- ▶ Since $\hat{\beta}_1 \sim N\left(\beta_1, \sigma^2 \frac{1}{SSX}\right)$, we have

$$\frac{\hat{\beta}_1 - \beta_1}{\sigma / \sqrt{SSX}} \sim N(0, 1)$$

- ▶ Because we don't know σ , we use $\hat{\sigma} = s$ which leads to :

$$\frac{\hat{\beta}_1 - \beta_1}{s / \sqrt{SSX}} \sim t_{n-2}$$

Inference for β_1

- ▶ The null is The parameter has no effect : $\beta_1 = 0$
- ▶ Then under the null

$$\frac{\hat{\beta}_1}{s/\sqrt{SSX}} \sim t_{n-2}$$

- ▶ If the p-value is small, this is evidence against the null.

Confidence interval

- ▶ If $z \sim N(0, 1)$ then $P(-1.96 < z < 1.96) = 0.95$.
- ▶ Confidence interval are based on this concept.
- ▶ If $z \sim N(\mu, \sigma^2)$ then

$$P(-1.96 < \frac{z - \mu}{\sigma} < 1.96) = 0.95$$

$$\Rightarrow P(-1.96\sigma < z - \mu < 1.96\sigma) = 0.95$$

$$\Rightarrow P(-1.96\sigma - z < -\mu < 1.96\sigma - z) = 0.95$$

$$\Rightarrow P(z - 1.96\sigma < \mu < z + 1.96\sigma) = 0.95$$

Confidence interval

- ▶ We say a confidence interval for μ is $(z - 1.96\sigma, z + 1.96\sigma)$
- ▶ or $CI_{0.95}(\mu) = z \pm 1.96\sigma$.
- ▶ This is a confidence interval for the unknown value of μ the true parameters.
- ▶ For $x \in CI_{0.95}(\mu)$ we would accept the null $H_0 : \mu = x$ with a significance level $1 - 0.95$.

Inference for β_1

- We have $\hat{\beta}_1 \sim N\left(\beta_1, \sigma^2 \frac{1}{SSX}\right)$:

$$\frac{\hat{\beta}_1 - \beta_1}{\sigma/\sqrt{SSX}} \sim N(0, 1)$$

$$P(-1.96 < \frac{\hat{\beta}_1 - \beta_1}{\sigma/\sqrt{SSX}} < 1.96) = 0.95$$

$$\Rightarrow P(-1.96\sigma/\sqrt{SSX} < \hat{\beta}_1 - \beta_1 < 1.96\sigma/\sqrt{SSX}) = 0.95$$

$$\Rightarrow P(-1.96\sigma/\sqrt{SSX} - \hat{\beta}_1 < -\beta_1 < 1.96\sigma/\sqrt{SSX} - \hat{\beta}_1) = 0.95$$

$$\Rightarrow P(\hat{\beta}_1 - 1.96\sigma/\sqrt{SSX} < \beta_1 < \hat{\beta}_1 + 1.96\sigma/\sqrt{SSX}) = 0.95$$

Inference for β_1

- ▶ If σ is unknown. We estimate with $s = \sum_{i=1}^n (y_i - \hat{y}_i)^2 / (n - 2)$
- ▶ Then we have :

$$\frac{\hat{\beta}_1 - \beta_1}{s / \sqrt{SSX}} \sim t_{n-2}$$

Inference for β_1

- This leads to the following confidence interval for β_1 for known σ :

$$(\hat{\beta}_1 - 1.96\sigma/\sqrt{SSX}, \hat{\beta}_1 + 1.96\sigma/\sqrt{SSX})$$

and for unknown σ :

$$(\hat{\beta}_1 - t_{n-2}(\alpha/2)s/\sqrt{SSX}, \hat{\beta}_1 + t_{n-2}(\alpha/2)s/\sqrt{SSX})$$

Conclusion

- ▶ By defining a model and assuming distributions we now have a model that allows us to establish an entire interval of possible values.
- ▶ We know exactly how *sure* we are for the proposed values for $\hat{\beta}_1$.
- ▶ We can then claim if predictor x has a statistically significant effect on y .
- ▶ But what if our assumptions are violated ? Our results are less trustworthy.
- ▶ That's the next topic we will cover.

Practice problem

- ▶ Find the expectation, variance and distribution for $\hat{\beta}_0$.
- ▶ A Modern Approach to Regression with R ch.2 : 4,5,7 solutions (here)
- ▶ Alison Gibbs' additional chapter 2 practice problems (everything except 3) (here)
- ▶ A Modern Approach to Regression with R ch.2 : 1(a,b) (do with R)

External Sources

- ▶ A Modern Approach to Regression with R ch.2
- ▶ A Modern Approach to Regression with R ch.5

Test1

- ▶ Rooms : A-R BA1160, S-Y BA1170, Z BA2135 (first letter on your student ID)
- ▶ Duration : 3 Hours (ish, 2h50 minutes to be precise). I think it is a 1h30-2h00 test.
- ▶ Types of questions : Some conceptual questions, some math (expectation an such) and some R output.
- ▶ Topics :
 - ▶ p-values, statistical significance and hypothesis testing.
 - ▶ ANOVA
 - ▶ Linear Regression