

MATRIX ANALYSIS

A Quick Guide

Huan Bui

Colby College
Physics & Statistics
Class of 2021

February 17, 2019

Preface

Greetings,

Matrix Analysis: A Quick Guide to is compiled based on my MA353: Matrix Analysis notes with professor Leo Livshits. The sections are based on a number of resources: *Linear Algebra Done Right* by Axler, *A Second Course in Linear Algebra* by Horn and Garcia, *Matrices and Linear Transformations* by Cullen, *Matrices: Methods and Applications* by Barnett, *Problems and Theorems in Linear Algebra* by Prasolov, *Matrix Operations* by Richard Bronson, and professor Leo Livshits' own textbook (in the making). Prerequisites: some prior exposure to a first course in linear algebra.

The development of this text will come in layers. The first layer, one that I am working on during the course of S'19 MA353, will be an overview of the key topics listed in the table of contents. As the semester progresses, I will be constantly updating the existing notes, as well as adding prof. Livshits' problems and my solutions to the problems. The second layer will come after the course is over, when concepts will have hopefully "come together."

I will decide how much narrative I should put into the text as text is developed over the semester. I'm thinking that I will only add detailed explanations wherever I find fit or necessary for my own studies. I will most likely keep the text as condensed as I can.

Enjoy!

Contents

Preface	1
1 List of Special Matrices & Their Properties	5
2 List of Operations	6
3 List of Algorithms	7
4 Complex Numbers	8
4.1 A different point of view	8
4.2 Relevant properties and definitions	10
5 Vector Spaces & Linear Functions	12
5.1 Review of Linear Spaces and Subspaces	12
5.2 Review of Linear Maps	12
5.3 Review of Kernels and Images	13
5.4 Atrices	13
5.5 Linear Independence, Span, and Bases	15
5.5.1 Linear Independence	15
5.5.2 Span	16
5.5.3 Bases	16
5.6 Linear Bijections and Isomorphisms	17
5.7 Finite-Dimensional Linear Spaces	17
5.7.1 Dimension	17
5.7.2 Rank-Nullity Theorem	18
5.7.3 Coordinatization & matricial representation of linear func- tions	18
5.8 Infinite-Dimensional Spaces	18
6 Sums of subspaces & Products of vector spaces	20
6.1 Direct Sums	20
6.2 Products and Quotients of Vector Spaces	20
6.3 Products of Vector Spaces	20
6.4 Products & Direct Sums	20
6.5 Quotients of Vector Spaces	21
6.6 Nullspaces	21
6.7 Ranges of Operator Powers	21
7 Idempotents & Resolutions of Identity	22
8 Block-representations of operators	23
8.1 Direct sums of operators	23
9 Invariant subspaces	24
9.1 Reducing subspaces	24

10 Polynomials applied to operators	25
10.1 Minimal polynomials of block- Δ^r operators	25
10.2 Minimal polynomials at a vector	25
11 Eigentheory	26
11.1 Spectral Mapping Theorem	26
12 Triangularization	27
12.1 Compression to invariant subspaces	27
12.2 Simultaneously Δ -ity of commuting families	27
13 Diagonalization	28
13.1 Spectral resolutions	28
13.2 Compressions to reducing subspaces	28
13.3 Simultaneous diagonalizability for commuting families	28
14 Primary decomposition over \mathbb{C} and generalized eigenspaces	29
15 Cyclic decomposition and Jordan form	30
15.1 Square roots of operators	30
15.2 Similarity of a matrix and its transpose	30
15.3 Similarity of a matrix and its conjugate	30
15.4 Jordan forms of \mathcal{AB} and \mathcal{BA}	30
15.5 Power-convergent operators	30
15.6 Power-bounded operators	30
15.7 Row-stochastic matrices	30
16 Determinant & Trace	31
16.1 Classical adjoints	31
16.2 Cayley-Hamilton theorem	31
17 Inner products and norms	32
17.1 Riesz representation theorem	33
17.2 Adjoints	33
17.3 Grammians	33
17.4 Orthogonal complements and orthogonal decompositions	33
17.5 Ortho-projections	33
17.6 Closest point solutions	33
17.7 Gram-Schmidt and orthonormal bases	33
18 Isometries and unitary operators	34
19 Ortho-triangularization	35
20 Spectral resolutions	36

21 Ortho-diagonalization; self-adjoint and normal operators; spectral theorems	37
22 Positive (semi-)definite operators	38
22.1 Classification of inner products	38
22.2 Positive square roots	38
23 Polar decomposition	39
24 Single value decomposition	40
24.1 Spectral/operator norm	40
24.2 Singular values and approximation	40
24.3 Singular values and eigenvalues	40
25 Problems and Solutions	41
25.1 Problem set 1	41
25.2 Problem set 2	50
25.3 Problem set 3	51

1 List of Special Matrices & Their Properties

1. **Hermitian/Self-adjoint:** $H = H^\dagger$. A Hermitian matrix is matrix that is equal to its own conjugate transpose:

$$H \text{ is Hermitian} \iff H_{ij} = \bar{H}_{ji}$$

Proposition 1.1.

- (a) H is Hermitian $\iff \langle w, Hv \rangle = \langle Hw, v \rangle$, where \langle, \rangle denotes the inner product.
- (b) H is Hermitian $\iff \langle v, Hv \rangle \in \mathbb{R}$.
- (c) H is Hermitian \iff it is *unitarily diagonalizable* with *real eigenvalues*.

Unitary: $U^*U = UU^* = I = U^\dagger U = UU^\dagger$. The real analogue of a unitary matrix is an orthogonal matrix. The following list contain the properties of U :

- (a) U preserves the inner product:

$$\langle Ux, Uy \rangle = \langle x, y \rangle.$$

- (b) U is normal: it commutes with $U^* = U^\dagger$.
- (c) U is diagonalizable:

$$U = VDV^*,$$

where D is diagonal and unitary, and V is unitary.

- (d) $|\det(U)| = 1$ (hence the real analogue to U is an orthogonal matrix)
- (e) Its eigenspaces are orthogonal.
- (f) U can be written as

$$U = e^{iH},$$

where H is a Hermitian matrix.

- (g) Any square matrix with unit Euclidean norm is the average of two unitary matrices.

2. **Cofactor:** ?

2 List of Operations

1. **Conjugate transpose** is what its name suggests.
2. **Classical adjoint/Adjugate/adjunct** of a square matrix is the transpose of its cofactor matrix.

3 List of Algorithms

4 Complex Numbers

4.1 A different point of view

We often think of complex numbers as

$$a + ib$$

where $a, b \in \mathbb{R}$ and $i = \sqrt{-1}$. While there is nothing “bad” about this way of thinking - in fact thinking of complex numbers as $a + ib$ allows us to very quickly and intuitively do arithmetics operations on them - a “matrix representation” of complex numbers can give us some insights on “what we actually do” when we perform complex arithmetics.

Let us think of

$$\begin{pmatrix} a & -b \\ b & a \end{pmatrix}$$

as a different representation of the same object - the same complex number “ $a + ib$.” Note that it does not make sense to say the matrix representation **equals** the complex number itself. But we shall see that a lot of the properties of complex numbers are carried into this matrix representation under interesting matricial properties.

$$\boxed{\begin{pmatrix} a & -b \\ b & a \end{pmatrix} \sim a + ib}$$

First, let us break the the matrix down:

$$a + ib = a \times 1 + i \times b \sim \begin{pmatrix} a & -b \\ b & a \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + b \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = aI + b\mathcal{I}.$$

Right away, we can make some “mental connections” between the representations:

$$\begin{aligned} I &\sim 1 \\ \mathcal{I} &\sim i. \end{aligned}$$

Now, we know that complex number multiplications commute:

$$(a + ib)(c + id) = (c + id)(a + ib).$$

Matrix multiplications are not commutative. So, we might wonder whether commutativity holds under the this new representation of complex numbers. Well, the answer is yes. We can readily verify that

$$(aI + b\mathcal{I})(cI + b\mathcal{I}) = (cI + b\mathcal{I})(aI + b\mathcal{I}).$$

How about additions? Let's check:

$$(a + ib) + (c + id) = (a + c) + i(b + d) \sim \begin{pmatrix} a + c & -(b + d) \\ (b + d) & a + c \end{pmatrix} = \begin{pmatrix} a & -b \\ b & a \end{pmatrix} + \begin{pmatrix} c & -d \\ d & c \end{pmatrix}.$$

Ah! Additions work. So, the new representation of complex numbers seems to be working flawlessly. However, we have yet to gain any interesting insights into the connections between the representations. To do that, we have to look into changing the form of the matrix. First, let's see what conjugation does:

$$(a + ib)^* = a - ib \sim \begin{pmatrix} a & b \\ -b & a \end{pmatrix} = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}^\top$$

Ah, so conjugation to a complex number in the traditional representation is the same as transposition in the matrix representations. What about the amplitude square? Let us call

$$M = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}.$$

We have

$$(a + ib)(a - ib) \sim \begin{pmatrix} a & -b \\ b & a \end{pmatrix} \begin{pmatrix} a & b \\ -b & a \end{pmatrix} = MM^\top = (a^2 + b^2)I = \det(M)I$$

Interesting. But observe that if $\det(M) \neq 0$

$$\frac{1}{\det(M)} MM^\top = I.$$

This tells us that

$$M^\top = M^{-1},$$

where M^{-1} is the inverse of M , and, not surprisingly, it corresponds to the reciprocal to the complex number $a + ib$. We can readily show that

$$M^{-1} \sim (a + ib)^{-1} = \frac{1}{a^2 + b^2}(a - ib).$$

Remember that we can also think of a complex number as a column vector:

$$c + id \sim \begin{pmatrix} c \\ d \end{pmatrix}.$$

Let us look back at complex number multiplication under matrix representation:

$$(a + ib)(c + id) = (ac - bd) + i(bc + ad) \sim \begin{pmatrix} a & -b \\ b & a \end{pmatrix} \begin{pmatrix} c \\ d \end{pmatrix} = \begin{pmatrix} ac - bd \\ bc + ad \end{pmatrix}.$$

Multiplication actually works in this “mixed” way of representing complex numbers as well. Now, observe that what we just did was performing a linear transformation on a vector in \mathbb{R}^2 . It is always interesting to look at the geometrical interpretation of this transformation. To do this, let us call N the “normalized” version of M :

$$N = \frac{1}{\sqrt{a^2 + b^2}} \begin{pmatrix} a & -b \\ b & a \end{pmatrix}.$$

We immediately recognize that N is an orthogonal matrix. This means N is an orthogonal transformation (length preserving). Now, it is reasonable to define

$$\begin{aligned} \cos \theta &= \frac{a}{\sqrt{a^2 + b^2}} \\ \sin \theta &= \frac{b}{\sqrt{a^2 + b^2}}. \end{aligned}$$

We can write N as

$$N = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix},$$

which is a physicists’ favorite matrix: the rotation by θ . So, let us write M in terms of N :

$$M = \begin{pmatrix} a & -b \\ b & a \end{pmatrix} = \sqrt{a^2 + b^2} N = \sqrt{a^2 + b^2} \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

We can interpret M as a rotation by θ , followed by a scaling by $\sqrt{a^2 + b^2}$. But what $\sqrt{a^2 + b^2}$ exactly is just the “length” or the “amplitude” of the complex number $a + ib$, if we think of it as an arrow in a plane.

4.2 Relevant properties and definitions

1. The *modulus* of $z = a + ib$ is the “amplitude” of z , denoted by $|z| = \sqrt{a^2 + b^2} = z\bar{z}$.
2. The modulus is *multiplicative*, i.e.

$$|wz| = |w||z|.$$

3. Triangle inequality:

$$|z + w| \leq |z| + |w|.$$

We can readily show this geometrically, or algebraically.

4. The *argument* of $z = a + ib$ is θ , where

$$\theta = \begin{cases} \tan^{-1} \left(\frac{b}{a} \right), & \text{if } a > 0 \\ \frac{\pi}{2} + k2\pi, k \in \mathbb{R} & \text{if } a = 0, b > 0 \\ -\frac{\pi}{2} + k2\pi, k \in \mathbb{R} & \text{if } a = 0, b < 0 \\ \text{Undefined} & \text{if } a = b = 0. \end{cases}$$

5. The *conjugate* of $a+ib$ is $a-ib$. Conjugation is *additive* and *multiplicative*, i.e.

$$\begin{aligned}z + w &= \bar{z} + \bar{w} \\ \bar{w}z &= \bar{w}\bar{z}.\end{aligned}$$

Note that we can also show the multiplicative property with the matrix representation as well:

$$\bar{w}z \sim (WZ)^\top = Z^\top W^\top \sim \bar{z}\bar{w} = \bar{w}\bar{z}.$$

6. Euler's identity, generalized to de Moivre's formula:

$$z^n = r^n e^{in\theta}.$$

5 Vector Spaces & Linear Functions

5.1 Review of Linear Spaces and Subspaces

Proposition 5.1. of linear spaces:

1. Commutativity and associativity of addition
2. Existence of an additively neutral element (null element). Zero multiples of elements give the null element: $0 \cdot V = \mathbf{0}$
3. Every element has an (unique) additively antipodal element
4. Scalar multiplication distributes over addition
5. Multiplicative identity:
6. $ab \cdot V = a \cdot (bV)$
7. $(a + b)V = aV + bV$

W is a subspace of V if

1. $S \subseteq V$
2. S is non-empty
3. S is closed under addition and scalar multiplication

Proposition 5.2. that are interesting/important/maybe-not-so-obvious:

1. If S is a subspace of V and $S \neq V$ then S is a proper subspace of V .
2. If X is a subspace of Y and Y is a subspace of Z , then X is a subspace of Z .
3. Non-trivial linear (non-singleton) spaces are infinite.

5.2 Review of Linear Maps

Consider linear spaces V and W and elements $v \in V$ and $w \in W$ and scalars $\alpha, \beta \in \mathbb{R}$, a function $F : V \rightarrow W$ is a linear map if

$$F[\alpha v + \beta w] = \alpha F[v] + \beta F[w].$$

Proposition 5.3.

1. $F[\mathbf{0}_V] = \mathbf{0}_W$
2. $G[w] = (\alpha \cdot F)[w] = \alpha \cdot F[w]$
3. Given $F : V \rightarrow W$ and $G : V \rightarrow W$, $H[v] = F[v] + G[v] = (F + G)[v]$ is called the sum of the functions F and G .

4. Linear combinations of linear maps are linear.
5. Compositions of linear maps are linear.
6. Compositions distributes over linear combinations of linear maps.
7. Inverses of linear functions (if they exist) are linear.
8. Inverse of a bijective linear function is a bijective linear function

5.3 Review of Kernels and Images

Definition 5.1. Let $F : V \rightarrow W$ be given. The kernel of F is defined as

$$\ker(F) = \{v \in V \mid F[v] = \mathbf{0}_W\}.$$

Proposition 5.4. Let $F : V \rightarrow W$ a linear map be given. Also, consider a linear map G such that $F \circ G$ is defined

1. F is null $\iff \ker(F) = V \iff \text{Im}(F) = \mathbf{0}_W$
2. $\ker(F)$ is a subspace of V
3. $\text{Im}(F)$ is a subspace of W
4. F is injective $\iff \ker(F) = \mathbf{0}_V$
5. $\ker(F) \subseteq \ker(F \circ G)$.
6. F injective $\implies \ker(F) = \ker(F \circ G)$

Definition 5.2. Let $F : V \rightarrow W$ be given. The image of F is defined as

$$\text{Im}(F) = \{w \in W \mid \exists v \in V, F[v] = w\}.$$

5.4 Atrices

Definition 5.3. Atrix functions: Let $V_1, \dots, V_m \in \mathbf{V}$ be given. Consider $f : \mathbb{R}^m \rightarrow \mathbf{V}$ be defined by

$$f \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{pmatrix} = \sum_{i=1}^m a_i V_i.$$

We denote f by

$$(V_1 \quad V_2 \quad \dots \quad V_m).$$

We refer to the V_i 's as the **columns** of f , even though there doesn't have to be any columns. Basically, f is simply a function that takes in an ordered list of coefficients and returns a linear combination of V_i with the respective coefficients. A matrix is a special atrix. Not every atrix is a matrix.

Proposition 5.5. of atrices

1. The V_i 's - the columns of an atrix - are the images of the standard basis tuples.
2. $\mathbf{e}_j \in \ker(V_1 \dots V_m) \iff V_j = \mathbf{0}_V$, where \mathbf{e}_j denotes a standard basis tuple with a 1 at the j^{th} position. To put in words, a kernel of an atrix contains a standard basis if and only if one of its columns is a null element.
3. $\text{Im}(f) \equiv \text{Im}(V_1 \dots V_m) = \text{span}(V_1 \dots V_m)$
4. B is a null atrix $\iff \ker(B) = \mathbb{R}^m \iff \text{Im}(B) = \mathbf{0}_V \iff V_j = \mathbf{0}_V \forall j = 1, 2, \dots, m$.
5. f is a linear function $\mathbb{R}^m \rightarrow V \iff f$ is an atrix function $\mathbb{R}^m \rightarrow V$.
6. An atrix A is bijective/invertible, then its inverse A^{-1} is a linear function, but is an atrix only if A is a matrix.
7. Linear combinations of atrices are atrices:

$$\begin{aligned} & \alpha \cdot (V_1 \quad \dots \quad V_m) + \beta \cdot (W_1 \quad \dots \quad W_m) \\ &= (\alpha V_1 + \beta W_1 \quad \dots \quad \alpha V_m + \beta W_m) \end{aligned}$$

8. Compositions of two atrices are NOT defined unless the atrix going first is a matrix. Consider $F : \mathbb{R}^m \rightarrow W$ and $G : \mathbb{R}^n \rightarrow T$. $F \circ G$ is only defined if $T = \mathbb{R}^m$. This makes G an $n \times m$ matrix. It follows that the atrix $F \circ G$ has the form

$$(f(g_1) \quad f(g_2) \quad \dots \quad f(g_m)).$$

9. Consider $F : \mathbb{R}^m \rightarrow V$ and $G : \mathbb{R}^k \rightarrow V$. $\text{Im}(F) \subseteq \text{Im}(G) \iff F = G \circ C$, with $C \in \mathbb{M}_{k \times m}$, i.e. C is an $k \times m$ matrix.
10. Consider an atrix $A : \mathbb{R}^m \rightarrow V$. $A = (V_1 \dots V_m)$. A is NOT injective.
 - $\iff \exists$ a non-trivial linear combination of the columns of A that gives $\mathbf{0}_V$
 - $\iff A = [\mathbf{0}_v]$ or $\exists j | V_j$ is linear combination of other columns of A
 - \iff The first column of A is $\mathbf{0}_V$ or $\exists j | V_j$ is a linear combination some of $V_i, i < j$.
11. If atrix $A : \mathbb{R} \rightarrow V$ has a single column then it is injective if the column is not $\mathbf{0}_V$.

Proposition 5.6. of elementary column operations for atrices. Elementary operations on the columns of F can be expressed as a composition of F and an appropriate elementary matrix E , $F \circ E$.

1. Swapping i^{th} and j^{th} columns: $F \circ E^{[i] \leftrightarrow [j]}$.
2. Scaling the j^{th} column by α : $F \circ E^{\alpha \cdot [j]}$.

3. Adjust the j^{th} column by adding to it $\alpha \times i^{th}$ column: $F \circ E^{[i] \xleftarrow{\pm} \alpha \cdot [j]}$.
4. Elementary column operations do not change the injectivity nor image of F .
5. If a column of F is a linear combination of some of the other columns then elementary column operations can turn it into $\mathbf{0}_V$.
6. Removing/Inserting null columns or columns that are linear combinations of other columns does not change the image of F .
7. Given matrix A , it is possible to eliminate (or not) columns of A to end up with a matrix B with $\text{Im}(B) = \text{Im}(A)$.
8. If B is obtained from insertion of columns into matrix A , then $\text{Im}(B) = \text{Im}(A) \iff$ the insert columns $\in \text{Im}(A)$.
9. Inserting columns to a surjective A does not destroy surjectivity of A . (A is already having extra or just enough columns)
10. A surjective $\iff A$ is obtained by inserting columns (or not) into an invertible matrix \iff deleting some columns of A (or not) gives an invertible matrix.
11. If A is injective, then column deletion does not destroy injectivity. (A is already “lacking” or having just enough columns)
12. The new matrix obtained from inserting columns from $\text{Im}(A)$ into A is injective $\iff A$ is injective.

5.5 Linear Independence, Span, and Bases

5.5.1 Linear Independence

- $X_1 \dots X_m$ are linearly independent
 $\iff F = [X_1 \dots X_m]$ injective
 $\iff \sum a_i X_i = 0 \iff a_i = 0 \forall i$
 $\iff \mathbf{0}_V \notin \{X_i\}$ and none are linear combinations of some of the others.
 $\iff X_1 \neq \mathbf{0}_V$ and X_j is not a linear combination of any of X_i 's for $i < j$.

Proposition 5.7.

1. The singleton list is linearly independent if its entry is not the null element
2. Sublists of a linearly independent list are linearly independent
3. List operations cannot create/destroy linearly independence.
4. If a linear map $L : X \rightarrow W$ injective, then $X_1 \dots X_m$ linearly independent $\iff L(X_1) \dots L(X_m)$ linearly independent.

5.5.2 Span

Proposition 5.8.

1. Spans are subspaces. $\text{span}(X_1 \dots X_m), X_j \in V$ is a subspace of V .
2. $\text{span}(X_1 \dots X_j) \subseteq \text{span}(X_1 \dots X_k)$ if $j \leq k$.
3. Adding elements to a list that spans V produces a list that spans V .
4. The following list operations do not change the span of V : removing the null/linearly dependent element, inserting a linearly dependent element, scaling element(s), adding (multiples) of an element to another element.
5. It is possible to reduce a list that spans to a list that spans AND have linearly independent elements.
6. Consider $A : V \rightarrow W$. If X_i span V then $A(X_i)$ span $\text{Im}(A)$. $i = 1, 2, \dots, m$.
7. If $A : V \rightarrow W$ invertible, then X_i span $V \iff A(X_i)$ span W , $i = 1, 2, \dots, m$.

5.5.3 Bases

Proposition 5.9.

1. A list is a basis of V if the elements are linearly independent and they span V .
2. A singleton linear space has no basis.
3. Re-ordering the elements of a basis gives another basis.
4. $\{X_1 \dots X_m\}$ is a basis of V if $(X_1 \dots X_m)$ is injective AND $\text{Im}(X_1 \dots X_m) = V$, i.e. $(X_1 \dots X_m)$ invertible.
5. If $\{X_i\}$ is a basis of V and $\{Y_i\}$ is a list of elements in W , then there exists a unique linear function $L : V \rightarrow W$ satisfying

$$L[X_i] = Y_i$$

6. If $A : V \rightarrow W$ bijective, then $\{V_i\}$ forms a basis of V and $\{A(X_i)\}$ forms a basis of W .
7. Elementary operations on bases give bases.

5.6 Linear Bijections and Isomorphisms

Definition 5.4. Let linear spaces V, W be given. V is **isomorphic** to W if $\exists F : V \rightarrow W$ bijective. We say $V \sim W$.

Proposition 5.10.

1. A non-zero scalar multiple of an isomorphism is an isomorphism.
2. A composition of isomorphism is an isomorphism.
3. “Isomorphism” behaves like an equivalence relation:
 - (a) Reflexivity: $V \sim V$.
 - (b) Symmetry: if $V \sim W$ then $W \sim V$.
 - (c) Transitivity: if $V \sim W$ and $W \sim Z$ then $V \sim Z$.
4. Consider $F : V \rightarrow W$ an isomorphism.
 - (a) Isomorphisms preserve linear independence. V_i ’s are linearly independent in $V \iff F(V_i)$ ’s are linearly independent in W .
 - (b) isomorphisms preserve spanning. V_i ’s span $V \iff F(V_i)$ ’s span W .
 - (c) Isomorphisms preserve bases. $\{V_i\}$ is a basis of $V \iff F\{V_i\}$ is a basis of W .
5. If $V \sim W$ then $\dim(V) = \dim(W)$ (finite or infinite).
6. If a linear map $A : V \rightarrow W$ is given and $A(X_i)$ ’s are linearly independent, then X_i ’s are linearly independent.
7. If a linear map $A : V \rightarrow W$ is injective and $\{X_i\}$ is a basis of V then $\{A(X_i)\}$ is a basis of $\text{Im}(A)$.

5.7 Finite-Dimensional Linear Spaces

5.7.1 Dimension

Proposition 5.11.

1. $\mathbb{R}^m \sim \mathbb{R}^n \iff m = n$.
2. Isomorphisms $F : \mathbb{R}^n \rightarrow V$ are bijective atrices.
3. Isomorphisms $G : W \rightarrow \mathbb{R}^m$ are inverses of bijective atrices.
4. Consider a non-singleton linear space V
 - (a) V has a basis with n elements.
 - (b) $V \sim \mathbb{R}^n$.
 - (c) $V \sim W$, where W is any linear space with a basis of n elements.

5. If V is a linear space with a basis with n elements, then any basis of V has n elements.
6. Linear space $V \sim W$ where W is n -dimensional if V is n -dimensional.
7. For a non-singleton linear space V , $V \sim \mathbb{R}^n \iff \dim(V) = n$.
8. $V \sim W \iff \dim(V) = \dim(W)$.
9. If W is a subspace of V , then $\dim(W) \leq \dim(V)$. Equality holds when $W = V$.

5.7.2 Rank-Nullity Theorem

Let finite-dimensional linear space V and linear map $F : V \rightarrow W$ be given. Then $\text{Im}(F)$ is finite-dimensional and

$$\dim(\text{Im}(F)) + \dim(\ker(F)) = \dim(V)$$

A stronger statement: If a linear map $F : V \rightarrow W$ has finite rank and finite nullity $\iff V$ is finite-dimensional, then

$$\text{Im}(F) + \ker(F) = \dim(V).$$

5.7.3 Coordinatization & matricial representation of linear functions

Consider a finite-dimensional linear space V with basis $\{V_i\}$, $i = 1, 2, \dots, m$. An element \tilde{V} in V can be expressed in exactly one way:

$$\tilde{V} = \sum_{i=1}^m a_i V_i,$$

where $\{a_i\}$ is unique. We call $\{a_i\}$ the coordinate tuple of \tilde{V} and a_i 's the coordinates.

Proposition 5.12.

1. Inverse of a bijective matrix outputs the coordinates. Suppose $A = [V_i]$. Then

$$\tilde{V} = \sum_{i=1}^m a_i V_i \iff A^{-1}(Z) = (a_1 \quad \dots \quad a_m)^\top$$

5.8 Infinite-Dimensional Spaces

Consider a non-singleton linear space V . The following statements are equivalent:

1. V is infinite-dimensional.

2. Every linearly independent list in V can be enlarged to a strictly longer linearly independent set in V .
3. Every linearly independent list in V can be enlarged to an arbitrarily long (finite) linearly independent set in V .
4. There are arbitrarily long (finite) linearly independent lists in V .
5. There are linearly independent lists in V of any (finite) length.
6. No list of finitely many elements of V spans V .

6 Sums of subspaces & Products of vector spaces

6.1 Direct Sums

Definition 6.1. Let U_j , $j = 1, 2, \dots, m$ are subspaces of V . $\sum_1^m U_j$ is a *direct sum* if each $u \in \sum U_j$ can be written in only one way as $u = \sum_1^m u_j$. The direct sum $\sum_i^m U_j$ is denoted as $U_1 \oplus \dots \oplus U_m$.

Proposition 6.1.

1. Condition for direct sum: If all U_j are subspaces of V , then $\sum_1^m U_j$ is a direct sum \iff the only way to write 0 as $\sum_1^m u_j$, where $u_j \in U_j$ is to take $u_j = 0$ for all j .
2. If U, W are subspaces of V and $U \cap W = \{0\}$ then $U + W$ is a direct sum.

6.2 Products and Quotients of Vector Spaces

6.3 Products of Vector Spaces

Definition 6.2. Product of vectors spaces

$$V_1 \times \dots \times V_m = \{(v_1, \dots, v_m) : v_j \in V_j, j = 1, 2, \dots, m\}.$$

Definition 6.3. Addition on $V_1 \times \dots \times V_m$:

$$(u_1, \dots, u_m) + (v_1, \dots, v_m) = (u_1 + v_1, \dots, v_m + u_m).$$

Definition 6.4. Scalar multiplication on $V_1 \times \dots \times V_m$:

$$\lambda(v_1, \dots, v_m) = (\lambda v_1, \dots, \lambda v_m).$$

Proposition 6.2.

1. Product of vectors spaces is a vector space.
 V_j are vectors spaces over $\mathbb{F} \implies V_1 \times \dots \times V_m$ is a vector space over \mathbb{F} .
2. Dimension of a product is the sum of dimensions:

$$\dim(V_1 \times \dots \times V_m) = \sum_1^m \dim(V_j)$$

6.4 Products & Direct Sums

Proposition 6.3.

1. Let U_1, \dots, U_m be subspaces of V . Define a linear map $\Gamma : U_1 \times \dots \times U_m \rightarrow U_1 + \dots + U_m$ by:

$$\Gamma(u_1, \dots, u_m) = \sum_1^m u_j.$$

$U_1 + \dots + U_m$ is a direct sum $\iff \Gamma$ is injective.

2. Let U_j be finite-dimensional and are subspaces of V .

$$U_1 \oplus \cdots \oplus U_m \iff \dim(U_1 + \cdots + U_m) = \sum_1^m \dim(U_j)$$

6.5 Quotients of Vector Spaces

6.6 Nullspaces

6.7 Ranges of Operator Powers

7 Idempotents & Resolutions of Identity

8 Block-representations of operators

8.1 Direct sums of operators

9 Invariant subspaces

9.1 Reducing subspaces

10 Polynomials applied to operators

10.1 Minimal polynomials of block- Δ^r operators

10.2 Minimal polynomials at a vector

11 Eigentheory

11.1 Spectral Mapping Theorem

12 Triangularization

12.1 Compression to invariant subspaces

12.2 Simultaneously Δ -ity of commuting families

13 Diagonalization

13.1 Spectral resolutions

13.2 Compressions to reducing subspaces

13.3 Simultaneous diagonalizability for commuting families

14 Primary decomposition over \mathbb{C} and generalized eigenspaces

15 Cyclic decomposition and Jordan form

15.1 Square roots of operators

15.2 Similarity of a matrix and its transpose

15.3 Similarity of a matrix and its conjugate

15.4 Jordan forms of AB and BA

15.5 Power-convergent operators

15.6 Power-bounded operators

15.7 Row-stochastic matrices

16 Determinant & Trace

16.1 Classical adjoints

16.2 Cayley-Hamilton theorem

17 Inner products and norms

- 17.1 Riesz representation theorem
- 17.2 Adjoints
- 17.3 Grammians
- 17.4 Orthogonal complements and orthogonal decompositions
- 17.5 Ortho-projections
- 17.6 Closest point solutions
- 17.7 Gram-Schmidt and orthonormal bases

18 Isometries and unitary operators

19 Ortho-triangularization

20 Spectral resolutions

21 Ortho-diagonalization; self-adjoint and normal operators; spectral theorems

22 Positive (semi-)definite operators

22.1 Classification of inner products

22.2 Positive square roots

23 Polar decomposition

24 Single value decomposition

24.1 Spectral/operator norm

24.2 Singular values and approximation

24.3 Singular values and eigenvalues

25 Problems and Solutions

25.1 Problem set 1

Problem. Prelim. 1.1. Suppose that \mathbf{V} is a finite-dimensional vector space. Consider the following subspace \mathbf{W} of the vector space $\mathbf{V} \times \mathbf{V}$:

$$\mathbf{W} = \left\{ \begin{pmatrix} v \\ -v \end{pmatrix} \middle| v \in \mathbf{V} \right\}.$$

Argue that

$$\dim(\mathbf{V}) = \dim(\mathbf{W}).$$

Solution.

We can equivalently show that $\dim(V) = \dim(W)$ by showing that \mathbf{V} is isomorphic to \mathbf{W} ($\mathbf{V} \sim \mathbf{W}$), i.e., there exists a bijective linear function $\mathcal{L} : \mathbf{V} \rightarrow \mathbf{W} \prec \mathbf{V} \times \mathbf{V}$.

We can define $\mathcal{L} : \mathbf{V} \rightarrow \mathbf{W}$ as

$$\mathcal{L}[v] = \begin{pmatrix} v \\ -v \end{pmatrix},$$

where $v \in \mathbf{V}$. Let $v_1, v_2 \in \mathbf{V}$ and $a, b \in \mathbb{C}$ be given, \mathcal{L} satisfies the linearity condition because

$$\mathcal{L}[av_1 + bv_2] = \begin{pmatrix} av_1 + bv_2 \\ -(av_1 + bv_2) \end{pmatrix} = a \begin{pmatrix} v_1 \\ -v_1 \end{pmatrix} + b \begin{pmatrix} v_2 \\ -v_2 \end{pmatrix} = a\mathcal{L}[v_1] + b\mathcal{L}[v_2],$$

where the second equality follows from the fact that \mathbf{W} is a subspace. Therefore, \mathcal{L} is a linear function.

Now, \mathcal{L} is surjective, by definition:

$$\text{Im}(\mathcal{L}) = \left\{ \begin{pmatrix} v \\ -v \end{pmatrix} \middle| v \in \mathbf{V} \right\} = \mathbf{W}.$$

\mathcal{L} is also injective, since $\ker(\mathcal{L}) = \mathbf{0}_V$:

$$\mathcal{L}[v] = \mathbf{0}_W \iff \begin{pmatrix} v \\ -v \end{pmatrix} = \mathbf{0}_W = \begin{pmatrix} \mathbf{0}_V \\ \mathbf{0}_V \end{pmatrix} \iff v = \mathbf{0}_V.$$

Therefore, \mathcal{L} is an isomorphism. It follows that $\mathbf{V} \sim \mathbf{W} \iff \dim(V) = \dim(W)$.

Problem. Prelim. 1.2. Argue that $\mathbf{V} + \{\mathbf{0}_{\mathbf{W}}\} = \mathbf{V}$, for any subspace \mathbf{V} of a vector space \mathbf{W} .

Solution.

We shall first argue that $\mathbf{0}_{\mathbf{W}} = \mathbf{0}_{\mathbf{V}}$. Let $v_+ \in \mathbf{V}$ be given, and let the unique additive antipodal element of v_+ be $v_- \in \mathbf{V}$. Because \mathbf{V} is a subspace of \mathbf{W} , $v_+ \in \mathbf{W}$ and $v_- \in \mathbf{W}$,

$$\mathbf{0}_{\mathbf{V}} = v_+ \overset{\mathbf{V}}{+} v_- = v_+ \overset{\mathbf{W}}{+} v_- = \mathbf{0}_{\mathbf{W}}.$$

Therefore, $\{\mathbf{0}_{\mathbf{W}}\} = \{\mathbf{0}_{\mathbf{V}}\}$. It then follows that

$$\mathbf{V} + \{\mathbf{0}_{\mathbf{W}}\} = \mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\}. \quad (1)$$

Now, it suffices to show that (i) \mathbf{V} is a subspace of $\mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\}$ and (ii) $\mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\}$ is a subspace of \mathbf{V} .

Consider $v \in \mathbf{V}$. It follows trivially that $v \in \mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\}$, so $\mathbf{V} \subseteq \mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\}$. But because both \mathbf{V} and $\mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\}$, \mathbf{V} is a subspace of $\mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\}$. Hence (i) is true.

Consider $v_0 \in \mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\}$, then for some $a_v, a_0 \in \mathbb{C}$, $v_{\mathbf{V}} \in \mathbf{V}$, and $v_0 \in \{\mathbf{0}_{\mathbf{V}}\}$,

$$v_0 = a_v v_{\mathbf{V}} + a_0 v_0.$$

But v_0 is identically $\mathbf{0}_{\mathbf{V}}$, so

$$v_0 = a_v v_{\mathbf{V}} + a_0 \mathbf{0}_{\mathbf{V}} = a_v v_{\mathbf{V}} \in \mathbf{V},$$

which implies $\mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\} \subseteq \mathbf{V}$. Therefore $\mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\}$ is a subspace of \mathbf{V} , i.e., (ii) is true.

Hence, by Eq. (1), facts (i), and (ii),

$$\mathbf{V} + \{\mathbf{0}_{\mathbf{V}}\} = \mathbf{V}.$$

Problem. 1. Suppose that $\mathbf{V}_1, \mathbf{V}_2, \dots, \mathbf{V}_{315}$ are subspaces of a vector space \mathbf{W} . Argue that the following claims are equivalent.

1. The subspace sum $\mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_{315}$ is direct.
2. If $x_i \in \mathbf{V}_i$ and

$$x_1 + x_2 + \dots + x_{315} = \mathbf{0}_{\mathbf{W}},$$

then $x_i = \mathbf{0}_{\mathbf{W}}$ for every i .

3. For any i , no non-null element of \mathbf{V}_i can be express as a sum of the elements of the other \mathbf{V}_j 's.
4. For any i , no non-null element of \mathbf{V}_i can be expressed as a sum of the elements of the preceding \mathbf{V}_j 's.

Solution. It suffices to show $1 \iff 2 \iff 3 \iff 4$.

- ($1 \implies 2$) If $\mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_{315}$ is a direct sum, then by definition the only way to write

$$\mathbf{0}_{\mathbf{W}} = x_1 + x_2 + \dots + x_{315}, \quad (2)$$

where $x_i \in \mathbf{V}_i$, is by requiring $x_i = \mathbf{0}_{\mathbf{W}}$ for all i . In other words, taking all $x_i = \mathbf{0}_{\mathbf{W}}$ for all i is the *one and only way* to satisfy Eq. (2) because $\mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_{315}$ is a direct sum.

- ($2 \implies 1$) Let $a, b \in \mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_{315}$ be given such that

$$\hat{h} = x_1 + x_2 + \dots + x_{315} = y_1 + y_2 + \dots + y_{315}, \quad (3)$$

where $x_i, y_i \in \mathbf{V}_i$. It suffices to show $x_i = y_i$, i.e, any representation of \hat{h} as Eq. (3) is unique. Consider

$$\mathbf{0}_{\mathbf{W}} = \hat{h} - \hat{h} = (x_1 - y_1) + (x_2 - y_2) + \dots + (x_{315} - y_{315})$$

By assuming the second statement and the fact that $x_i, y_i \in \mathbf{V}_i$, $x_i = y_i$ for all i . Therefore, any representation of \hat{h} is unique, as desired by the first statement.

- ($3 \implies 2$) Suppose that the second statement is incorrect. Without loss of generality, suppose that for $i = 1$, $\mathbf{0}_{\mathbf{W}} \neq x_1 \in \mathbf{V}_1$

$$x_1 + x_2 + \dots + x_{315} = \mathbf{0}_{\mathbf{W}}.$$

It immediately follows that

$$x_1 = -(x_2 + x_3 + \dots + x_{315}) \neq \mathbf{0}_{\mathbf{W}},$$

which implies the third statement is also incorrect. By contraposition, $3 \implies 2$.

- (2 \implies 3) Suppose that $x_1 = -x_2 \neq \mathbf{0}_W$, $x_1 \in V_1$, $x_2 \in V_2$. Then with $x_j = \mathbf{0}_W$ for all $j = 3, 4, \dots, 315$,

$$\begin{aligned} x_1 + x_2 + x_3 + \dots + x_{315} &= x_1 - x_1 + x_3 + \dots + x_{315} \\ &= x_3 + \dots + x_{315} \\ &= \mathbf{0}_W. \end{aligned}$$

But $x_1 = -x_2 \neq \mathbf{0}_W$. Therefore the second statement no longer holds. By contraposition, it implies the third statement.

- (3 \iff 4) This is evidently true since “sum of the elements of the preceding V_j ” is a subset of “sums of the elements of the other V_j ’s.” Conversely, if no non-null element of V_i can be expressed as a sum of the elements of the preceding V_j ’s, then the third statement is also true. (Imagine going through an ordered list, checking for a counter example, and not succeeding).

Problem. 2. Sub-sums of direct sums are direct:

Suppose that $\mathbf{V}_1, \mathbf{V}_2, \dots, \mathbf{V}_{315}$ are subspaces of a vector space \mathbf{W} , and the subspace sum $\mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_{315}$ is direct.

1. Suppose that for each i , \mathbf{Z}_i is a subspace of \mathbf{V}_i . Argue that the subspace sum $\mathbf{Z}_1 + \mathbf{Z}_2 + \dots + \mathbf{Z}_{315}$ is direct.
2. Argue that the sum $\mathbf{V}_2 + \mathbf{V}_5 + \mathbf{V}_7 + \mathbf{V}_{12}$ is also direct.

Solution.

1. By the previous problem, because $\mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_{315}$ is direct, for any i , no non-null element of \mathbf{V}_i can be expressed as a sum of the elements of the other \mathbf{V}_j 's. Since \mathbf{Z}_i is a subspace of \mathbf{V}_i , no $\mathbf{0}_{\mathbf{W}} \neq z_i \in \mathbf{Z}_i \prec \mathbf{V}_i$ can be expressed as a sum of the elements of the other \mathbf{Z}_j 's $\prec \mathbf{V}_j$'s. Hence, the subspace sum $\mathbf{Z}_1 + \mathbf{Z}_2 + \dots + \mathbf{Z}_{315}$ is also direct.
2. This is true by (1). Since each \mathbf{V}_j is a subspace of itself, the subspace sum of distinct \mathbf{V}_j 's is direct. Therefore, the sum $\mathbf{V}_2 + \mathbf{V}_5 + \mathbf{V}_7 + \mathbf{V}_{12}$ is direct.

Problem. 3. Associativity of directness of subspace sums: Suppose that

$$\mathbf{Y}, \mathbf{V}_1, \mathbf{V}_2, \dots, \mathbf{V}_5, \mathbf{U}_1, \mathbf{U}_2, \dots, \mathbf{U}_{12}, \mathbf{Z}_1, \mathbf{Z}_2, \dots, \mathbf{Z}_4, \mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{53}$$

are subspaces of a vector space \mathbf{W} . Argue that the following claims are equivalent.

1. The subspace sum

$$\begin{aligned} & \mathbf{Y} + \mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_5 + \mathbf{U}_1 + \mathbf{U}_2 + \dots \\ & + \mathbf{U}_{12} + \mathbf{Z}_1 + \mathbf{Z}_2 + \dots + \mathbf{Z}_4 + \mathbf{X}_1 + \mathbf{X}_2 + \dots + \mathbf{X}_{53} \end{aligned}$$

is direct.

2. The subspace sums

$$\begin{aligned} [\mathbf{V} &:=] \mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_5 \\ [\mathbf{U} &:=] \mathbf{U}_1 + \mathbf{U}_2 + \dots + \mathbf{U}_{12} \\ [\mathbf{Z} &:=] \mathbf{Z}_1 + \mathbf{Z}_2 + \dots + \mathbf{Z}_4 \\ [\mathbf{X} &:=] \mathbf{X}_1 + \mathbf{X}_2 + \dots + \mathbf{X}_{53} \\ \mathbf{Y} &+ \mathbf{V} + \mathbf{U} + \mathbf{X} + \mathbf{Z} \end{aligned}$$

are all direct.

Solution.

1. ($1 \implies 2$) If the subspace sum

$$\begin{aligned} & \mathbf{Y} + \mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_5 + \mathbf{U}_1 + \mathbf{U}_2 + \dots \\ & + \mathbf{U}_{12} + \mathbf{Z}_1 + \mathbf{Z}_2 + \dots + \mathbf{Z}_4 + \mathbf{X}_1 + \mathbf{X}_2 + \dots + \mathbf{X}_{53} \end{aligned}$$

is direct, then by part 2 the previous problem, the subspace sums

$$\begin{aligned} [\mathbf{V} &:=] \mathbf{V}_1 + \mathbf{V}_2 + \dots + \mathbf{V}_5 \\ [\mathbf{U} &:=] \mathbf{U}_1 + \mathbf{U}_2 + \dots + \mathbf{U}_{12} \\ [\mathbf{Z} &:=] \mathbf{Z}_1 + \mathbf{Z}_2 + \dots + \mathbf{Z}_4 \\ [\mathbf{X} &:=] \mathbf{X}_1 + \mathbf{X}_2 + \dots + \mathbf{X}_{53} \\ \mathbf{Y} &+ \mathbf{V} + \mathbf{U} + \mathbf{X} + \mathbf{Z} \end{aligned}$$

are all direct.

2. ($2 \implies 1$) This follows trivially because the subspace sum

$$\mathbf{Y} \oplus \mathbf{V} \oplus \mathbf{U} \oplus \mathbf{X} \oplus \mathbf{Z}$$

can be written as

$$\begin{aligned} & \mathbf{Y} \oplus (\mathbf{V}_1 \oplus \mathbf{V}_2 \oplus \dots \oplus \mathbf{V}_5) \oplus (\mathbf{U}_1 \oplus \mathbf{U}_2 \oplus \dots \oplus \mathbf{U}_{12}) \\ & \oplus (\mathbf{Z}_1 \oplus \mathbf{Z}_2 \oplus \dots \oplus \mathbf{Z}_4) \oplus (\mathbf{X}_1 \oplus \mathbf{X}_2 \oplus \dots \oplus \mathbf{X}_{53}). \end{aligned} \quad (4)$$

Because vector space addition is associative, (4) is equivalent to saying the subspace sum in the first statement is direct.

Problem. 4. Direct sums preserve linear independence: Suppose that $\mathbf{U}, \mathbf{V}, \mathbf{W}$ are subspaces of a vector space \mathbf{Z} , and the sum $\mathbf{U}, \mathbf{V}, \mathbf{W}$ is direct.

1. Suppose that U_1, U_2, \dots, U_{13} is a linearly independent list in \mathbf{U} , V_1, V_2, \dots, V_6 is a linearly independent list in \mathbf{V} , and W_1, W_2, \dots, W_{134} is a linearly independent list in \mathbf{W} . Argue that the concatenated list

$$U_1, U_2, \dots, U_{13}, V_1, V_2, \dots, V_6, W_1, W_2, \dots, W_{134}$$

is linearly independent.

2. Suppose that U_1, U_2, \dots, U_{13} is a basis of \mathbf{U} , V_1, V_2, \dots, V_6 is a basis of \mathbf{V} , and W_1, W_2, \dots, W_{134} is a basis of \mathbf{W} . Argue that the concatenated list

$$U_1, U_2, \dots, U_{13}, V_1, V_2, \dots, V_6, W_1, W_2, \dots, W_{134}$$

is a basis of $\mathbf{U} \oplus \mathbf{V} \oplus \mathbf{W}$.

Solution.

1. It suffices to consider a linearly independent $U_1 \in \mathbf{U}$ and \mathbf{V} . Since $\mathbf{U} + \mathbf{V} + \mathbf{W}$ is a direct sum, by the previous problem, $\mathbf{U} + \mathbf{V}$ is a direct sum. Therefore, $U_1 \neq \mathbf{0}_Z$ cannot be expressed as a sum of the elements of \mathbf{V} . Therefore, U_1 is linearly independent of all V_i 's $\in \mathbf{V}$. The same is true with regards to U_1 and all W_i 's $\in \mathbf{W}$.

It follows, without loss of generality, that if U_1, U_2, \dots, U_{13} is a linearly independent list in \mathbf{U} , V_1, V_2, \dots, V_6 is a linearly independent list in \mathbf{V} , and W_1, W_2, \dots, W_{134} is a linearly independent list in \mathbf{W} , then the list

$$U_1, U_2, \dots, U_{13}, V_1, V_2, \dots, V_6, W_1, W_2, \dots, W_{134}$$

is linearly independent.

2. Since U_1, U_2, \dots, U_{13} is a basis of \mathbf{U} , V_1, V_2, \dots, V_6 is a basis of \mathbf{V} , and W_1, W_2, \dots, W_{134} is a basis of \mathbf{W} , the concatenated list

$$U_1, U_2, \dots, U_{13}, V_1, V_2, \dots, V_6, W_1, W_2, \dots, W_{134}$$

is linearly independent. It, then, suffices to show that the concatenated list spans $\mathbf{U} \oplus \mathbf{V} \oplus \mathbf{W}$.

Consider $H \in \mathbf{U} \oplus \mathbf{V} \oplus \mathbf{W}$, H can be written as

$$H = u_h U_h + v_h V_h + w_h W_h,$$

where $u_h, v_h, w_h \in \mathbb{C}$, $U_h \in \mathbf{U}$, $V_h \in \mathbf{V}$, $W_h \in \mathbf{W}$. Now, U_h can be written as a linear combination of U_1, U_2, \dots, U_{13} because U_1, U_2, \dots, U_{13} form a basis of \mathbf{U} . Similarly, V_h, W_h can also be written as linear combination

of basis elements of their respective subspaces. It follows that H can be expressed as a linear combination of the given concatenated list, i.e, the concatenated list spans $\mathbf{U} \oplus \mathbf{V} \oplus \mathbf{W}$.

Therefore, the concatenated list

$$U_1, U_2, \dots, U_{13}, V_1, V_2, \dots, V_6, W_1, W_2, \dots, W_{134}$$

is a basis of $\mathbf{U} \oplus \mathbf{V} \oplus \mathbf{W}$.

Problem. 5. Suppose that x_1, x_2, \dots, x_{14} are non-null elements of a linear space \mathbf{W} .

1. Argue that the following claims are equivalent.

- (a) x_1, x_2, \dots, x_{14} are linearly independent.
- (b) The subspace sum

$$\text{span}(x_1) + \text{span}(x_2) + \cdots + \text{span}(x_{14})$$

is direct.

2. Argue that the following claims are equivalent.

- (a) x_1, x_2, \dots, x_{14} is a basis of \mathbf{W} .
- (b) $\mathbf{W} = \text{span}(x_1) \oplus \text{span}(x_2) \oplus \cdots \oplus \text{span}(x_{14})$

Solution.

1. (a) $(a \implies b)$ If x_1, x_2, \dots, x_{14} are linearly independent, then any x_i is neither null nor can be expressed as a linear combination of the other x_j 's. Therefore, no non-null $x_i \in \text{span}(x_i)$ can be written as a sum of other x_j 's $\in \text{span}(x_j)$. This is equivalent to

$$\text{span}(x_1) + \text{span}(x_2) + \cdots + \text{span}(x_{14})$$

being direct.

- (b) $(b \implies a)$ The subspace sum

$$\text{span}(x_1) + \text{span}(x_2) + \cdots + \text{span}(x_{14})$$

being direct implies that no non-null $x_i \in \text{span}(x_i)$ can be written as a sum of other x_j 's $\in \text{span}(x_j)$. Therefore, the list x_1, x_2, \dots, x_{14} is linearly independent.

2. Because x_i is a basis of $\text{span}(x_i)$, by the previous problem, the concatenated list x_1, x_2, \dots, x_{14} is a basis of $\text{span}(x_1) \oplus \text{span}(x_2) \oplus \cdots \oplus \text{span}(x_{14})$.
(†)

- (a) $(a \implies b)$ Let $w \in \text{span}(x_1) \oplus \text{span}(x_2) \oplus \cdots \oplus \text{span}(x_{14})$ be given, then by (†), w can be expressed as a combination of the x_i 's, where $x_i \in \text{span}(x_i)$. But this is equivalent to $w \in \mathbf{W}$ because x_1, x_2, \dots, x_{14} is a basis of \mathbf{W} . Therefore, $\mathbf{W} = \text{span}(x_1) \oplus \text{span}(x_2) \oplus \cdots \oplus \text{span}(x_{14})$.
- (b) $(b \implies a)$ It follows from fact (†) that if $\mathbf{W} = \text{span}(x_1) \oplus \text{span}(x_2) \oplus \cdots \oplus \text{span}(x_{14})$, then x_1, x_2, \dots, x_{14} is a basis of \mathbf{W} .

25.2 Problem set 2

25.3 Problem set 3