

QUANTUM MECHANICS

A Quick Guide

Huan Q. Bui

Colby College

PHYSICS & MATHEMATICS

Statistics

Class of 2021

June 5, 2019

Preface

Greetings,

Quantum Mechanics, A Quick Guide to... is my reading notes from Shankar's *Principles of Quantum Mechanics, Second Edition*. Additional material will come from my class notes and my comments/interpretations/solutions. Enjoy!

Contents

Preface	1
1 Mathematical Introduction	4
1.1 Linear Vector Spaces	4
1.2 Inner Product Spaces	5
2 Review of Classical Mechanics	7
3 All is Not Well with Classical Mechanics	8
4 The Postulates –a General Discussion	9
5 Simple Problems in One Dimension	10
6 The Classical Limit	11
7 The Harmonic Oscillator	12
8 The Path Integral Formulation of Quantum Theory	13
9 The Heisenberg Uncertainty Relation	14
10 Systems with N Degrees of Freedom	15
11 Symmetries and Their Consequences	16
12 Rotational Invariance and Angular Momentum	17
13 The Hydrogen Atom	18
14 Spin	19
15 Additional of Angular Momentum	20
16 Variational and WKB Methods	21
17 Time-Independent Perturbation Theory	22
18 Time-Dependent Perturbation Theory	23
19 Scattering Theory	24
20 The Dirac Equation	25
21 Path Integrals–II	26

22 Selected Problems and Solutions	27
22.1 Mathematical Introduction	27
22.2 Review of Classical Mechanics	28
22.3 All is Not Well with Classical Mechanics	29
22.4 The Postulates –a General Discussion	30
22.5 Simple Problems in One Dimension	31
22.6 The Classical Limit	32
22.7 The Harmonic Oscillator	33
22.8 The Path Integral Formulation of Quantum Theory	34
22.9 The Heisenberg Uncertainty Relation	35
22.10 Systems with N Degrees of Freedom	36
22.11 Symmetries and Their Consequences	37
22.12 Rotational Invariance and Angular Momentum	38
22.13 The Hydrogen Atom	39
22.14 Spin	40
22.15 Addition of Angular Momentum	41
22.16 Variational and WKB Methods	42
22.17 Time-Independent Perturbation Theory	43
22.18 Time-Dependent Perturbation Theory	44
22.19 Scattering Theory	45
22.20 The Dirac Equation	46
22.21 Path Integrals–II	47

1 Mathematical Introduction

1.1 Linear Vector Spaces

We should familiar with defining characteristics of linear vector spaces at this point. Here are some important definitions/theorems again:

Definition 1.1. A linear vector space \mathbf{V} is a collection of objects called *vectors* for which there exists

1. A definite rule for summing, and
2. A definite rule for scaling, with the following features:
 - Closed under addition: for $x, y \in \mathbf{V}$, $x + y \in \mathbf{V}$.
 - Closed under scalar multiplication: $x \in \mathbf{V}$, then $ax \in \mathbf{V}$ for some scalar a .
 - Scalar multiplication is distributive.
 - Scalar multiplication is associative.
 - Addition is commutative.
 - Addition is associative.
 - There exists a (unique) null element in \mathbf{V} .
 - There exists a (unique) additive inverse.

Vector spaces are defined over some field. The field can be real numbers, complex numbers, or it can also be finite. As for good practice, we will begin to label vectors with Dirac bra-ket notation. So, for instance, $|v\rangle \in \mathbf{V}$ denotes vector $v \in \mathbf{V}$. Basic manipulations of these vectors are intuitive:

1. $|0\rangle$ is unique, and is the null element.
2. $0|V\rangle = |0\rangle$.
3. $|-V\rangle = -|V\rangle$.
4. $|-V\rangle$ is a unique additive inverse of $|V\rangle$.

The reasons for choosing to use the Dirac notation will become clear later on. Another important basic concept is *linear (in)dependence*. Of course, there are a number of equivalent statement for linear independence. We shall just give one here:

Definition 1.2. A set of vectors is said to be linearly independent if the only linear relation

$$\sum_{i=1}^n a_i |i\rangle = |0\rangle \tag{1}$$

is the trivial one where the components $a_i = 0$ for any i .

The next two basic concepts are *dimension* and *basis*.

Definition 1.3. A vector space \mathbf{V} has dimension n if it can accommodate a maximum of n linearly independent vectors. We denote this n -dimensional vector space as \mathbf{V}^n .

We can show that

Theorem 1.1. Any vector $|v\rangle \in \mathbf{V}^n$ can be written (uniquely) as a linear combination of any n linearly independent vectors.

Definition 1.4. A set of n linearly independent vectors in a n -dimensional space is called a *basis*. So if $|1\rangle, \dots, |n\rangle$ form a basis for \mathbf{V}^n , then any $|v\rangle \in \mathbf{V}$ can be written uniquely as

$$|v\rangle = \sum_{i=1}^n a_i |i\rangle. \quad (2)$$

It is nice to remember the following:

$$\boxed{\text{Linear Independence} = \text{Basis} + \text{Span}} \quad (3)$$

When a collection of vectors span a vector space \mathbf{V} , it just means that any $|v\rangle \in \mathbf{V}$ can be written as a linear combination of (some of) these vectors.

The algebra of linear combinations is quite intuitive. If $|v\rangle = \sum_i a_i |i\rangle$ and $|w\rangle = \sum_i b_i |i\rangle$ then

1. $|v + w\rangle = \sum_i (a_i + b_i) |i\rangle$.
2. $c|v\rangle = c \sum_i a_i |i\rangle = \sum_i ca_i |i\rangle$.

A linear algebra text will of course provide a much better coverage of these topics.

1.2 Inner Product Spaces

A generalization of the familiar dot product is the *inner product* or the *scalar product*. An inner product between two vectors $|v\rangle$ and $|w\rangle$ is denoted $\langle v|w\rangle$. An inner product has to satisfy the following properties:

1. Conjugate symmetry (or skew-symmetry): $\langle v|w\rangle = \langle w|v\rangle^*$.
2. Positive semi-definiteness: $\langle v|v\rangle \geq 0$.
3. Linearity in ket: $\langle v|aw + bz\rangle = a\langle v|w\rangle + b\langle v|z\rangle$.
4. Conjugate-linearity in bra: $\langle av + bz|w\rangle = \bar{a}\langle v|w\rangle + \bar{b}\langle z|w\rangle$.

Definition 1.5. An inner product space is a vector space with an inner product.

Definition 1.6. $\langle v|w\rangle = 0 \iff |v\rangle \perp |w\rangle$.

Definition 1.7. The *norm* (or length) of $|v\rangle$ is defined as

$$\|v\| = \sqrt{\langle v|v\rangle}. \quad (4)$$

Unit vectors have unit norm. Unit vectors are said to be *normalized*.

Definition 1.8. A set of basis vectors all of unit norm, which are pairwise orthogonal will be called an *orthonormal basis* or ONB.

Let $|v\rangle = \sum_i a_i |i\rangle$ and $|w\rangle = \sum_i b_i |j\rangle$, then

$$\langle v|w\rangle = \sum_i a_i^* b_i \langle i|j\rangle. \quad (5)$$

2 Review of Classical Mechanics

3 All is Not Well with Classical Mechanics

4 The Postulates –a General Discussion

5 Simple Problems in One Dimension

6 The Classical Limit

7 The Harmonic Oscillator

8 The Path Integral Formulation of Quantum Theory

9 The Heisenberg Uncertainty Relation

10 Systems with N Degrees of Freedom

11 Symmetries and Their Consequences

12 Rotational Invariance and Angular Momentum

13 The Hydrogen Atom

14 Spin

15 Additional of Angular Momentum

16 Variational and WKB Methods

17 Time-Independent Perturbation Theory

18 Time-Dependent Perturbation Theory

19 Scattering Theory

20 The Dirac Equation

21 Path Integrals–II

22 Selected Problems and Solutions

22.1 Mathematical Introduction

22.2 Review of Classical Mechanics

22.3 All is Not Well with Classical Mechanics

22.4 The Postulates –a General Discussion

22.5 Simple Problems in One Dimension

22.6 The Classical Limit

22.7 The Harmonic Oscillator

22.8 The Path Integral Formulation of Quantum Theory

22.9 The Heisenberg Uncertainty Relation

22.10 Systems with N Degrees of Freedom

22.11 Symmetries and Their Consequences

22.12 Rotational Invariance and Angular Momentum

22.13 The Hydrogen Atom

22.14 Spin

22.15 Additional of Angular Momentum

22.16 Variational and WKB Methods

22.17 Time-Independent Perturbation Theory

22.18 Time-Dependent Perturbation Theory

22.19 Scattering Theory

22.20 The Dirac Equation

22.21 Path Integrals–II