

Name: **Huan Q. Bui**  
Course: **8.309 - Classical Mechanics III**  
Problem set: **#1**

## 1. Two Particles in a Gravitational Field

(a) In the center of mass (COM) frame, the Lagrangian is given by

$$\begin{aligned}
\mathcal{L} &= T - U \\
&= T - U_{\text{attraction}} - U_g \\
&= \left[ \frac{1}{2} m_1 \dot{\vec{r}}_1^2 + \frac{1}{2} m_2 \dot{\vec{r}}_2^2 \right] - \left[ -\frac{Gm_1 m_2}{|\vec{r}|} - g(m_1 x_1 + m_2 x_2) \right] \\
&= \frac{1}{2} (m_1 + m_2) \dot{\vec{R}}_{\text{COM}}^2 + \frac{1}{2} \frac{m_1 m_2}{m_1 + m_2} \dot{\vec{r}}^2 + \frac{Gm_1 m_2}{r} + g(m_1 + m_2) X_{\text{COM}},
\end{aligned}$$

where  $X_{\text{COM}}$  is the  $x$ -component of  $\vec{R}_{\text{COM}}$ . To obtain this Lagrangian, we have solved for  $\vec{r}_1$  and  $\vec{r}_2$  in terms of  $\vec{R}_{\text{COM}}$  and  $\vec{r}$  from the following definitions:

$$\begin{cases} \vec{R}_{\text{COM}} = (m_1 \vec{r}_1 + m_2 \vec{r}_2) / (m_1 + m_2) \\ \vec{r} = \vec{r}_2 - \vec{r}_1 \end{cases} \implies \begin{cases} \vec{r}_1 = \vec{R}_{\text{COM}} - m_2 \vec{r} / (m_1 + m_2) \\ \vec{r}_2 = \vec{R}_{\text{COM}} + m_1 \vec{r} / (m_1 + m_2) \\ X_{\text{COM}} = (m_1 x_1 + m_2 x_2) / (m_1 + m_2) \end{cases}$$

Calling the total mass  $m_1 + m_2 = M$  and reduced mass  $m_1 m_2 / (m_1 + m_2) = \mu$ , we have

$$\mathcal{L} = \underbrace{\frac{1}{2} M \dot{\vec{R}}_{\text{COM}}^2 + g M X_{\text{COM}}}_{\text{COM}} + \underbrace{\frac{1}{2} \mu \dot{\vec{r}}^2 + \frac{Gm_1 m_2}{r}}_{\text{relative}}$$

The Lagrangian splits into two parts which describe the center-of-mass and relative dynamics, respectively. This makes sense physically because both bodies are essentially in “free fall” with each other. The center of mass of the system is decoupled from the relative motion, i.e. we can go to a frame in which the center of mass is stationary, and the only dynamics left is the relative motion of the masses.

(b) Going to the center of mass frame, we have the following Lagrangian

$$\mathcal{L}_r = \frac{1}{2} \mu \dot{\vec{r}}^2 + \frac{Gm_1 m_2}{r}.$$

To avoid taking derivatives of the basis vectors in spherical coordinates, we may write  $\dot{\vec{r}} = \dot{x}^2 + \dot{y}^2 + \dot{z}^2$  where  $(x, y, z) = r(\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$  and let Mathematica compute the more familiar derivatives. The result is

$$\mathcal{L}_r = \frac{1}{2} \mu \left[ \dot{r}^2 + r^2 \left( \dot{\theta}^2 + \dot{\phi}^2 \sin^2 \theta \right) \right] + \frac{Gm_1 m_2}{r}$$

The corresponding Euler-Lagrange equations are

$$\begin{aligned}
\frac{d}{dt} \left( \frac{\partial \mathcal{L}_r}{\partial \dot{r}} \right) &= \frac{\partial \mathcal{L}_r}{\partial r} \implies \mu \ddot{r} = -\frac{Gm_1 m_2}{r^2} + \mu r \left( \dot{\theta}^2 + \dot{\phi}^2 \sin^2 \theta \right) \\
\frac{d}{d\theta} \left( \frac{\partial \mathcal{L}_r}{\partial \dot{\theta}} \right) &= \frac{\partial \mathcal{L}_r}{\partial \theta} \implies 2\dot{r}\dot{\theta} + r\ddot{\theta} = r\dot{\phi}^2 \cos \theta \sin \theta \\
\frac{d}{d\phi} \left( \frac{\partial \mathcal{L}_r}{\partial \dot{\phi}} \right) &= \frac{\partial \mathcal{L}_r}{\partial \phi} \implies \mu r \sin \theta \left[ 2 \left( \dot{r} \sin \theta + r\dot{\theta} \cos \theta \right) \dot{\phi} + r\ddot{\phi} \sin \theta \right] = 0
\end{aligned}$$

- (c) The Hamiltonian corresponding to this Lagrangian is obtained via the Legendre transform. To do this, we first find the canonical momenta  $p_r, p_\theta, p_\phi$  in Mathematica using  $p_i = \partial \mathcal{L}_r / \partial \dot{q}_i$ .

$$p_r = \mu \dot{r} \quad p_\theta = \mu r^2 \dot{\theta} \quad p_\phi = \mu r^2 \sin^2 \theta \dot{\phi}$$

The Hamiltonian is

$$\begin{aligned} \mathcal{H}_r &= \left( p_r \dot{r} + p_\theta \dot{\theta} + p_\phi \dot{\phi} \right) - \mathcal{L}_r \\ &= \frac{1}{2} \mu \left[ \dot{r}^2 + r^2 \left( \dot{\theta}^2 + \dot{\phi}^2 \sin^2 \theta \right) \right] - \frac{G m_1 m_2}{r} \end{aligned}$$

Alternatively, we can get this Hamiltonian (which is the total energy) by recognizing that the kinetic part of the Lagrangian is quadratic and the potential is not velocity dependent.

To find the Hamiltonian equations of motion, we first express  $\mathcal{H}_r$  in terms of the canonical momenta:

$$\mathcal{H}_r = \frac{p_r^2}{2\mu} + \frac{p_\theta^2}{2\mu r^2} + \frac{p_\phi^2}{2\mu r^2 \sin^2 \theta} - \frac{G m_1 m_2}{r}$$

With this, we find

$$\dot{r} = \frac{\partial \mathcal{H}_r}{\partial p_r} = \frac{p_r}{\mu}; \quad \dot{\theta} = \frac{\partial \mathcal{H}_r}{\partial p_\theta} = \frac{p_\theta}{\mu r^2}; \quad \dot{\phi} = \frac{p_\phi}{\mu r^2 \sin^2 \theta}$$

and

$$\dot{p}_r = -\frac{\partial \mathcal{H}_r}{\partial r} = \frac{p_r^2}{\mu r^3} + \frac{p_\phi^2}{\mu r^3 \sin^2 \theta} - \frac{G m_1 m_2}{r^2}; \quad \dot{p}_\theta = -\frac{\partial \mathcal{H}_r}{\partial \theta} = \frac{p_\phi^2 \cos \theta}{\mu r^2 \sin^3 \theta}; \quad \dot{p}_\phi = -\frac{\partial \mathcal{H}_r}{\partial \phi} = 0.$$

- (d) Mathematica code:

```
(* Problem 1 *)

(* KE, PE, and Lagrangian *)

In[1]:= KE = (\[Mu]/2)*(D[x[t], t]^2 + D[y[t], t]^2 + D[z[t], t]^2);
In[2]:= PE = -G*m1*m2/r[t];
In[3]:= L = KE - PE

Out[3]= (G m1 m2)/r[t] +
1/2 \[Mu] (Derivative[1][x][t]^2 + Derivative[1][y][t]^2 +
Derivative[1][z][t]^2)

In[4]:= L =
L /. {x[t] -> r[t]*Sin[\[Theta][t]]*Cos[\[Phi][t]],
y[t] -> r[t]*Sin[\[Theta][t]]*Sin[\[Phi][t]],
z[t] -> r[t]*Cos[\[Theta][t]],
x'[t] -> D[r[t]*Sin[\[Theta][t]]*Cos[\[Phi][t]], t],
y'[t] -> D[r[t]*Sin[\[Theta][t]]*Sin[\[Phi][t]], t],
z'[t] -> D[r[t]*Cos[\[Theta][t]], t]} // FullSimplify

Out[4]= 1/2 ((2 G m1 m2)/
r[t] + \[Mu] Derivative[1][r][t]^2 + \[Mu] r[t]^2
(Derivative[1][\[Theta][t]]^2 +
Sin[\[Theta][t]^2 Derivative[1][\[Phi][t]]^2))

In[5]:= (* The 'r' equation *)
In[6]:= D[D[L, r'[t]], t] // FullSimplify
Out[6]= \[Mu] (r^[Prime]\[Prime])[t]
In[7]:= D[L, r[t]] // FullSimplify
```

```

Out[7]= -((G m1 m2)/
r[t]^2) + \[Mu] r[
t] (Derivative[1][\[Theta]][t]^2 +
Sin\[Theta][t]^2 Derivative[1][\[Phi]][t]^2)

In[8]:= (* The 'Theta' equation *)

In[9]:= D[D[L, \[Theta]'[t]], t] // FullSimplify

Out[9]= \[Mu] r[
t] (2 Derivative[1][r][t] Derivative[1][\[Theta]][t] +
r[t] (\[Theta]^\[Prime]\[Prime])[t])

In[10]:= D[L, \[Theta][t]] // FullSimplify

Out[10]= \[Mu] Cos\[Theta][t] r[t]^2 Sin\[Theta][t] Derivative[
1][\[Phi]][t]^2

(* The 'Phi' equation *)

In[11]:= D[D[L, \[Phi]'[t]], t] // FullSimplify

Out[11]= \[Mu] r[
t] Sin\[Theta][
t] (2 (Sin\[Theta][t] Derivative[1][r][t] +
Cos\[Theta][t] r[t] Derivative[1][\[Theta]][t]) Derivative[
1][\[Phi]][t] +
r[t] Sin\[Theta][t] (\[Phi]^\[Prime]\[Prime])[t])

In[12]:= D[L, \[Phi][t]] // FullSimplify

Out[12]= 0

(* Canonical momenta *)

In[13]:= pr[t] = D[L, r'[t]]

Out[13]= \[Mu] Derivative[1][r][t]

In[14]:= p\[Theta][t] = D[L, \[Theta]'[t]]

Out[14]= \[Mu] r[t]^2 Derivative[1][\[Theta]][t]

In[15]:= p\[Phi][t] = D[L, \[Phi]'[t]]

Out[15]= \[Mu] r[t]^2 Sin\[Theta][t]^2 Derivative[1][\[Phi]][t]

(* Lagrangian to Hamiltonian *)

In[17]:= H = (pr[t]*r'[t] + p\[Theta][t]*\[Theta]'[t] +
p\[Phi][t]*\[Phi]'[t]) - L // Expand

Out[17]= -((G m1 m2)/r[t]) + 1/2 \[Mu] Derivative[1][r][t]^2 +
1/2 \[Mu] r[t]^2 Derivative[1][\[Theta]][t]^2 +
1/2 \[Mu] r[t]^2 Sin\[Theta][t]^2 Derivative[1][\[Phi]][t]^2

In[63]:= (* Velocities: new instances of P to put in Hamiltonian *)

In[18]:= velocities =
Solve[{Pr[t] == D[L, r'[t]], P\[Theta][t] == D[L, \[Theta]'[t]],
P\[Phi][t] == D[L, \[Phi]'[t]]}, {r'[t], \[Theta]'[t], \[Phi]'[
t]}][[1]]

Out[18]= {Derivative[1][r][t] -> Pr[t]/\[Mu],
Derivative[1][\[Theta]][t] -> P\[Theta][t]/(\[Mu] r[t]^2),
Derivative[1][\[Phi]][t] -> (
Csc\[Theta][t]^2 P\[Phi][t])/(\[Mu] r[t]^2)}

(* Write Hamiltonian in terms of momenta: *)

In[19]:= H = H /. velocities // Expand

Out[19]= Pr[t]^2/(2 \[Mu]) + P\[Theta][t]^2/(2 \[Mu] r[t]^2) + (
Csc\[Theta][t]^2 P\[Phi][t]^2)/(2 \[Mu] r[t]^2) - (G m1 m2)/r[t]

(*Hamiltonian EOMs*)

In[20]:= D[r[t], t] == D[H, Pr[t]]

Out[20]= Derivative[1][r][t] == Pr[t]/\[Mu]

```

```

In[21]:= D[\[Theta][t], t] == D[H, P\[Theta][t]]

Out[21]= Derivative[1][\[Theta]][t] == P\[Theta][t]/(\[Mu] r[t]^2)

In[22]:= D\[Phi][t], t] == D[H, P\[Phi][t]]

Out[22]= Derivative[1][\[Phi]][t] == (
Csc[\[Theta][t]]^2 P\[Phi][t])/(\[Mu] r[t]^2)

In[26]:= D[Pr[t], t] == -D[H, r[t]] // Expand

Out[26]= Derivative[1][Pr][t] ==
P\[Theta][t]^2/(\[Mu] r[t]^3) + (
Csc[\[Theta][t]]^2 P\[Phi][t]^2)/(\[Mu] r[t]^3) - (G m1 m2)/r[t]^2

In[27]:= D[P\[Theta][t], t] == -D[H, \[Theta][t]] // Expand

Out[27]= Derivative[1][P\[Theta]][t] == (
Cot[\[Theta][t]] Csc[\[Theta][t]]^2 P\[Phi][t]^2)/(\[Mu] r[t]^2)

In[83]:= D[P\[Phi][t], t] == -D[H, \[Phi][t]] // Expand

Out[83]= Derivative[1][P\[Phi]][t] == 0

```

## 2. Double Pendulum in a Plane with Gravity

(a) In rectangular coordinates:

$$\begin{aligned}\mathcal{L} &= T - U \\ &= \frac{1}{2}m_1(\dot{x}_1^2 + \dot{y}_1^2) + \frac{1}{2}m_2(\dot{x}_2^2 + \dot{y}_2^2) - m_1gy_1 - m_2gy_2.\end{aligned}$$

With

$$\begin{aligned}x_1 &= l_1 \sin \theta_1; & x_2 &= l_2 \sin \theta_2 + l_1 \sin \theta_1 \\ y_1 &= -l_1 \cos \theta_1; & y_2 &= -l_2 \cos \theta_2 - l_1 \cos \theta_1\end{aligned}$$

we have

$$\mathcal{L} = \frac{1}{2}(m_1 + m_2)l_1^2\dot{\theta}_1^2 + \frac{1}{2}l_1^2m_2\dot{\theta}_2^2 + l_1l_2m_2\cos(\theta_1 - \theta_2)\dot{\theta}_1\dot{\theta}_2 + gl_1(m_1 + m_2)\cos \theta_1 + gl_2m_2\cos \theta_2$$

The equations of motion are:

$$\begin{aligned}\frac{d}{dt}\left(\frac{\partial \mathcal{L}}{\partial \dot{\theta}_1}\right) &= \frac{\partial \mathcal{L}}{\partial \theta_1} \implies l_2m_2\sin(\theta_1 - \theta_2)\dot{\theta}_2^2 + (m_1 + m_2)(g\sin \theta_1 + l_1\ddot{\theta}_1) + l_2m_2\cos(\theta_1 - \theta_2)\ddot{\theta}_2 = 0 \\ \frac{d}{dt}\left(\frac{\partial \mathcal{L}}{\partial \dot{\theta}_2}\right) &= \frac{\partial \mathcal{L}}{\partial \theta_2} \implies g\sin \theta_2 - l_1\sin(\theta_1 - \theta_2)\dot{\theta}_1^2 + l_1\cos(\theta_1 - \theta_2)\ddot{\theta}_1 + l_2\ddot{\theta}_2 = 0.\end{aligned}$$

(b) Now take  $m_1 = m_2 = m$ . Following a similar procedure as before, we first find the canonical momenta using  $p_i = \partial \mathcal{L} / \partial \dot{q}_i$ , then find the Hamiltonian by Legendre-transforming the Lagrangian. Equivalently, we can simply take the total energy, as the kinetic part of the Lagrangian is quadratic and the potential is velocity independent.

$$\mathcal{H} = l_1^2m\dot{\theta}_1^2 + \frac{1}{2}l_2^2m\dot{\theta}_2^2 + l_1l_2m\cos(\theta_1 - \theta_2)\dot{\theta}_1\dot{\theta}_2 - 2gl_1m\cos \theta_1 - gl_2m\cos \theta_2$$

To write  $\mathcal{H}$  in terms of the canonical momenta, we need to find how they are related to the velocities. Setting  $p_{\theta_i} = \partial \mathcal{L} / \partial \dot{\theta}_i$  we find

$$\dot{\theta}_1 = \frac{-l_2p_{\theta_1} + l_1\cos(\theta_1 - \theta_2)p_{\theta_2}}{l_1^2l_2m[\cos^2(\theta_1 - \theta_2) - 2]} \quad \dot{\theta}_2 = \frac{l_2\cos(\theta_1 - \theta_2)p_{\theta_1} - 2l_1p_{\theta_2}}{l_1l_2^2m[\cos^2(\theta_1 - \theta_2) - 2]}$$

In terms of  $p_{\theta_1}$  and  $p_{\theta_2}$ , the Hamiltonian is

$$\mathcal{H} = -\frac{l_1^2\left(g l_2^2 m^2 [\cos(2(\theta_1 - \theta_2)) - 3](2l_1 \cos \theta_1 + l_2 \cos \theta_2 + 22p_{\theta_2}^2) - 2l_1 l_2 p_{\theta_1} p_{\theta_2} \cos(\theta_1 - \theta_2) + l_2^2 p_{\theta_1}^2\right)}{l_1^2 l_2^2 m [\cos(2(\theta_1 - \theta_2)) - 3]}$$

The remain equations of motion are the “ $\dot{p}$ ” equations:

$$\begin{aligned}\dot{p}_{\theta_1} &= \frac{-2gl_1^3l_2^2m^2\sin \theta_1[\cos(2(\theta_1 - \theta_2)) - 3]^2 + 2\sin(2(\theta_1 - \theta_2))(2l_1^2p_{\theta_2}^2 + l_2^2p_{\theta_1}^2)}{l_1^2l_2^2m[\cos(2(\theta_1 - \theta_2)) - 3]^2} \\ &\quad + \frac{-2l_1l_2p_{\theta_1}p_{\theta_2}\sin(\theta_1 - \theta_2)(\cos(2(\theta_1 - \theta_2)) + 5)}{l_1^2l_2^2m[\cos(2(\theta_1 - \theta_2)) - 3]^2}\end{aligned}$$

and

$$\begin{aligned}\dot{p}_{\theta_2} &= gl_2m\sin \theta_2 \\ &\quad + \frac{2\sin(\theta_1 - \theta_2)\left(-4l_1^2p_{\theta_2}^2\cos(\theta_1 - \theta_2) + l_1l_2p_{\theta_1}p_{\theta_2}[\cos(2(\theta_1 - \theta_2)) + 5] - 2l_2^2p_{\theta_1}^2\cos(\theta_1 - \theta_2)\right)}{l_1^2l_2^2m[\cos(2(\theta_1 - \theta_2)) - 3]^2}\end{aligned}$$

(c) Mathematica code:

```
(* Problem 2 *)

(* KE, PE, and Lagrangian *)

In[1]:= KE = (1/2)*m1*(D[x1[t], t]^2 + D[y1[t], t]^2) + (1/2)*
m2*(D[x2[t], t]^2 + D[y2[t], t]^2);

In[2]:= PE = m1*g*y1[t] + m2*g*y2[t];

In[3]:= L = KE - PE;

In[4]:= L = L /. {
x1[t] -> l1*Sin[[Theta]1[t]],
y1[t] -> -l1*Cos[[Theta]1[t]],
x2[t] -> l2*Sin[[Theta]2[t]] + l1*Sin[[Theta]1[t]],
y2[t] -> -l2*Cos[[Theta]2[t]] - l1*Cos[[Theta]1[t]],
x1'[t] -> D[l1*Sin[[Theta]1[t]], t],
y1'[t] -> D[-l1*Cos[[Theta]1[t]], t],
x2'[t] -> D[l2*Sin[[Theta]2[t]] + l1*Sin[[Theta]1[t]], t],
y2'[t] -> D[-l2*Cos[[Theta]2[t]] - l1*Cos[[Theta]1[t]], t]} //
FullSimplify

Out[4]= 1/2 (2 g (l1 (m1 + m2) Cos[[Theta]1[t]] +
l2 m2 Cos[[Theta]2[t]]) +
l1^2 (m1 + m2) Derivative[1][[Theta]1][t]^2 +
2 l1 l2 m2 Cos[[Theta]1[t] - [Theta]2[t]] Derivative[
1][[Theta]1][t] Derivative[1][[Theta]2][t] +
l2^2 m2 Derivative[1][[Theta]2][t]^2)

(* Lagrangian EOMs *)

In[5]:= D[D[L, [[Theta]1'[t]], t] == D[L, [[Theta]1[t]] // FullSimplify

Out[5]= l1 (l2 m2 Sin[[Theta]1[t] - [Theta]2[t]] Derivative[
1][[Theta]2][
t]^2 + (m1 + m2) (g Sin[[Theta]1[t]] +
l1 ([Theta]1^[Prime][Prime])[t]) +
l2 m2 Cos[[Theta]1[t] - [Theta]2[
t]] ([Theta]2^[Prime][Prime])[t]) == 0

In[6]:= D[D[L, [[Theta]2'[t]], t] == D[L, [[Theta]2[t]] // FullSimplify

Out[6]= l2 m2 (g Sin[[Theta]2[t]] -
l1 Sin[[Theta]1[t] - [Theta]2[t]] Derivative[1][[Theta]1][
t]^2 + l1 Cos[[Theta]1[t] - [Theta]2[
t]] ([Theta]1^[Prime][Prime])[t] +
l2 ([Theta]2^[Prime][Prime])[t]) == 0

(* Take m1 = m2 = m *)

In[7]:= D[D[L, [[Theta]1'[t]], t] == D[L, [[Theta]1[t]] /. {m1 -> m,
m2 -> m} // FullSimplify

Out[7]= l1 m (2 g Sin[[Theta]1[t]] +
l2 Sin[[Theta]1[t] - [Theta]2[t]] Derivative[1][[Theta]2][
t]^2 + 2 l1 ([Theta]1^[Prime][Prime])[t] +
l2 Cos[[Theta]1[t] - [Theta]2[t]] ([Theta]2^[Prime][Prime])[
t]) == 0

In[8]:= D[D[L, [[Theta]2'[t]], t] == D[L, [[Theta]2[t]] /. {m1 -> m,
m2 -> m} // FullSimplify

Out[8]= l2 m (g Sin[[Theta]2[t]] -
l1 Sin[[Theta]1[t] - [Theta]2[t]] Derivative[1][[Theta]1][
t]^2 + l1 Cos[[Theta]1[t] - [Theta]2[
t]] ([Theta]1^[Prime][Prime])[t] +
l2 ([Theta]2^[Prime][Prime])[t]) == 0

(*Hamiltonian*)

In[9]:= H = (D[L, [[Theta]1'[t]]*D[[Theta]1[t], t] +
D[L, [[Theta]2'[t]]*D[[Theta]2[t], t]) - L /. {m1 -> m,
m2 -> m} // Expand

Out[9]= -2 g l1 m Cos[[Theta]1[t]] - g l2 m Cos[[Theta]2[t]] +
l1^2 m Derivative[1][[Theta]1][t]^2 +
l1 l2 m Cos[[Theta]1[t] - [Theta]2[t]] Derivative[1][[Theta]1][
t] Derivative[1][[Theta]2][t] +
1/2 l2^2 m Derivative[1][[Theta]2][t]^2
```

```

(*solve for velocities in terms of momenta to write H in terms of \
momenta*)

In[10]:= D[L, \[Theta]1'[t]] /. {m1 -> m, m2 -> m} // FullSimplify

Out[10]= 11 m (2 11 Derivative[1][\[Theta]1][t] +
12 Cos[\[Theta]1[t] - \[Theta]2[t]] Derivative[1][\[Theta]2][t])

In[11]:= velocities =
Solve[{P\[Theta]1[t] == D[L, \[Theta]1'[t]],
P\[Theta]2[t] == D[L, \[Theta]2'[t]]}, {\[Theta]1'[
t], \[Theta]2'[t]}][[1]] /. {m1 -> m, m2 -> m} // FullSimplify

Out[11]= {Derivative[1][\[Theta]1][t] -> (-12 P\[Theta]1[t] +
11 Cos[\[Theta]1[t] - \[Theta]2[t]] P\[Theta]2[t])/
(11^2 12 m (-2 + Cos[\[Theta]1[t] - \[Theta]2[t]]^2)),
Derivative[1][\[Theta]2][t] -> (
12 Cos[\[Theta]1[t] - \[Theta]2[t]] P\[Theta]1[t] -
2 11 P\[Theta]2[t])/
(11 12^2 m (-2 + Cos[\[Theta]1[t] - \[Theta]2[t]]^2))}

In[12]:= H = H /. velocities // FullSimplify

Out[12]= -((12^2 P\[Theta]1[t]^2 -
2 11 12 Cos[\[Theta]1[t] - \[Theta]2[t]] P\[Theta]1[
t] P\[Theta]2[t] +
11^2 (g 12^2 m^2 (-3 +
Cos[2 (\[Theta]1[t] - \[Theta]2[t]))] (2 11 Cos[\[Theta]1[
t]] + 12 Cos[\[Theta]2[t]])) +
2 P\[Theta]2[t]^2))/(11^2 12^2 m (-3 +
Cos[2 (\[Theta]1[t] - \[Theta]2[t]))]))

(* Hamiltonian EOMs *)

In[16]:= Solve[P\[Theta]1'[t] == -D[H, \[Theta]1[t]],
P\[Theta]1'[t]][[1]] // FullSimplify

Out[16]= {Derivative[1][P\[Theta]1][
t] -> (-2 g 11^3 12^2 m^2 (-3 +
Cos[2 (\[Theta]1[t] - \[Theta]2[t]))]^2 Sin[\[Theta]1[t]] -
2 11 12 (5 + Cos[2 (\[Theta]1[t] - \[Theta]2[t]))] P\[Theta]1[
t] P\[Theta]2[t] Sin[\[Theta]1[t] - \[Theta]2[t]] +
2 (12^2 P\[Theta]1[t]^2 + 2 11^2 P\[Theta]2[t]^2) Sin[
2 (\[Theta]1[t] - \[Theta]2[t]))]/(11^2 12^2 m (-3 +
Cos[2 (\[Theta]1[t] - \[Theta]2[t]))]^2))}

In[17]:= Solve[P\[Theta]2'[t] == -D[H, \[Theta]2[t]],
P\[Theta]2'[t]][[1]] // FullSimplify

Out[17]= {Derivative[1][P\[Theta]2][
t] -> (2 (-2 12^2 Cos[\[Theta]1[t] - \[Theta]2[t]] P\[Theta]1[
t]^2 + 11 12 (5 +
Cos[2 (\[Theta]1[t] - \[Theta]2[t]))] P\[Theta]1[
t] P\[Theta]2[t] -
4 11^2 Cos[\[Theta]1[t] - \[Theta]2[t]] P\[Theta]2[
t]^2) Sin[\[Theta]1[t] - \[Theta]2[t]]/(11^2 12^2 m (-3 +
Cos[2 (\[Theta]1[t] - \[Theta]2[t]))]^2) -
g 12 m Sin[\[Theta]2[t]]}

In[66]:= \[Theta]1'[t] == D[H, P\[Theta]1[t]] // FullSimplify

Out[66]= 11 Derivative[1][\[Theta]1][t] == (
2 12 P\[Theta]1[t] -
2 11 Cos[\[Theta]1[t] - \[Theta]2[t]] P\[Theta]2[t])/
(3 11 12 m - 11 12 m Cos[2 \[Theta]1[t] - 2 \[Theta]2[t]])

In[67]:= \[Theta]2'[t] == D[H, P\[Theta]2[t]] // FullSimplify

Out[67]= Derivative[1][\[Theta]2][t] == (
2 (12 Cos[\[Theta]1[t] - \[Theta]2[t]] P\[Theta]1[t] -
2 11 P\[Theta]2[t])/
(11 12^2 m (-3 + Cos[2 (\[Theta]1[t] - \[Theta]2[t]))])

```

### 3. Point Mass on a Hoop: Goldstein Ch.2 Problem #18.

By the geometry of the problem, we the system may be described by one (spherical) coordinate  $\theta$  defined as usual.  $r$  is fixed at  $r = a$  and  $\phi(t) = \omega t$ , where  $\omega$  is fixed.

$$(x, y, z) = (a \sin \theta \cos \omega t, a \sin \theta \sin \omega t, a \cos \theta).$$

With this, the Lagrangian is

$$\begin{aligned} \mathcal{L} &= \frac{1}{2}m [\dot{x}^2 + \dot{y}^2 + \dot{z}^2] - mgz \\ &= \frac{1}{2}am [-2g \cos \theta + a\omega^2 \sin^2 \theta + a\dot{\theta}^2]. \end{aligned}$$

There is only one Lagrangian equation of motion:

$$\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{\theta}} \right) = \frac{\partial \mathcal{L}}{\partial \theta} \implies a\ddot{\theta} = (g + a\omega^2 \cos \theta) \sin \theta.$$

It is clear from the functional form of  $\mathcal{L}$  that a constant of motion is  $r$  since  $r = a$  fixed. Moreover, since  $\mathcal{L}$  is time-independent, energy is conserved, therefore is also a constant of motion. In this case, the energy function is identically the Hamiltonian because the Lagrangian is quadratic in the kinetic part and velocity-independent in the potential part. Thus, we have

$$\text{const} = \mathcal{H} = \frac{\partial \mathcal{L}}{\partial \dot{\theta}} \dot{\theta} - \mathcal{L} = \frac{1}{2}am (a\dot{\theta}^2 - a\omega^2 \sin^2 \theta + 2g \cos \theta).$$

We now want to find the critical value  $\omega_0$  described in the problem statement. Since the particle is stationary,  $\dot{\theta} = 0$ . The particle is at a stationary point exactly when it is at a local minimum of the effective potential. From the Hamiltonian, we can read off the effective potential as

$$V(\theta) = \frac{1}{2}am (-a\omega^2 \sin^2 \theta + 2g \cos \theta).$$

We find that

$$0 = \frac{\partial V(\theta)}{\partial \theta} = \frac{1}{2}am (-a\omega^2 \sin 2\theta - 2g \sin \theta) = am \sin \theta (-a\omega^2 \cos \theta - g)$$

so the stationary points are  $\theta = 0, \pi$  or  $\theta = \arccos(-\omega_0^2/\omega^2)$  where  $\omega_0 \equiv \sqrt{g/a}$ . Of these three points,  $\theta = 0$  is always unstable since it for  $\theta \approx 0$   $V(\theta)$  looks like  $\cos \theta$ , so the particle will move away from  $\theta = 0$ .

When  $\omega_0 \geq \omega$ , the only equilibrium is  $\theta = \pi$  since  $\partial V(\theta)/\partial \theta \leq 0$  for all  $\theta \in [0, \pi]$ . When  $\omega_0 \leq \omega$ , the equilibrium point becomes  $\theta = \theta_0 = \arccos(-\omega_0^2/\omega^2)$  because  $\partial V/\partial \theta \geq 0$  for  $\theta \geq \theta_0$ .

Mathematica code:

```
(*Problem 3*)
In[20]:= KE = (m/2)*(D[x[t], t]^2 + D[y[t], t]^2 + D[z[t], t]^2);
In[21]:= PE = m*g*z[t];
In[22]:= L = KE - PE
Out[22]= -g m z[t] +
1/2 m (Derivative[1][x][t]^2 + Derivative[1][y][t]^2 +
Derivative[1][z][t]^2)
(*Lagrangian*)
In[23]:= L = L /. {x[t] -> a*Sin[Theta[t]]*Cos[Omega*t],
y[t] -> a*Sin[Theta[t]]*Sin[Omega*t],
z[t] -> a*Cos[Theta[t]],
x'[t] -> D[a*Sin[Theta[t]]*Cos[Omega*t], t],
y'[t] -> D[a*Sin[Theta[t]]*Sin[Omega*t], t],
z'[t] -> D[a*Cos[Theta[t]], t]} // FullSimplify
Out[23]= 1/2 a m (-2 g Cos[Theta[t]] + a Omega^2 Sin[Theta[t]]^2 +
a Derivative[1][Theta][t]^2)
```



```

In[19]:= (*Hamiltonian*)
In[25]:= H = D[L, \[Theta]'[t]]*\[Theta]'[t] - L // FullSimplify
Out[25]= 1/2 a m (2 g Cos[\[Theta][t]] - a \[Omega]^2 Sin[\[Theta][t]]^2 +
a Derivative[1][\[Theta]][t]^2)
In[26]:= 1/2 a m (2 g Cos[\[Theta][t]] - a \[Omega]^2 Sin[\[Theta][t]]^2 +
a Derivative[1][\[Theta]][t]^2) // TeXForm
Out[26]//TeXForm=
\frac{1}{2} a m \left(a \theta '(t)^2-a \omega ^2 \sin ^2(\theta
(t))+2 g \cos (\theta (t))\right)
(*Lagrangian EOM*)
In[88]:= FullSimplify[D[D[L, \[Theta]'[t]], t] == D[L, \[Theta][t]]]
Out[88]= a^2 m (\[Theta]^\[Prime][\[Prime]][t] ==
a m (g + a \[Omega]^2 Cos[\[Theta][t]]) Sin[\[Theta][t]]

```

#### 4. Spring System on a Plane

(a) The Lagrangian in Cartesian coordinates is

$$\mathcal{L} = \frac{1}{2}m_1(\dot{x}_1^2 + \dot{y}_1^2) + \frac{1}{2}m_2(\dot{x}_2^2 + \dot{y}_2^2) - \frac{1}{2}k\left(\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} - b\right)^2$$

(b) New coordinates are  $(x_{\text{COM}}, y_{\text{COM}}, r, \theta)$ , where  $(x_{\text{COM}}, y_{\text{COM}})$  describes the position of the center of mass of the system, while  $(r, \theta)$  together describe the relative position vector of the two masses. These new coordinates are defined by

$$x_{\text{COM}} = \frac{m_1x_1 + m_2x_2}{m_1 + m_2}; \quad y_{\text{COM}} = \frac{m_1y_1 + m_2y_2}{m_1 + m_2}; \quad r = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}; \quad \theta = \arctan \frac{y_2 - y_1}{x_2 - x_1}$$

The Lagrangian in these coordinates is:

$$\mathcal{L} = -\frac{k}{2}(b - r)^2 + \frac{m_1m_2(\dot{r}^2 + r^2\dot{\theta}^2)}{2(m_1 + m_2)} + \frac{1}{2}(m_1 + m_2)(\dot{x}_{\text{COM}}^2 + \dot{y}_{\text{COM}}^2)$$

The equations of motion are:

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{x}_{\text{COM}}} = \frac{\partial \mathcal{L}}{\partial x_{\text{COM}}} \implies \ddot{x}_{\text{COM}} = 0$$

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{y}_{\text{COM}}} = \frac{\partial \mathcal{L}}{\partial y_{\text{COM}}} \implies \ddot{y}_{\text{COM}} = 0$$

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{r}} = \frac{\partial \mathcal{L}}{\partial r} \implies \ddot{r} = \frac{k(m_1 + m_2)(b - r)}{m_1m_2} + r\dot{\theta}^2$$

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{\theta}} = \frac{\partial \mathcal{L}}{\partial \theta} \implies \ddot{\theta} = -\frac{2\dot{r}\dot{\theta}}{r}$$

(c) The cyclic coordinates are  $x_{\text{COM}}$ ,  $y_{\text{COM}}$ , and  $\theta$ , since  $\mathcal{L}$  does not explicitly depend on them. The conserved generalized momenta are thus

$$p_{x_{\text{COM}}} = \frac{\partial \mathcal{L}}{\partial \dot{x}_{\text{COM}}} = (m_1 + m_2)\dot{x}_{\text{COM}}; \quad p_{y_{\text{COM}}} = \frac{\partial \mathcal{L}}{\partial \dot{y}_{\text{COM}}} = (m_1 + m_2)\dot{y}_{\text{COM}}; \quad p_{\theta} = \frac{\partial \mathcal{L}}{\partial \dot{\theta}} = \frac{m_1m_2}{m_1 + m_2}r^2\dot{\theta}.$$

We hope to show that there is a solution which rotates but does not oscillate. To this end, we will use energy conservation as a constraint on the possible dynamics of the system. In particular, since we are interested in how  $r(t)$ , we will look at the  $r$ -equation and the force which dictates  $r$ 's behavior. Letting  $\mu = m_1m_2/(m_1 + m_2)$ ,

$$\mu\ddot{r} = k(b - r) + r\dot{\theta}^2.$$

Consider this a one-body problem with Newton's second law  $\vec{F} = m\vec{a}$ . This force is associated with the potential  $V(r)$  where

$$V(r) = \frac{1}{2}k(b - r)^2 - \frac{1}{2}\mu r^2\dot{\theta}^2.$$

On the other hand, we may compute the total energy of the system. After recognizing that the Lagrangian is quadratic in the kinetic terms and has velocity independent potentials, the total energy is the Hamiltonian:

$$\mathcal{H} = \frac{1}{2}\mu\dot{r}^2 - \frac{1}{2}\mu r^2\dot{\theta}^2 + \frac{1}{2}k(b - r)^2 + \text{constant}$$

where we have used the fact that  $\ddot{x}_{\text{COM}} = \ddot{y}_{\text{COM}} = 0$ . Since  $\mathcal{H}$  has no explicit time dependence,  $d\mathcal{H}/dt = 0$ , and thus energy is conserved. In particular, when we compare  $\mathcal{H}$  to  $V(r)$ , we find that

$$\frac{1}{2}\mu\dot{r}^2 + V(r) = \text{constant}$$

Now, suppose  $\dot{r}(0) = 0$  (does not oscillate) but  $\dot{\theta} \neq 0$  (rotates). If  $k > \mu\dot{\theta}^2$  then  $\partial^2 V/\partial r > 0$ , ensuring a stable equilibrium. Thus, if we start at  $r(0) = r_0 = bk/(k - \mu\dot{\theta}^2)$  where  $V(r_0) = 0$  then  $\dot{r}$  remains at zero for all  $t > 0$ . Therefore, there exists a solution which rotates but does not oscillate. When  $\dot{\theta}$  gets large, i.e., when  $\dot{\theta} > \sqrt{k/\mu}$ , then the concavity of  $V(r)$  changes to negative and the stable equilibrium no longer exists.  $r$  diverges to infinite in this case.

(d) Mathematica code:

```
(*Problem 4*)

In[50]:= KE = (1/2)*(m1 + m2)*(D[xCOM[t], t]^2 +
D[yCOM[t], t]^2) + (1/2)*(m1*
m2)*(D[r[t], t]^2 + r[t]^2*D[\[Theta][t], t]^2)/(m1 + m2);

In[51]:= PE = (k/2)*(r[t] - b)^2;

In[52]:= L = KE - PE;

(*Lagrangian*)

In[53]:= L // FullSimplify

Out[53]= 1/2 (-k (b - r[t])^2 + (m1 + m2) (Derivative[1][xCOM][t]^2 +
Derivative[1][yCOM][t]^2) + (
m1 m2 (Derivative[1][r][t]^2 +
r[t]^2 Derivative[1][\[Theta][t]^2))/(m1 + m2))

(*Lagrangian EOMs*)

(*xCOM equation*)

In[54]:= Solve[FullSimplify[D[D[L, xCOM'[t]], t] == D[L, xCOM[t]]],
xCOM''[t]][[1]] // Expand

Out[54]= {(xCOM^[Prime][Prime])[t] -> 0}

(*yCOM equation*)

In[55]:= Solve[FullSimplify[D[D[L, yCOM'[t]], t] == D[L, yCOM[t]]],
yCOM''[t]][[1]] // Expand

Out[55]= {(yCOM^[Prime][Prime])[t] -> 0}

(*r equation*)

In[65]:= Solve[FullSimplify[D[D[L, r'[t]], t] == D[L, r[t]]],
r''[t]][[1]] // FullSimplify

Out[65]= {(r^[Prime][Prime])[t] -> (k (m1 + m2) (b - r[t]))/(
m1 m2) + r[t] Derivative[1][\[Theta][t]^2}

In[15]:= (*theta equation*)

In[57]:= Solve[
FullSimplify[
D[D[L, \[Theta]'[t]], t] == D[L, \[Theta][t]], \[Theta]''[t]][[
1]] // Expand

Out[57]= {(\[Theta]^[Prime][Prime])[t] -> -((
2 Derivative[1][r][t] Derivative[1][\[Theta][t])/r[t])}

(*Cyclic coordinates*)

In[58]:= D[L, xCOM[t]]

Out[58]= 0

In[59]:= D[L, yCOM[t]]

Out[59]= 0
```

```

In[60]:= D[L, \[Theta][t]]
Out[60]= 0

(*Conserved generalized momenta*)

In[61]:= D[L, xCOM'[t]] // Simplify
Out[61]= (m1 + m2) Derivative[1][xCOM][t]

In[62]:= D[L, yCOM'[t]] // Simplify
Out[62]= (m1 + m2) Derivative[1][yCOM][t]

In[63]:= D[L, \[Theta]'[t]] // Simplify
Out[63]= (m1 m2 r[t]^2 Derivative[1][\[Theta]][t])/(m1 + m2)

```

## 6. Routhian Mechanics

We start with the definition of a particular Routhian:

$$R(q_1, \dots, q_n, p_1, \dots, p_s, \dot{q}_{s+1}, \dots, \dot{q}_n, t) = \sum_{k=1}^s p_k \dot{q}_k - \mathcal{L}(q_1, \dots, q_n, \dot{q}_1, \dots, \dot{q}_n, t).$$

For  $i = 1, \dots, n$  we have

$$\boxed{\frac{\partial R}{\partial q_i}} = -\frac{\partial \mathcal{L}}{\partial q_i} = -\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{q}_i} \right) = -\frac{d}{dt} p_i = \boxed{-\dot{p}_i}$$

For  $i = 1, \dots, s$ , we have

$$\boxed{\frac{\partial R}{\partial p_i}} = \dot{q}_i$$

For  $i = s + 1, \dots, n$ , we have

$$\boxed{\frac{\partial R}{\partial \dot{q}_i}} = -\frac{\partial \mathcal{L}}{\partial \dot{q}_i} = \boxed{-p_i} \implies \boxed{\frac{d}{dt} \frac{\partial R}{\partial \dot{q}_i}} = -\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{q}_i} \right) = -\frac{d}{dt} p_i = \boxed{\frac{\partial R}{\partial q_i}}$$

Finally,

$$\frac{\partial R}{\partial t} = -\frac{\partial \mathcal{L}}{\partial t}.$$

## 7. Extra Problem: Equivalent Lagrangians

We claim that

$$\mathcal{L}' = \mathcal{L} + \frac{dF(q, t)}{dt}$$

where  $F$  is a differential function, satisfies the Euler-Lagrange equations.

*Proof.* On the one hand,

$$\frac{d}{dt} \frac{\partial \mathcal{L}'}{\partial \dot{q}} = \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} + \frac{d}{dt} \frac{\partial}{\partial \dot{q}} \frac{dF}{dt} = \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} + \frac{d}{dt} \frac{\partial F}{\partial q} = \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} + \frac{\partial}{\partial q} \left( \frac{\partial F}{\partial q} \right) \dot{q} + \frac{\partial^2 F}{\partial t \partial q},$$

where we have used the fact that  $F$  does not depend on  $\dot{q}$  for the second equality (i.e.,  $dF/dt$  generates a factor of  $\dot{q}$ , which gets taken away by  $\partial/\partial \dot{q}$ ).

On the other hand,

$$\frac{\partial \mathcal{L}'}{\partial q} = \frac{\partial \mathcal{L}}{\partial q} + \frac{\partial}{\partial q} \frac{dF}{dt} = \frac{\partial \mathcal{L}}{\partial q} + \frac{\partial}{\partial q} \left( \frac{\partial F}{\partial q} \right) \dot{q} + \frac{\partial^2 F}{\partial q \partial t}.$$

Since  $\partial^2 F / \partial q \partial t = \partial^2 F / \partial t \partial q$  and that  $\mathcal{L}$  satisfies the Euler-Lagrange equations, we see that

$$\frac{d}{dt} \frac{\partial \mathcal{L}'}{\partial \dot{q}} = \frac{\partial \mathcal{L}'}{\partial q}$$

as desired. □