

# **STATISTICAL INFERENCE**

- A Quick Guide -

Huan Q. Bui

Colby College

PHYSICS & MATHEMATICS  
Statistics

Class of 2021

April 18, 2020

## Preface

Greetings,

This guide is based on SC482: Statistical Inference, taught by Professor Liam O'Brien. The guide consists of lecture notes and material from *Introduction to Mathematical Statistics, 8th edition* by Hogg, McKean, and Craig. A majority of the text will be reading notes and solutions to selected problems.

As this is intended only to be a reference source, I might not be as meticulous with my explanations as I have been in some other guides.

Enjoy!

# Contents

Preface . . . . .	2
<b>1 Special Distributions</b>	<b>5</b>
1.1 The Binomial and Related Distributions . . . . .	6
1.1.1 Negative Binomial & Geometric Distribution . . . . .	6
1.2 Multinomial Distribution . . . . .	7
1.3 Hypergeometric Distribution . . . . .	7
1.4 The Poisson Distribution . . . . .	7
1.5 The $\Gamma, \chi^2, \beta$ distributions . . . . .	8
1.5.1 The $\Gamma$ and exponential distribution . . . . .	8
1.5.2 The $\chi^2$ distribution . . . . .	9
1.5.3 The $\beta$ distribution . . . . .	9
1.6 The Normal distribution . . . . .	10
1.6.1 Contaminated Normal . . . . .	11
1.7 The Multivariate Normal . . . . .	11
1.8 The $t$ - and $F$ -distributions . . . . .	12
1.8.1 The $t$ -distribution . . . . .	12
1.8.2 The $F$ -distribution . . . . .	13
1.8.3 The Student's Theorem . . . . .	14
<b>2 Elementary Statistical Inferences</b>	<b>17</b>
2.1 Sampling & Statistics . . . . .	18
2.1.1 Point estimators . . . . .	18
2.1.2 Histogram estimates of pmfs and pdfs . . . . .	19
2.2 Confidence Intervals . . . . .	21
2.2.1 CI for difference in means . . . . .	22
2.2.2 CI for difference in proportions . . . . .	23
2.3 Order Statistics . . . . .	23
2.3.1 Quantiles . . . . .	24
2.3.2 CI for quantiles . . . . .	24
2.4 Introduction to Hypothesis Testing . . . . .	25
2.5 Additional comments about statistical test . . . . .	26
2.5.1 Observed Significance Level, $p$ -value . . . . .	26
2.6 Chi-Square Tests . . . . .	26
2.7 The Method of Monte Carlo . . . . .	29

2.7.1	Inverse Transform . . . . .	29
2.7.2	Accept-Reject Generation Algorithm . . . . .	30
2.7.3	Evaluating definite integrals . . . . .	30
2.8	Bootstrapping . . . . .	32
2.8.1	Bootstrapping for Hypothesis Testing . . . . .	32
<b>3</b>	<b>Consistency and Limiting Distributions</b>	<b>35</b>
3.1	Convergence in Probability . . . . .	35
3.1.1	Sampling and Statistic . . . . .	36
3.2	Convergence in Distribution . . . . .	36
3.2.1	Bounded in Probability . . . . .	37
3.2.2	$\Delta$ -method . . . . .	37
3.2.3	Moment Generating Function Technique . . . . .	38
3.3	Central Limit Theorem . . . . .	38
<b>4</b>	<b>Maximum Likelihood Methods</b>	<b>41</b>
4.1	Maximum Likelihood Estimation . . . . .	42
4.2	Rao-Cramér Lower Bound and Efficiency . . . . .	45
4.3	Maximum Likelihood Tests . . . . .	50
4.4	Multiparameter Case: Estimation . . . . .	53
4.5	Multiparameter Case: Testing . . . . .	55
4.6	The EM algorithm . . . . .	56
<b>5</b>	<b>Sufficiency</b>	<b>57</b>
5.1	Measures of Quality of Estimators . . . . .	57
5.2	A Sufficient Statistic for a Parameter . . . . .	57
<b>6</b>	<b>Problems</b>	<b>59</b>
6.1	Problem Set 1 . . . . .	60
6.2	Problem Set 2 . . . . .	72
6.3	Problem set 3 . . . . .	86
6.4	Problem set 4 . . . . .	93
6.5	Problem set 5 . . . . .	101
6.6	Problem set 6 . . . . .	113
6.7	Problem set 7 . . . . .	127

**Part 1**

# **Special Distributions**

## 1.1 The Binomial and Related Distributions

If we let the random variable  $X$  equal the number of observed successes in  $n$  independent Bernoulli trials, each with success probability of  $p$ , then  $X$  follows the binomial distribution.

A binomial pmf is given by

$$p(x) = \begin{cases} \binom{n}{x} p^x (1-p)^{n-x} & x = 0, 1, 2, \dots \\ 0, & \text{else} \end{cases} \quad (1.1)$$

Using the binomial expansion formula, we can easily check that

$$\sum_x p(x) = 1 \quad (1.2)$$

The mgf of a binomial distribution is obtained by:

$$M_{\text{bin}}(t) = E[e^{tx}] = \sum_x e^{tx} p(x) = [(1-p) + pe^t]^n \quad \forall t \in \mathbb{R} \quad (1.3)$$

With this, we can find the mean and variance for  $p(x)$ :

$$\mu = M'(0) = n, \quad \sigma^2 = M''(0) = np(1-p) \quad (1.4)$$

**Theorem 1.1.1.** Let  $X_1, X_2, \dots, X_m$  be independent binomial random variables such that  $X_i \sim \text{bin}(n_i, p)$ ,  $i = 1, 2, \dots, m$ . Then

$$Y = \sum_{i=1}^m X_i \sim \text{bin}\left(\sum_{i=1}^m n_i, p\right) \quad (1.5)$$

*Proof:* We prove this via the mgf for  $Y$ . By independence, we have that

$$M_Y(t) = \prod_{i=1}^m (1 - p + pe^t)^{n_i} = (1 - p + pe^t)^{\sum_{i=1}^m n_i} \quad (1.6)$$

The mgf completely determines the distribution which  $Y$  follows, so we're done.  $\square$

### 1.1.1 Negative Binomial & Geometric Distribution

Consider a sequence of independent Bernoulli trials with constant probability  $p$  of success. The random variable  $Y$  which denotes the total number of failures in this sequence before the  $r$ th success follows the negative binomial distribution.

A negative binomial pmf is given by

$$p_Y(t) = \begin{cases} \binom{(y+r-1)}{r-1} p^r (1-p)^y & y = 0, 1, 2, \dots \\ 0, & \text{else} \end{cases} \quad (1.7)$$

The mgf of this distribution is

$$M(t) = p^r [1 - (1-p)e^t]^{-r} \quad (1.8)$$

When  $r = 1$ ,  $Y$  follows the geometric distribution, whose pmf is given by

$$p_Y(y) = p(1-p)^y, \quad y = 0, 1, 2, \dots \quad (1.9)$$

The mgf of this distribution is

$$M(t) = p[1 - (1-p)e^t]^{-1} \quad (1.10)$$

## 1.2 Multinomial Distribution

We won't worry about this for now.

## 1.3 Hypergeometric Distribution

We won't worry about this for now.

## 1.4 The Poisson Distribution

The Poisson distribution gives the probability of observing  $x$  occurrences of some rare events characterized by rate  $\lambda > 0$ . The pmf is given by

$$p(x) = \begin{cases} \frac{\lambda^x e^{-\lambda}}{x!}, & x = 0, 1, 2, \dots \\ 0, & \text{else} \end{cases} \quad (1.11)$$

We say a random parameter with the pmf of the form of  $p(x)$  follows the Poisson distribution with parameter  $\lambda$ .

The mgf of a Poisson distribution is given by

$$M(t) = e^{-\lambda(e^t - 1)} \quad (1.12)$$

From here, we can find the mean and variance:

$$\mu = M'(0) = \lambda, \quad \sigma^2 = M''(0) = \lambda \quad (1.13)$$

**Theorem 1.4.1.** If  $X_1, \dots, X_n$  are independent random variables, each  $X_i \sim \text{Poi}(\lambda_i)$ , then

$$Y = \sum_{i=1}^n X_i \sim \text{Poi}\left(\sum_{i=1}^n \lambda_i\right) \quad (1.14)$$

*Proof:* We once again prove this via the mgf of  $Y$ :

$$M_Y(t) = \prod_{i=1}^n e^{\lambda_i(e^t - 1)} = e^{\sum_{i=1}^n \lambda_i(e^t - 1)} \quad (1.15)$$

□

## 1.5 The $\Gamma, \chi^2, \beta$ distributions

The gamma function of  $\alpha > 0$  is given by

$$\Gamma(\alpha) = \int_0^\infty y^{\alpha-1} e^{-y} dy, \quad (1.16)$$

which gives  $\Gamma(1) = 1$  and  $\Gamma(\alpha) = (\alpha - 1)\Gamma(\alpha - 1)$ .

### 1.5.1 The $\Gamma$ and exponential distribution

A continuous random variable  $X \sim \Gamma(\alpha, \beta)$  where  $\alpha > 0$  and  $\beta > 0$  whenever its pdf is

$$f(x) = \begin{cases} \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} e^{-x/\beta}, & 0 < x < \infty \\ 0, & \text{else} \end{cases} \quad (1.17)$$

The mgf for  $X$  is obtained via the change of variable  $y = x(1 - \beta t)/\beta$ , where  $t < 1/\beta$ :

$$M(t) = \int_0^\infty \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} e^{-x(1-\beta t)/\beta} dx = \frac{1}{(1 - \beta t)^\alpha} \quad (1.18)$$

From here, we can find the mean and variance:

$$\mu = M'(0) = \alpha\beta, \quad \sigma^2 = \alpha\beta^2 \quad (1.19)$$

The  $\Gamma(1, \beta)$  distribution is a special case, and it is called the **exponential distribution** with parameter  $1/\beta$ .

**Theorem 1.5.1.** Let  $X_1, \dots, X_n$  be independent random variables, with  $X_i \sim \Gamma(\alpha_i, \beta)$ . Then

$$Y = \sum_{i=1}^n X_i \sim \Gamma\left(\sum_{i=1}^n \alpha_i, \beta\right) \quad (1.20)$$

*Proof:* Can you guess via which device we prove the statement above?  $\square$

### 1.5.2 The $\chi^2$ distribution

The  $\chi^2$  distribution is a special case of the gamma distribution where  $\alpha = r/2, r \in \mathbb{N}^*$  and  $\beta = 2$ . If a continuous r.v.  $X \sim \chi^2(r)$  then its pdf is

$$f(x) = \begin{cases} \frac{1}{\Gamma(r/2)2^{r/2}} x^{r/2-1} e^{-x/2}, & 0 < x < \infty \\ 0, & \text{else} \end{cases} \quad (1.21)$$

Its mgf is

$$M(t) = (1 - 2t)^{-r/2}, \quad t < \frac{1}{2} \quad (1.22)$$

**Theorem 1.5.2.** Let  $X \sim \chi^2(r)$  and  $k > -r/2$  be given. Then  $E[X^k]$  exists and is given by

$$E[X^k] = \frac{2^k \Gamma(r/2 + k)}{\Gamma(r/2)} \quad (1.23)$$

*Proof:* is proof is purey computational and is left to the reader.  $\square$

From here, we note that all moments of the  $\chi^2$  distribution exist.

**Theorem 1.5.3.** Let  $X_1, \dots, X_n$  be r.v. with  $X_i \sim \chi^2(r_i)$ . Then

$$Y = \sum_{i=1}^n X_i \sim \chi^2\left(\sum_{i=1}^n r_i\right) \quad (1.24)$$

*Proof:* we once again find the mgf for  $Y$ .  $\square$

### 1.5.3 The $\beta$ distribution

The  $\beta$  distribution differs from the other continuous ones we've discussed so far because its support are bounded intervals.

I will skip most of the details here, except mentioning that we can derive the beta distribution from a pair of independent  $\Gamma$  random variables. Suppose  $Y = X_1/(X_1 + X_2)$  where  $X_i \sim \Gamma(\alpha, \beta)$  then the pdf of  $Y$  is that of the beta distribution:

$$g(y) = \begin{cases} \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} y^{\alpha-1} (1-y)^{\beta-1}, & 0 < y < 1 \\ 0, & \text{else} \end{cases} \quad (1.25)$$

The mean and variance of  $Y$  are

$$\mu = \frac{\alpha}{\alpha + \beta}, \quad \sigma^2 = \frac{\alpha\beta}{(\alpha + \beta + 1)(\alpha + \beta)^2} \quad (1.26)$$

## 1.6 The Normal distribution

I have dedicated a large chunk in the [QFT](#) notes to evaluating Gaussian integrals, so I won't go into that here.

$X \sim \mathcal{N}(\mu, \sigma^2)$  whenever its pdf is

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2} \frac{(x-\mu)^2}{\sigma^2}\right), \quad -\infty < x < \infty \quad (1.27)$$

where  $\mu$  and  $\sigma^2$  are the mean and variance of  $X$ , respectively.

The mgf of  $X$  can be obtained via the substitution  $X = \sigma Z + \mu$ :

$$M(t) = \exp\left(\mu t + \frac{1}{2}\sigma^2 t^2\right) \quad (1.28)$$

We note the following correspondence for  $X = \sigma Z + \mu$ :

$$X \sim \mathcal{N}(\mu, \sigma^2) \iff Z \sim \mathcal{N}(0, 1) \quad (1.29)$$

**Theorem 1.6.1.**  $X \sim \mathcal{N}(\mu, \sigma^2) \implies V = (X - \mu)^2 / \sigma^2 \sim \chi^2(1)$ , i.e. a standardized, squared normal follows a chi-square distribution.

*Proof:* The proof isn't too hard. Let us write  $V$  as  $W^2$  and so  $W \sim \mathcal{N}(0, 1)$ . We consider the cdf  $G(v)$  for  $V$ , with  $v \geq 0$ :

$$G(v) = P(W^2 \leq v) = P(-\sqrt{v} \leq W \leq \sqrt{v}) = 2 \int_0^{\sqrt{v}} \frac{1}{\sqrt{2\pi}} e^{-w^2/2} dw \quad (1.30)$$

with  $G(v) = 0$  whenever  $v < 0$ . From here, we can see that the pdf for  $v$ , under the change of notation  $w \rightarrow \sqrt{y}$ , is

$$g(v) = G'(v) = \frac{d}{dv} \left\{ \int_0^v \frac{1}{\sqrt{2\pi}\sqrt{y}} e^{-y/2} dy \right\}, \quad 0 \geq v \quad (1.31)$$

or 0 otherwise. This means

$$g(v) = \begin{cases} \frac{1}{\sqrt{\pi}\sqrt{2}} v^{1/2-1} e^{-v/2}, & 0 < v < \infty \\ 0, & \text{else} \end{cases} \quad (1.32)$$

Using the fact that  $\Gamma(1/2) = \sqrt{\pi}$  and by verifying that  $g(v)$  integrates to unity we show  $V \sim \chi^2(1)$ .  $\square$

**Theorem 1.6.2.** Let  $X_1, \dots, X_n$  be independent r.v. with  $X_i \sim \mathcal{N}(\mu_i, \sigma_i^2)$ . Then for constants  $a_1, \dots, a_n$

$$Y = \sum_{i=1}^n a_i X_i \sim \mathcal{N} \left( \sum_{i=1}^n a_i \mu_i, \sum_{i=1}^n a_i^2 \sigma_i^2 \right) \quad (1.33)$$

*Proof:* We once again prove this kind of theorems via the mgf for  $Y$ :

$$\begin{aligned} M(t) &= \prod_{i=1}^n \exp \left( t a_i \mu_i + \frac{1}{2} a_i^2 \sigma_i^2 \right) \\ &= \exp \left\{ t \sum_{i=1}^n a_i \mu_i + \frac{1}{2} t^2 \sum_{i=1}^n a_i^2 \sigma_i^2 \right\} \end{aligned} \quad (1.34)$$

which is the mgf for the normal with the corresponding mean and variance above.  $\square$

**Corollary:** Let  $X_1, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$ . Then

$$\bar{X} = \frac{\sum_{i=1}^n X_i}{n} \sim \mathcal{N}(\mu, \sigma^2/n) \quad (1.35)$$

*Proof:* the proof is left to the reader.

### 1.6.1 Contaminated Normal

We won't worry about this for now.

## 1.7 The Multivariate Normal

I'll just jump straight to the  $n$ -dimensional generalization. Evaluations of high-dimensional Gaussian integrals and moments can also be found in the [QFT](#) notes.

We say an  $n$ -dimensional random vector  $\mathbf{X}$  has a multivariate normal distribution if its mgf is

$$M_{\mathbf{X}}(t) = \exp \left( \mathbf{t}^\top \boldsymbol{\mu} + \frac{1}{2} \mathbf{t}^\top \boldsymbol{\Sigma} \mathbf{t} \right) \quad (1.36)$$

for all  $\mathbf{t} \in \mathbb{R}^n$ , where  $\boldsymbol{\Sigma}$  is a symmetric, positive semi-definite matrix and  $\boldsymbol{\mu} \in \mathbb{R}^n$ . For short, we say  $\mathbf{X} \sim \mathcal{N}_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ .

**Theorem 1.7.1.** Suppose  $\mathbf{X} \sim \mathcal{N}_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  where  $\boldsymbol{\Sigma}$  is positive definite. Then

$$\mathbf{Y} = (\mathbf{X} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{X} - \boldsymbol{\mu}) \sim \chi^2(1) \quad (1.37)$$

**Theorem 1.7.2.** If  $\mathbf{X} \sim \mathcal{N}_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  then

$$\mathbf{Y} = \mathbf{AX} + \mathbf{b} \sim \mathcal{N}_n(\mathbf{A}\boldsymbol{\mu} + \mathbf{b}) \quad (1.38)$$

*Proof:* The proof once again uses the mgf for  $\mathbf{Y}$ , but also some linear algebra manipulations.  $\square$

There are many other theorems and results related to this topic, but I won't go into them for now.

## 1.8 The $t$ - and $F$ -distributions

These two distributions are useful in certain problems in statistical inference.

### 1.8.1 The $t$ -distribution

Suppose  $W \sim \mathcal{N}(0, 1)$  and  $V \sim \chi^2(r)$  and that they are independent. Then the joint pdf of  $W$  and  $V$ , called  $h(w, v)$ , is the product of the pdf's of  $W$  and  $V$ :

$$h(w, v) = \begin{cases} \frac{1}{\sqrt{2\pi}} e^{-w^2/2} \frac{1}{\Gamma(r/2)2^{r/2}} v^{r/2-1} e^{-v/2}, & w \in \mathbb{R}, v > 0 \\ 0, & \text{else} \end{cases} \quad (1.39)$$

Now we define a new variable  $T = W/\sqrt{V/r}$  and consider the transformation:

$$t = \frac{w}{\sqrt{v/r}} \quad u = v \quad (1.40)$$

which bijectively maps the parameter space  $(w, v) = \mathbb{R} \times \mathbb{R}^+$  to  $(t, u) = \mathbb{R} \times \mathbb{R}^+$ . The absolute value of the Jacobian of the transformation is given by

$$|J| = \left| \det \begin{pmatrix} \partial_t w & \partial_u w \\ \partial_t v & \partial_u v \end{pmatrix} \right| = \frac{\sqrt{u}}{\sqrt{r}}. \quad (1.41)$$

With this, the joint pdf of  $T$  and  $U \equiv V$  is given by

$$g(t, u) = |J| h\left(\frac{t\sqrt{u}}{\sqrt{r}}, u\right) = \begin{cases} \frac{u^{r/2-1}}{\sqrt{2\pi}\Gamma(r/2)2^{r/2}} \exp\left[-\frac{u}{2}\left(1+\frac{t^2}{r}\right)\right] \frac{\sqrt{u}}{\sqrt{r}}, & t \in \mathbb{R}, u \in \mathbb{R}^+ \\ 0, & \text{else} \end{cases} \quad (1.42)$$

By integrating out  $u$  we obtain the marginal pdf for  $T$ :

$$\begin{aligned} g_1(t) &= \int_{-\infty}^{\infty} g(t, u) du \\ &= \int_0^{\infty} \frac{u^{(r+1)/2-1}}{\sqrt{2\pi r}\Gamma(r/2)2^{r/2}} \exp\left[-\frac{u}{2}\left(1+\frac{t^2}{r}\right)\right] du. \end{aligned} \quad (1.43)$$

Via the substitution  $z = u[1 + (t^2/r)]/2$  we can evaluate the integral to find for  $t \in \mathbb{R}$

$$g_1(t) = \boxed{\frac{\Gamma[(r+1)/2]}{\sqrt{\pi r}\Gamma(r/2)} \frac{1}{(1+t^2/r)^{(r+1)/2}}} \quad (1.44)$$

A r.v.  $T$  with this pdf is said to follow the  $t$ -distribution (or the Student's  $t$ -distribution) with  $r$  degrees of freedom. The  $t$ -distribution is symmetric about 0 and has a unique maximum at 0. As  $r \rightarrow \infty$ , the  $t$ -distribution converges to  $\mathcal{N}(0, 1)$ .

The mean of  $T \sim \text{Stu}(r)$  is zero. The variance can be found to be  $\text{Var}(T) = E[T^2] = \frac{r}{r-2}$ , so long as  $r > 2$ .

### 1.8.2 The F-distribution

Let  $U \sim \chi^2(r_1), V \sim \chi^2(r_2)$  be given. Then the joint pdf of  $U$  and  $V$  is once again the product of their pdf's:

$$h(u, v) = \begin{cases} \frac{1}{\Gamma(r_1/2)\Gamma(r_2/2)2^{(r_1+r_2)/2}} u^{r_1/2-1} v^{r_2/2-1} e^{-(u+v)/2}, & u, v \in \mathbb{R}^+ \\ 0, & \text{else} \end{cases} \quad (1.45)$$

Define the new random variable

$$W = \frac{U/r_1}{V/r_2} \quad (1.46)$$

whose pdf  $g_1(w)$  we are interested in finding. Consider the transformation

$$w = \frac{u/r_1}{v/r_2}, \quad z = v \quad (1.47)$$

which bijectively maps  $(u, v) = \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow (w, z) = [\mathbb{R}^+ \times \mathbb{R}^+]$ . Like last time, the absolute value of the Jacobian can be found to be

$$|J| = \frac{r_1}{r_2} z. \quad (1.48)$$

The joint pdf  $g(w, z)$  of the random variables  $W$  and  $Z = V$  is obtained from by scaling  $h(u, v)$  by  $|J|$  and applying the variable transformation:

$$g(w, z) = \frac{1}{\Gamma(r_1/2)\Gamma(r_2/2)2^{\frac{r_1+r_2}{2}}} \left( \frac{r_1 zw}{r_2} \right)^{\frac{r_1-2}{2}} z^{\frac{r_2-2}{2}} \exp \left[ -\frac{z}{2} \left( \frac{r_1 w}{r_2} + 1 \right) \right] \frac{r_1 z}{r_2} \quad (1.49)$$

so long as  $(w, z) \in \mathbb{R}^+ \times \mathbb{R}^+$  and 0 otherwise. We then proceed to find the marginal pdf  $g_1(w)$  of  $W$  by integrating out  $z$ . By considering the change of variables:

$$y = \frac{z}{2} \left( \frac{r_1 w}{r_2} + 1 \right) \quad (1.50)$$

we can evaluate the integral and find the marginal pdf of  $W$  to be

$$g_1(w) = \begin{cases} \frac{\Gamma[(r_1+r_2)/2](r_1/r_2)^{r_1/2}}{\Gamma(r_1/2)\Gamma(r_2/2)} \frac{w^{r_1/2-1}}{(1+r_1 w/r_2)^{(r_1+r_2)/2}}, & w \in \mathbb{R}^+ \\ 0, & \text{else} \end{cases} \quad (1.51)$$

$W$ , which is the ratio of two independent chi-square variables  $U, V$ , is said to follow an  $F$ -distribution with degrees of freedom  $r_1$  and  $r_2$ . We call the ratio  $W = (U/r_1)/(V/r_2)$  the “ $F$ ” ratio.

The mean of  $W$  is  $E[F] = \frac{r_2}{r_2-2}$ . When  $r_2$  is large,  $E[F] \rightarrow 1$ .

### 1.8.3 The Student's Theorem

Here we will create the connection between the normal distribution and the  $t$ -distribution. This is an important result for the later topics on inference for normal random variables.

**Theorem 1.8.1.** Let  $X_1, \dots, X_n$  be iid r.v. with  $X_i \sim \mathcal{N}(\mu, \sigma^2) \forall i$ . Define the r.v.’s

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \quad (1.52)$$

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2. \quad (1.53)$$

Then

- (a)  $\bar{X} \sim \mathcal{N}(\mu, \sigma^2/n)$ .
- (b)  $\bar{X}$  and  $S^2$  are independent.
- (c)  $(n-1)S^2/\sigma^2 \sim \chi^2(n-1)$ .
- (d) The variable  $\bar{T} = (\bar{X} - \mu)/(S/\sqrt{n})$  follows the Student's  $t$ -distribution with  $n-1$  degrees of freedom.

$$\bar{T} = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim \text{Stu}(n-1). \quad (1.54)$$

*Proof:* The proof is good, so I will reproduce it here. Because  $X_i \sim \mathcal{N}(\mu, \sigma^2) \forall i$ ,  $\mathbf{X} \sim \mathcal{N}_n(\mu\mathbf{1}, \sigma^2\mathbf{I})$ , where  $\mathbf{1}$  denotes the  $n$ -vector whose components are all 1. Now, consider  $\mathbf{v}^\top = (1/n)\mathbf{1}^\top$ . We see that  $\bar{X} = \mathbf{v}^\top \mathbf{X}$ . Define the random vector  $\mathbf{Y} = (X_1 - \bar{X}, \dots, X_n - \bar{X})^\top$  and consider the (true) equality:

$$\mathbf{W} = \begin{pmatrix} \bar{X} \\ \mathbf{Y} \end{pmatrix} = \underbrace{\begin{pmatrix} \mathbf{v}^\top \\ \mathbb{I} - \mathbf{1}\mathbf{v}^\top \end{pmatrix}}_{\text{the transformation}} \mathbf{X} \quad (1.55)$$

which just restates our definitions nicely. We see that  $\mathbf{W}$  is a result of a linear transformation of multivariate normal random vector, and so it follows that  $\mathbf{W} \sim \mathcal{N}_{n+1}$  with mean

$$E[\mathbf{W}] = \begin{pmatrix} \mathbf{v}^\top \\ \mathbb{I} - \mathbf{1}\mathbf{v}^\top \end{pmatrix} \mu\mathbf{1} = \begin{pmatrix} \mu \\ \mathbf{0}_n \end{pmatrix} \quad (1.56)$$

and the covariance matrix

$$\Sigma = \begin{pmatrix} \mathbf{v}^\top \\ \mathbb{I} - \mathbf{1}\mathbf{v}^\top \end{pmatrix} \sigma^2 \mathbb{I} \begin{pmatrix} \mathbf{v}^\top \\ \mathbb{I} - \mathbf{1}\mathbf{v}^\top \end{pmatrix}^\top = \sigma^2 \begin{pmatrix} \frac{1}{n} & \mathbf{0}_n^\top \\ \mathbf{0}_n & \mathbb{I} - \mathbf{1}\mathbf{v}^\top \end{pmatrix} \quad (1.57)$$

From here, part (a) is proven. Next, observe that  $\Sigma$  is diagonal, and so all covariances are zero. This means  $\bar{X}$  is independent of  $\mathbf{Y}$ . But because  $S^2 = (n-1)^{-1}\mathbf{Y}^\top \mathbf{Y}$ ,  $\bar{X}$  is independent of  $S^2$  as well. So, (b) is proven.

Now, consider the r.v.

$$V = \sum_{i=1}^n \left( \frac{X_i - \mu}{\sigma} \right)^2 \quad (1.58)$$

Each summand of  $V$  is a square of an  $\mathcal{N}(0, 1)$  r.v., and so each follows a  $\chi^2(1)$ . Because  $V$  is a sum of squares of  $n$  such  $\chi^2(1)$ 's,  $V \sim \chi^2(n)$ . Next, we can rewrite  $V$  as

$$V = \sum_{i=1}^n \left( \frac{(X_i - \bar{X}) + (\bar{X} - \mu)}{\sigma} \right)^2 = \frac{(n-1)S^2}{\sigma^2} + \left( \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \right)^2. \quad (1.59)$$

By (b), the summands in the last equation are independent. The second term is a square of a  $\mathcal{N}(0, 1)$ , so it follows a  $\chi^2(1)$ . Taking mgfs of both sides, we get

$$(1 - 2t)^{-n/2} = \underbrace{E[\exp\{t(n-1)S^2/\sigma^2\}]}_{M(c)}(1 - 2t)^{-1/2}. \quad (1.60)$$

Solving for the mgf of  $(n-1)S^2/\sigma^2$  we get part (c). Finally, writing  $T$  as

$$T = \frac{(\bar{X} - \mu)/(\sigma/\sqrt{n})}{\sqrt{(n-1)S^2/(\sigma^2(n-1))}} \quad (1.61)$$

and using (a)-(c) gives us (d). *Hint:* consider what distributions the numerator and denominator of  $T$  follow.  $\square$

## **Part 2**

# **Elementary Statistical Inferences**

## 2.1 Sampling & Statistics

In statistical inferences, our ignorance about the pdf/pmf of a random variable  $X$  can be classified in two ways:

- The pdf/pmf is unknown.
- The pdf/pmf is assumed/known but its parameter vector  $\theta$  is not.

We consider the second class of classification for now.

**Definition:** If the random variables  $X_1, X_2, \dots, X_n$  are iid, then these random variables constitute a **random sample** of size  $n$  from the common distribution.

**Definition:** Let  $X_1, \dots, X_n$  denote a sample on a random variable  $X$ . Let  $T = T(X_1, \dots, X_n)$  be a function of the sample. Then  $T$  is called a **statistic**.

### 2.1.1 Point estimators

**Definition:** (*Unbiasedness*) Let  $X_1, \dots, X_n$  denote a sample on a random variable  $X$  with pdf  $f(x; \theta)$ ,  $\theta \in \Omega$ . Let  $T = T(X_1, \dots, X_n)$  be a statistic. We say that  $T$  is an *unbiased estimator* of  $\theta$  if  $E[T] = \theta$ .

We now introduce the concept of the **maximum likelihood estimator (mle)**. The information in the sample and the parameter  $\theta$  are involved in the joint distribution of the random sample. We write this as

$$L(\theta) = L(\theta; x_1, \dots, x_n) = \prod_{i=1}^n f(x_i; \theta). \quad (2.1)$$

This is called the **likelihood function** of the random sample. A measure of the center of  $L(\theta)$  seems to be an appropriate estimate of  $\theta$ . We often use the value of  $\theta$  at which  $L(\theta)$  is maximized. If this value is unique, then it is called the **maximum likelihood estimator (mle)**, denoted as  $\hat{\theta}$ :

$$\hat{\theta} = \text{Argmax} L(\theta). \quad (2.2)$$

We often work with the log of the likelihood in practice, which is the function  $l(\theta) = \log(L(\theta))$ . The logarithm is a strictly increasing function, so its maximum is obtained exactly when the maximum of  $L(\theta)$  is obtained. In most models, the pdf and pmf are differentiable functions of  $\theta$ , in which cases  $\hat{\theta}$  solves the equation:

$$\partial_\theta l(\theta) = 0 \quad (2.3)$$

This is equivalent to saying  $\hat{\theta}$  maximizes  $l(\theta)$ . If  $\theta$  is a vector of parameters, this results in a system of equations to be solved simultaneously. These equations are called the **estimating equations**, (EE).

### 2.1.2 Histogram estimates of pmfs and pdfs

Let  $X_1, \dots, X_n$  be a random sample on a random variable  $X$  with cdf  $F(x)$ . A histogram of the sample is an estimate of the pmf or pdf depending on whether  $X$  is discrete or continuous. We make no assumptions on the form of the distribution of  $X$ . In particular, we don't assume the parametric form of the distribution, hence the histogram is often called the **nonparametric** estimator.

#### The distribution of $X$ is discrete

Assume  $X$  is a discrete r.v. with pmf  $p(x)$ . Consider a sample  $X_1, \dots, X_n$ . Suppose  $X \in \mathcal{D} = \{a_1, \dots, a_n\}$ , then intuitively the estimate of  $p(a_j)$  is the relative frequency of  $a_j$ . More formally, for  $j = 1, \dots, m$  we define the statistic

$$I_j(X_i) = \begin{cases} 1 & X_i = a_j \\ 0 & X_i \neq a_j \end{cases} \quad (2.4)$$

Then the estimate of  $p(a_j)$  is the average

$$\hat{p}(a_j) = \frac{1}{n} \sum_{i=1}^n I_j(X_i) \quad (2.5)$$

The estimators  $\{\hat{p}(a_1), \dots, \hat{p}(a_m)\}$  constitute the nonparametric estimate of the pmf  $p(x)$ . We note that  $I_j(X_i)$  has a Bernoulli distribution with probability  $p(a_j)$ , and so

$$E[\hat{p}(a_j)] = \frac{1}{n} \sum_{i=1}^n E[I_j(X_i)] = \frac{1}{n} \sum_{i=1}^n p(a_j) = p(a_j), \quad (2.6)$$

which means  $\hat{p}(a_j)$  is an *unbiased estimator* of  $p(a_j)$ .

Now, suppose that the space of  $X$  is infinite, i.e.,  $\mathcal{D} = \{a_1, \dots\}$  then in practice we select a value, say  $a_m$ , and make the groupings

$$\{a_1\}, \{a_2\}, \dots, \{a_m\}, \tilde{a}_{m+1} = \{a_{m+1}, \dots\} \quad (2.7)$$

Let  $\hat{p}(\tilde{a}_{m+1})$  be the proportion of the sample items that are greater than or equal to  $a_{m+1}$ . Then the estimates  $\{\hat{p}(a_1), \dots, \hat{p}(a_{m+1})\}$  form our estimate of  $p(x)$ . To merge groups, the rule of thumb is to select  $m$  so that the frequency of the category  $a_m$  exceeds twice the combined frequencies of the categories  $a_{m+1}, a_{m+2}, \dots$

A histogram is a *barplot* pf  $\hat{p}(a_j)$  versus  $a_j$ . When  $a_j$  contains no ordinal information (e.g. hair colors, etc) then such histograms consist of nonabutting bars and are called *bar charts*. When the space  $\mathcal{D}$  is ordinal, then the histograms is an abutting bar chart plotted in the natural order of the  $a_j$ 's.

### The distribution of $X$ is continuous

Assume  $X$  is a continuous r.v. with pdf  $f(x)$ . Consider a sample  $X_1, \dots, X_n$ . We first sketch an estimate for this pdf at a specified value of  $x$ . For a given  $h > 0$ , we consider the interval  $(x-h, x+h)$ . By MVT, we have for  $\xi, |x - \xi| < h$ :

$$P(|X - x| < h) = \int_{x-h}^{x+h} f(t) dt \approx 2hf(x). \quad (2.8)$$

The LHS is the proportion of the sample items that fall in the interval  $(x - h, x + h)$ . This suggests the use of the estimate of  $f(x)$  at a given  $x$ :

$$\hat{f}(x) = \frac{\#\{|X_i - x| < h\}}{2hn}. \quad (2.9)$$

More formally, the indicator statistic is, for  $i = 1, \dots, n$

$$I_i(x) = \begin{cases} 1 & x - h < X_i < x + h \\ - & \text{else} \end{cases}, \quad (2.10)$$

from which we obtain the nonparametric estimator of  $f(x)$ :

$$\hat{f}(x) = \frac{1}{2hn} \sum_{i=1}^n I_i(x). \quad (2.11)$$

Since the sample items are iid:

$$E[\hat{f}(x)] = \frac{1}{2hn} nf(\xi) 2h = f(\xi) \rightarrow f(x) \quad \text{as } h \rightarrow 0. \quad (2.12)$$

Therefore  $\hat{f}(x)$  is *approximately* (as opposed to *exact* in the discrete case) an unbiased estimator of  $f(x)$ .  $I_i$  is called the **rectangular kernel** with **bandwidth**  $2h$ .

Provided realized values  $x_1, \dots, x_n$  of the random sample of  $X$  with pdf  $f(x)$ , there are many ways to obtain a histogram estimate of  $f(x)$ . First, select an integer  $m$ , an  $h > 0$ , and a value  $a < \min(x_i)$ , so that the  $m$  intervals cover the range of the sample. These intervals form our classes. Let  $A_j = (a + (2j-3)h, a + (2j-1)h]$  for  $j = 1, \dots, m$ . Let  $\hat{f}_h(x)$  denote our histogram estimate. For  $a - h < x \leq (2m-1)h$ ,  $x$  is in one and only one  $A_j$ . Then for  $x \in A_j$ , we define

$$\hat{f}_h(x) = \frac{\#\{x_i \in A_j\}}{2hn} \geq 0. \quad (2.13)$$

We see that

$$\begin{aligned}
\int_{-\infty}^{\infty} \hat{f}_h(x) dx &= \int_{a-h}^{a+(2m-1)h} \hat{f}_h(x) dx \\
&= \sum_{j=1}^m \int_{A_j} \frac{\#\{x_i \in A_j\}}{2hn} dx \\
&= \frac{1}{2hn} \sum_{j=1}^m \#\{x_i \in A_j\} [h(2j-1 - 2j+3)] \\
&= \frac{2h}{2hn} n \\
&= 1.
\end{aligned} \tag{2.14}$$

So  $\hat{f}_h(x)$  satisfies the properties of a pdf.

## 2.2 Confidence Intervals

**Definition:** Let  $X_1, \dots, X_n$  be a sample on a r.v.  $X$  which has the pdf  $f(x; \theta), \theta \in \Omega$ . Let  $0 < \alpha < 1$  be specified. Let  $L = L(X_1, \dots, X_n)$  and  $U = U(X_1, \dots, X_n)$  be two statistics. We say that the interval  $(L, U)$  is a  $(1 - \alpha)100\%$  **confidence interval** for  $\theta$  if

$$1 - \alpha \equiv P_{\theta}[\theta \in (L, U)]. \tag{2.15}$$

That is, the probability that the interval includes  $\theta$  is  $1 - \alpha$ , which is called the **confidence coefficient** or the **confidence level** of the interval.

Under normality, the confidence interval for  $\mu$  is given by

$$\boxed{(\bar{x} - t_{\alpha/2, n-1}s/\sqrt{n}, \bar{x} + t_{\alpha/2, n-1}s/\sqrt{n})} \tag{2.16}$$

where  $t_{\alpha/2, n-1}$  is the upper  $\alpha/2$  critical points of a  $t$ -distribution with  $n - 1$  df. This CI is referred to as the  $(1 - \alpha)100\%$   **$t$ -interval** for  $\mu$ .  $s$  is referred to as the **standard error** of  $\bar{X}$ .

**The Central Limit Theorem:** Let  $X_1, \dots, X_n$  denote the observations of a random sample from a distribution that has mean  $\mu$  and finite variance  $\sigma^2$ . Then the distribution function of the r.v.  $W_n = (\bar{X} - \mu)/(\sigma/\sqrt{n})$  converges to  $\Phi$ , the distribution function of the  $\mathcal{N}(0, 1)$  distribution, as  $n \rightarrow \infty$ .

When the sample is large, the CI for  $\mu$  can be given by

$$\boxed{(\bar{x} - z_{\alpha/2}s/\sqrt{n}, \bar{x} + z_{\alpha/2}s/\sqrt{n})} \tag{2.17}$$

In general, for the same  $\alpha$ , the  $t$ -CI is larger (and hence more conservative) than the  $z$ -CI. When  $\sigma$  is known, we replace  $s$  by  $\sigma$ .

The larger sample CI for  $p$  is given by

$$\boxed{(\hat{p} - z_{\alpha/2} \sqrt{\hat{p}(1-\hat{p})/n}, \hat{p} + z_{\alpha/2} \sqrt{\hat{p}(1-\hat{p})/n})} \quad (2.18)$$

where  $\sqrt{\hat{p}(1-\hat{p})/n}$  is called the standard error of  $\hat{p}$ .

### 2.2.1 CI for difference in means

By independence of samples,

$$\text{Var}(\hat{\Delta}) = \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2} \quad (2.19)$$

where  $\hat{\Delta} = \bar{X} - \bar{Y}$ . We can readily show that  $\hat{\Delta}$  is an unbiased estimator of  $\Delta = \mu_1 - \mu_2$ . Let the sample variances

$$S_j^2 = \frac{1}{n_j - 1} \sum_{i=1}^{n_j} (X_i - \bar{X})^2 \quad (2.20)$$

be given. Then the random variable follows the  $\mathcal{N}(0, 1)$ :

$$\boxed{Z = \frac{\hat{\Delta} - \Delta}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}} \sim \mathcal{N}(0, 1)} \quad (2.21)$$

The approximate  $(1 - \alpha)100\%$  CI for  $\Delta = \mu_1 - \mu_2$  is then given by

$$\left( (\bar{x} - \bar{y}) - z_{\alpha/2} \sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}, (\bar{x} - \bar{y}) + z_{\alpha/2} \sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}} \right) \quad (2.22)$$

This is a large sample  $(1 - \alpha)100\%$  CI for  $\mu_1 - \mu_2$ .

Now, suppose  $X \sim \mathcal{N}(\mu_1, \sigma^2)$  and  $Y \sim \mathcal{N}(\mu_2, \sigma^2)$  (i.e.  $X$  and  $Y$  are normally distributed with the same variance) are independent. We want to show  $\Delta \sim t$ -distribution. We know that  $\bar{X} \sim \mathcal{N}(\mu_1, \sigma^2/n_1)$  and  $\bar{Y} \sim \mathcal{N}(\mu_2, \sigma^2/n_2)$ , so it is true that

$$\frac{(\bar{X} - \bar{Y}) - (\mu_1 - \mu_2)}{\sigma \sqrt{1/n_1 + 1/n_2}} \sim \mathcal{N}(0, 1). \quad (2.23)$$

This quantity will later be the numerator of our  $T$ -statistic. Now, let

$$S_p^2 = \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2} \quad (2.24)$$

then  $S_p^2$ , the **pooled estimator** of  $\sigma^2$ , is also an unbiased estimator of  $\sigma^2$ . Because  $(n_i - 1)S_i^2/\sigma^2 \sim \chi^2(n - 1)$ , we have that  $(n - 2)S_p^2/\sigma^2 \sim \chi^2(n - 2)$ . And so

$$T = \frac{\frac{(\bar{X} - \bar{Y}) - (\mu_1 - \mu_2)}{\sigma\sqrt{1/n_1 + 1/n_2}}}{\sqrt{(n - 2)S_p^2/(n - 2)\sigma^2}} = \frac{(\bar{X} - \bar{Y}) - (\mu_1 - \mu_2)}{S_p\sqrt{1/n_1 + 1/n_2}} \sim t_{n-2} \quad (2.25)$$

From here, it is easy to work out the  $(1 - \alpha)100\%$  CI for  $\mu_1 - \mu_2$ :

$$\left( (\bar{x} - \bar{y}) - t_{\alpha/2, n-2} s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}, (\bar{x} - \bar{y}) + t_{\alpha/2, n-2} s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \right) \quad (2.26)$$

There is some difficulty when the unknown variances  $\sigma$  in the distributions of  $X$  and  $Y$  are not equal.

### 2.2.2 CI for difference in proportions

Our estimator of the difference in proportions  $p_1 - p_2$  is  $\bar{X} - \bar{Y} \equiv \hat{p}_1 - \hat{p}_2$  where  $X \sim b(1, p_1)$  and  $Y \sim b(1, p_2)$ . Of course, we know that  $\sigma_1^2 = p_1(1 - p_1)$  and  $\sigma_2^2 = p_2(1 - p_2)$ . From here, the approximate  $(1 - \alpha)100\%$  confidence interval for  $p_1 - p_2$  is

$$(\hat{p}_1 - \hat{p}_2) \pm z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} \quad (2.27)$$

## 2.3 Order Statistics

Let  $X_1, X_2, \dots, X_n$  denote a random sample from a distribution of the continuous type having a pdf  $f(x)$  that has support  $S = (a, b)$ , where  $\infty \leq a < b \leq \infty$ . Let  $Y_1 < Y_2 < \dots < Y_n$  represent  $X_1, X_2, \dots, X_n$  when the latter are arranged in ascending order of magnitude. We call  $Y_i, i = 1, 2, \dots, n$ , the  $i$ th order statistic of the random sample  $X_1, X_2, \dots, X_n$ . We have a theorem which gives the joint pdf of  $Y_1, Y_2, \dots, Y_n$ .

**Theorem 2.3.1.** The joint pdf of  $Y_1, Y_2, \dots, Y_n$  is given by

$$g(y_1, y_2, \dots, y_n) = \begin{cases} n!f(y_1)\dots f(y_n) & a < y_1 < \dots < y_n < b \\ 0 & \text{else} \end{cases} . \quad (2.28)$$

*Proof:* The support of  $X_1, \dots, X_n$  can be partitioned into  $n!$  mutually disjoint sets that map onto the support of the  $Y_i$ 's. Obviously the Jacobian for

each transformation is either 1 or  $-1$ .

$$\begin{aligned} g(y_1, y_2, \dots, y_n) &= \sum_{i=1}^{n!} |J_i| f(y_1) \dots f(y_n) \\ &= \begin{cases} n! f(y_1) \dots f(y_n) & a < y_1 < \dots < y_n < b \\ 0 & \text{else} \end{cases} \end{aligned} \quad (2.29)$$

### 2.3.1 Quantiles

Let  $X$  be a random variable with a continuous cdf  $F(x)$ . For  $0 < p < 1$ , define the  $p$ th quantile of  $X$  to be  $\xi_p = F^{-1}(p)$ . Let  $X_1, X_2, \dots, X_n$  be a random sample from the distribution of  $X$  and let  $Y_1 < Y_2 < \dots < Y_n$  be the corresponding order statistics. Let  $k$  be the greatest integer less than or equal to  $p(n+1)$ . We next define an estimator of  $\xi_p$  after making the following observation. The area under the pdf  $f(x)$  to the left of  $Y_k$  is  $F(Y_k)$ . The expected value of this area is

$$E[F(Y_k)] = \int_a^b F(y_k) g_k(y_k) dy_k \quad (2.30)$$

where  $g_k(y_k)$  is the pdf of  $Y_k$ . Consider the transformation  $z = F(y_k)$ , then the integral becomes

$$E[F(Y_k)] = \int_0^1 \frac{n!}{(k-1)!(n-k)!} z^k (1-z)^{n-k} dz = \dots = \frac{k}{n+1} \quad (2.31)$$

where we recognize the similarity between the integral and the integral of a beta pdf. So, on the average, there is  $k/(n+1)$  of the total area to the left of  $Y_k$ . Because  $p = k/(n+1)$ , it seems reasonable to take  $Y_k$  as an estimator of the quantile  $\xi_p$ . Hence, we call  $Y_k$  the  **$p$ th sample quantile**. It is also called the  **$100p$ th percentile** of the sample.

A **five-number** summary of the data consists of the following five sample quantiles: the minimum ( $Y_1$ ), the first quartile ( $Y_{.25(n+1)}$ ), the median, the third quartile ( $Y_{.75(n+1)}$ ), and the maximum ( $Y_n$ ). For this section, we use the notation  $Q_1$ ,  $Q_2$ , and  $Q_3$  to denote, respectively, the first quartile, median, and third quartile of the sample.

The five-number summary is the basis for a useful and quick plot of the data. This is called a **boxplot** of the data. In the **box and whisker** plots, we also define a potential outlier. Let  $h = 1.5(Q_3 - Q_1)$ . The **lower/upper fence** is defined by  $L/UF = Q_{1/3} \mp h$ . Points lying outside the  $(LF, UF)$  interval are called **potential outliers**.

### 2.3.2 CI for quantiles

Let  $X$  be a continuous random variable with cdf  $F(x)$ . For  $0 < p < 1$ , define the  $100p$ th distribution percentile to be  $\xi_p$ , where  $F(\xi_p) = p$ . For a sample of

size  $n$  on  $X$ , let  $Y_1 < Y_2 < \dots < Y_n$  be the order statistics. Let  $k = [(n+1)p]$ . Then the 100 $p$ th sample percentile  $Y_k$  is a point estimate of  $\xi_p$ .

Let  $i < [(n+1)p] < j$ , and consider the order statistics  $Y_i < Y_j$  and the event  $Y_i < \xi_p < Y_j$ . The event  $Y_i < \xi_p < Y_j$  is equivalent to obtaining between  $i$  (inclusive) and  $j$  (exclusive) successes in  $n$  independent trials. So,

$$P(Y_i < \xi_p < Y_j) = \sum_{w=i}^{j-1} \binom{n}{w} p^w (1-p)^{n-w}. \quad (2.32)$$

For the median, we denote  $\xi_{1/2}$  the median of  $F(x)$ , i.e.  $\xi_{1/2}$  solves  $F(x) = 1/2$ . Let  $Q_2$  denote the sample median, which is a point estimator of  $\xi_{1/2}$ . Take  $c_{\alpha/2}$  such that  $P[S \leq c_{\alpha/2}] = \alpha/2$  where  $S \sim b(n, 1/2)$ . Then note also that  $P[S \leq c_{\alpha/2}] = \alpha/2$ . From here we have

$$P[Y_{c_{\alpha/2}} < \xi_{1/2} < Y_{n-c_{\alpha/2}}] = 1 - \alpha. \quad (2.33)$$

So, if  $y_{\alpha/2+1}$  and  $y_{n-\alpha/2}$  are the realized values of the order statistics  $Y_{c_{\alpha/2}+1}$  and  $Y_{n-c_{\alpha/2}}$  then the interval

$$(y_{c_{\alpha/2}}, y_{n-c_{\alpha/2}}) \quad (2.34)$$

is a  $(1 - \alpha)100\%$  confidence interval for  $\xi_{1/2}$ .

## 2.4 Introduction to Hypothesis Testing

Suppose a r.v.  $X \sim f(x; \theta)$ , where  $\theta \in \Omega$ . Suppose that  $\theta \in \omega_0$  or  $\theta \in \omega_1$  where  $\omega_0$  and  $\omega_1$  are disjoint subsets of  $\Omega$  and  $\omega_0 \cup \omega_1 = \Omega$ . We label these hypotheses as

$$\begin{aligned} H_0 : \theta &\in \omega_0 \\ H_1 : \theta &\in \omega_1. \end{aligned} \quad (2.35)$$

$H_0$  is called the **null hypothesis**.  $H_1$  is called the **alternative hypothesis**. Type I error occurs when we decide that  $\theta \in \omega_1$  when in fact  $\theta \in \omega_0$ . Type II error occurs when we decide the opposite.

We require the **critical region**,  $C$ , to complete the testing structure for the general problem. Consider the r.v.  $X$  and the hypotheses given above.  $C$  is such that

$$\begin{aligned} \text{Reject } H_0 \text{ if } (X_1, \dots, X_n) &\in C \\ \text{Reject } H_1 \text{ if } (X_1, \dots, X_n) &\in C^c. \end{aligned} \quad (2.36)$$

**Type I** error occurs if  $H_0$  is rejected when it is true. **Type II** error occurs if  $H_0$  is retained when  $H_1$  is true.

**Definition:** We say a critical region  $C$  is of **size**  $\alpha$  if

$$\boxed{\alpha = \max_{\theta \in \omega_0} P_\theta[(X_1, \dots, X_n)] \in C} \quad (2.37)$$

Over all critical regions of size  $\alpha$ , we want to consider critical regions that have lower probabilities of Type II error, i.e., for  $\theta \in \omega_1$ , we want to maximize

$$1 - P_\theta[\text{Type II Error}] = P_\theta[(X_1, \dots, X_n) \in C] \quad (2.38)$$

The probability on the right side of the equation above is called the **power** of the test at  $\theta$ . It is the probability that the test detects the alternative when  $\theta \in \omega_1$  is the true parameter. Minimizing Type II error requires maximizing the test power. The **power function** of  $C$  is

$$\gamma_C(\theta) = P_\theta[(X_1, \dots, X_n) \in C]; \quad \theta \in \omega_1. \quad (2.39)$$

Given two critical regions  $C_1, C_2$  both of size  $\alpha$ .  $C_1$  is better than  $C_2$  if  $\gamma_{C_1}(\theta) \geq \gamma_{C_2}(\theta)$  for all  $\theta \in \omega_1$ .

A **simple** hypothesis completely specifies the underlying distribution. A **composite** hypothesis can be composed of many simple hypotheses and hence do not completely specify the distribution.  $\alpha$  is often referred to as the **significance level** of the test associated with that critical region.

## 2.5 Additional comments about statistical test

### 2.5.1 Observed Significance Level, $p$ -value

Suppose  $H_0 : \mu = \mu_0$  and  $H_1 : \mu > \mu_0$ , where  $\mu_0$  is maximized. Then we reject  $H_0$  in favor of  $H_1$  if  $\bar{X} \geq k$  where  $\bar{X}$  is the sample mean. The  $p$ -value is the probability that under  $H_0$ ,  $\bar{X} \geq \bar{x}$ :

$$\boxed{p\text{-value} = P_{H_0}(\bar{X} \geq \bar{x})} \quad (2.40)$$

If  $\alpha > p$  then we reject  $H_0$  in favor of  $H_1$ . Else, we fail to reject  $H_1$ .

## 2.6 Chi-Square Tests

Let r.v.  $X_i \sim \mathcal{N}(\mu_i, \sigma_i^2)$  for  $i = 1, \dots, n$ . Let  $X_1, \dots, X_n$  be mutually independent. The joint pdf is then

$$\frac{1}{\sigma_1 \dots \sigma_n (\sqrt{2\pi})^n} \exp \left[ -\frac{1}{2} \sum_{i=1}^n \left( \frac{x_i - \mu_i}{\sigma_i} \right)^2 \right] \quad (2.41)$$

The r.v.

$$\sum_{i=1}^n \left( \frac{x_i - \mu_i}{\sigma_i} \right)^2 \quad (2.42)$$

has a  $\chi^2(n)$  distribution.

We will now consider some r.v.s that have approximate  $\chi^2$  distribution. Suppose  $X_1 \sim b(n, p_1)$ . Consider the r.v. defined by

$$Y = \frac{X_1 - p_1}{\sqrt{np_1(1-p_1)}} \quad (2.43)$$

which has, as  $n \rightarrow \infty$ , an approximate  $\mathcal{N}(0, 1)$ . We know from earlier discussions that  $Y^2 \sim \mathcal{N}(0, 1)$ . Now, let  $X_2 = n - X_1$  and  $p_2 = 1 - p_1$ . Let  $Q_1 = Y^2$ . Then we have

$$Q_1 = \frac{(X_1 - np_1)^2}{np_1(1-p_1)} = \frac{(X_1 - np_1)^2}{np_1} + \frac{(X_1 - np_1)^2}{n(1-p_1)} = \frac{(X_1 - np_1)^2}{np_1} + \frac{(X_2 - np_2)^2}{np_2}. \quad (2.44)$$

In general, let  $X_1, \dots, X_{k-1}$  have a multinomial distribution with the parameters  $n$  and  $p_1, \dots, p_{k-1}$ . Let  $X_k = n - (X_1 + \dots + X_{k-1})$  and let  $p_k = 1 - (p_1 + \dots + p_{k-1})$ . Define  $Q_{k-1}$  by

$$Q_{k-1} = \sum_{i=1}^k \frac{(X_i - np_i)^2}{np_i} \quad (2.45)$$

As  $n \rightarrow \infty$ ,  $Q_{k-1} \sim \chi^2(k-1)$ . This makes the r.v.  $Q_{k-1}$  a basis of the tests of certain statistical hypotheses. For instance, when the joint pdf of  $X_1, X_2, \dots, X_{k-1}$  (and  $X_k = n - X_1 - \dots - X_{k-1}$ ) is a multinomial pmf with parameters  $n$  and  $p_1, \dots, p_{k-1}$  (and  $p_k = 1 - p_1 - \dots - p_{k-1}$ ), we can consider the simple null hypothesis  $H_0 : p_1 = p_{10}, \dots, p_{k-1} = p_{(k-1)0}$  where  $p_{10}, \dots, p_{(k-1)0}$  are specified numbers. Under this null, the r.v.

$$Q_{k-1} = \sum_1^k \frac{(X_i - np_{i0})^2}{np_{i0}} \quad (2.46)$$

has an approximate  $\chi^2(k-1)$ . Intuitively, when  $H_0$  is true,  $np_{i0}$  must be the expected value of  $X_i$ , which means  $Q_{k-1}$  is not too large. Thus, we reject  $H_0$  if  $Q_{k-1} \geq c$ . The critical value  $c$  is specified by the significance level  $\alpha$ ,  $\mathbf{c} = \text{qchisq}(1-\alpha, k-1)$ . This is frequently called the **goodness-of-fit** test.

We can have a chi-square test for **homogeneity**. Consider two multinomial distributions with parameters  $n_j, p_{1j}, \dots, p_{kj}$  and  $j = 1, 2$ . Let  $X_{ij}$  where  $i =$

$1, \dots, k$  and  $j = 1, 2$  be frequencies. Suppose  $n_1, n_2$  large and the observations are independent, then the r.v.

$$\sum_{j=1}^2 \sum_{i=1}^n \frac{(X_{ij} - n_j p_{ij})^2}{n_j p_{ij}} \sim \chi^2(2k - 2) \quad (2.47)$$

because it is the sum of two independent r.v.'s each of which  $\sim \chi^2(k - 1)$ . The null hypothesis we consider is

$$H_0 : p_{11} = p_{12}; \dots; p_{k1} = p_{k2}, \quad (2.48)$$

where each  $p_{i1} = p_{i2}$  where  $i = 1, \dots, k$  is unspecified. It turns out that the mle of  $p_{i1} = p_{i2}$  is given by

$$\theta = \frac{X_{i1} + X_{i2}}{n_1 + n_2} \quad (2.49)$$

which makes intuitive sense. Note that we need only  $k - 1$  points estimates, and so the r.v.

$$Q_{k-1} = \sum_{j=1}^2 \sum_{i=1}^k \frac{\{X_{ij} - n_j[(X_{i1} + X_{i2})/(n_1 + n_2)]\}^2}{n_j[(X_{i1} + X_{i2})/(n_1 + n_2)]} \sim \chi^2(2k - 2 - (k - 1) = k - 1). \quad (2.50)$$

With this, we can test if two multinomial distributions are the same.

We can also test for **independence**. Suppose the result of an experiment is classified by only two attributes  $A$  (of  $a$  possible outcomes) and  $B$  (of  $b$  possible outcomes). These events are  $A_1, \dots, A_a$  for attribute  $A$  and  $B_1, \dots, B_b$  for attribute  $B$ . Then consider  $p_{ij} = P(A_i \cap B_j)$ . Say the experiment is repeated  $n$  independent times and  $X_{ij}$  denotes the frequency of the event  $A_i \cap B_j$ . There are  $k = ab$  such events, so the r.v.

$$Q_{ab-1} = \sum_{j=1}^b \sum_{i=1}^a \frac{(X_{ij} - np_{ij})^2}{np_{ij}} \sim \chi^2(ab - 1), \quad (2.51)$$

provided  $n$  is large. To test for independence,  $H_0 : P(A_i \cap B_j) = P(A_i)P(B_j)$  for all  $i, j$ . To test  $H_0$ , we cannot compute  $Q_{ab-1}$ , but instead compute

$$\sum_{j=1}^b \sum_{i=1}^a \frac{[X_{ij} - n(X_{i.}/n)(X_{.j}/n)]^2}{n(X_{i.}/n)(X_{.j}/n)} \sim \chi^2(ab - 1 - (a + b - 2) = (a - 1)(b - 1)) \quad (2.52)$$

where

$$\hat{p}_{i.} = \frac{X_{i.}}{n}, \quad X_{i.} = \sum_{j=1}^b X_{ij}, i = 1, \dots, a \quad (2.53)$$

$$\hat{p}_{.j} = \frac{X_{.j}}{n}, \quad X_{.j} = \sum_{i=1}^a X_{ij}, j = 1, \dots, b \quad (2.54)$$

Just a sanity check, the chi-square statistic always has the form of  $\sum \text{Expected} - \text{Observed}^2/\text{Expected}$ . All tests' statistics have this form. The differences are subtle and are context-based.

## 2.7 The Method of Monte Carlo

The idea of Monte Carlo methods is to use random numbers to simulate random phenomena and to make numerical approximations. In general, we use Monte Carlo methods for

- Inverse transform sampling: take a random uniform  $(0, 1)$  and transform it into a different distribution.
- Accept-Reject Algorithm, which is a method that uses a random uniform generator to produce a set of random numbers that follows some other distribution.
- To approximate the value of definite integrals.

### 2.7.1 Inverse Transform

For example, we want to simulate coin flips of a coin that is biased and comes up heads with probability  $p$ . Here's the algorithm:

- Generate a random uniform  $(0, 1) \rightarrow u_1$
- If  $u_1 < p \implies$  heads, else tails.
- Repeat

For any multinomial distribution with probabilities  $p_1, \dots, p_k$  we can follow this process:

- Generate a random uniform  $u - 1 \sim (0, 1)$ .
- If  $u_1 \leq p_1 \implies$  assign outcome 1.
- Elif  $u_1 \leq p_1 + p_2 \implies$  assign outcome 2.
- ...
- Elif  $u_1 \leq p_1 + \dots + p_{k-1} \implies$  assign outcome  $k - 1$ .
- Else, assign outcome  $k$ .

Basically, what we're doing here is using the CDF to check against  $u$ .

In the continuous case, we generate  $u \sim (0, 1)$  but we want a random variable with some other density  $X \sim f_X(x)$ . Assuming that  $X = T(U)$ , starting with the CDF of  $X$ :

$$\begin{aligned} F(x) &= P(X \leq x) \\ &= P(T(u) \leq x) \\ &= P(u \leq T^{-1}(x)) \\ &= F_u(T^{-1}(x)) \\ &= T^{-1}(x) \end{aligned} \tag{2.55}$$

where the last equality follows from the fact that  $F_u(u)$  is just the identity function. So we have

$$F(x) = T^{-1}(x) \tag{2.56}$$

which means  $F$  and  $T$  are inverses of each other.

For example, we can use the inverse transform to generate random  $\text{Exp}(\beta)$ :

$$f_X(x) = \frac{1}{\beta} e^{-x/\beta}, \quad x \in \mathbb{R}^+ \tag{2.57}$$

We want to find  $F_X(x)$  first:

$$F_X(x) = \int_0^x \frac{1}{\beta} e^{-x'/\beta} dx' = 1 - e^{-x/\beta}. \tag{2.58}$$

And so it is easy to see that

$$T^{-1}(x) = 1 - e^{-x/\beta} \implies u = 1 - e^{-x/\beta} \implies x = \beta \log(1 - u). \tag{2.59}$$

From here, we can generate a sample of uniform  $u$ 's to get  $X \sim \text{Exp}(\beta)$ .

Note that a disadvantage to this method is the fact that for this to work we must be able to write down the inverse CDF in some closed form. The advantage, though, is that this method is very efficient if it works. This is because for each  $u$  we generate we get an  $x$ . This is not the case for the methods we will discuss next.

### 2.7.2 Accept-Reject Generation Algorithm

### 2.7.3 Evaluating definite integrals

This is based on the idea that

$$E[g(x)] = \int_{-\infty}^{\infty} g(x)f(x)dx. \tag{2.60}$$

For example,

$$\int_0^1 e^{-x^2/2} dx = E[e^{-x^2/2}] \quad (2.61)$$

where  $x \sim U(0, 1)$ .

We could estimate  $g(\cdot)$  by

$$\bar{g} = \frac{1}{J} \sum_{i=1}^J [g(x^{(j)})] \quad (2.62)$$

where  $x^{(j)} \sim f(x)$ . If we take enough random variables,  $\bar{g} \rightarrow E[g]$ . We could also use the CLT to calculate error bounds:

$$\bar{g} \pm z_{\alpha/2} SE(\bar{g}), \quad (2.63)$$

where

$$SE(\bar{g}) = \frac{1}{J^2} \sum_{i=1}^J \left( g(x^{(j)}) - \bar{g} \right)^2. \quad (2.64)$$

For example, we can try to evaluate

$$\int_0^\infty x^4 e^{-x^2/2} dx. \quad (2.65)$$

We identify  $g(x) = x^4$ , and  $e^{-x}$  as the density for  $\text{Exp}(1)$ . What we can do is generate a sample of random  $\text{Exp}(1)$ , lu into  $x^4$  then take the average value of  $x^4$ .

## 2.8 Bootstrapping

The basic idea of bootstrapping is this: We have a sample of data  $x_1, \dots, x_n = \vec{X}$ . We will replicate sample infinitely many times. This will be a model for the population.

Note that this process doesn't work well in the case that our sample is poorly representative. In practice, we obtain many bootstrap samples  $X^*$ 's where each  $X^*$  is a resample from our original sample where we randomly select  $X_i^*$  from  $\vec{X}$  **with replacement**.

The bootstrap sample is the same size as the original sample.

If we're interested in estimating a parameter using an estimator  $\hat{\theta} = f(\vec{X})$ , then we can calculate an estimate  $\hat{\theta}^* = f(\vec{X}^*)$  from each bootstrap sample. The bootstrap distribution of  $\hat{\theta}^*$  models the sampling distribution of  $\hat{\theta}$ .

For example, let  $x_1, \dots, x_n$  be a random sample from some population with an unknown mean  $\mu$ . We can think about taking a simple bootstrap sample,  $X_i^*$ , then calculate its mean. Of course, this is an unbiased estimator for  $\mu$ .

If we take many bootstrap samples  $X_i^*$  and calculate the mean of each, we could generate a bootstrap for  $\bar{X}^*$ . In an estimation setting, the most common use of bootstrap distribution is to estimate the SE of  $\hat{\theta}$ .

We can also find CI using percentiles/normal approximation this way.

### 2.8.1 Bootstrapping for Hypothesis Testing

Say we have  $H_0 : \mu = \mu_0$  and  $H_a : \mu > \mu_0$ . To test these hypotheses with bootstrapping, we shift the original data such that the shifted mean is  $\mu_0$ , i.e. we do  $X_i - \bar{X} + \mu_0$  to all observations.

From here, we bootstrap and generate a sampling distribution for the mean of the bootstrap samples. Then, we look and count how many bootstraps means are greater than the observed means and find the associated  $p$ -value.

We note that the advantage of doing this is we don't make any distributional assumptions. The disadvantage (kind of) is that this process can be a little too computationally expensive. However, with modern computers, this is no longer a major problem.

For example, say we want to compare two means.  $H_0 : \Delta = 0$ ,  $H_a : \Delta \neq 0$ . Say we have two samples (of different sample sizes) from cdfs  $F(x)$  and  $F(x-\Delta)$ ,

respectively. Under the null, these samples come from the same distribution. So, we can follow these steps:

- Combine the samples into a single sample.
- Take one bootstrap of size  $n_1$  and one of size  $n_2$ .
- Calculate the difference in means.
- Repeat many times
- Count how many bootstrap differences are further from 0 than the original observed difference in means.
- Extract the p-value (either by percentile or by SE).



## Part 3

# Consistency and Limiting Distributions

### 3.1 Convergence in Probability

**Definition:** Let  $\{X_n\}$  be a sequence of r.v. and let  $X$  be a r.v. defined on a sample pace.  $X_n$  converges in probability to  $X$  if, for all  $\epsilon < 0$ ,

$$\lim_{n \rightarrow \infty} P[|X_n - X| \geq \epsilon] = 0, \quad (3.1)$$

i.e.,

$$\lim_{n \rightarrow \infty} P[|X_n - x| < \epsilon] = 1. \quad (3.2)$$

If so, we write

$$X_n \xrightarrow{P} X. \quad (3.3)$$

**Theorem 3.1.1.** (handy theorem) If  $\hat{\theta}_n$  is an unbiased estimator of  $\theta$ , then  $\hat{\theta}_n \xrightarrow{P} \hat{\theta}$  if  $\lim_{n \rightarrow \infty} \text{Var}(\hat{\theta}_n) = 0$ . In which case, we call  $\hat{\theta}$  a consistent estimator of  $\theta$ .

**Theorem 3.1.2.** (Weak Law of Large Numbers). Let  $\{X_n\}$  be a sequence of iid r.v. having common mean  $\mu$  and variance  $\sigma^2$ , then

$$\bar{X} \equiv \frac{1}{n} \sum_{i=1}^n X_i \xrightarrow{P} \mu. \quad (3.4)$$

*Proof:* The proof uses Chebychev's inequality. Let  $\epsilon > 0$  be given, then

$$P[|X_n - \bar{X}| \geq \epsilon] = P[|\bar{X} - \mu| \geq (\epsilon\sqrt{n}/\sigma)(\sigma/\sqrt{n})] \leq \frac{\sigma^2}{n\epsilon^2} \rightarrow 0, \quad n \rightarrow \infty \quad (3.5)$$

□

**Theorem 3.1.3.** If  $X_n \xrightarrow{P} X$  and  $Y_n \xrightarrow{P} Y$ , then

$$X_n + Y_n \xrightarrow{P} X + Y. \quad (3.6)$$

*Proof:* The proof is quite easy. It uses the fact that  $P$  is monotone relative to set containment and the triangle inequality.  $\square$

**Theorem 3.1.4.** If  $X_n \xrightarrow{P} X$  then  $aX_n \xrightarrow{a} aX$ .

*Proof:* The proof is also very easy, so I won't show it here.  $\square$

**Theorem 3.1.5.** If  $X_n \xrightarrow{P} a$  and the real function  $g$  is continuous at  $a$  then

$$g(X_n) \xrightarrow{P} g(a). \quad (3.7)$$

*Proof:* The proof is analysis-like. It's not so hard so I (again) won't show it here.

**Theorem 3.1.6.** If  $X_n \xrightarrow{P} X$  and  $Y_n \xrightarrow{P} Y$ , then

$$X_n Y_n \xrightarrow{P} XY. \quad (3.8)$$

*Proof:* This proof uses the result from the previous theorem. The key is to write  $X_n Y_n$  as a combination of  $X_n^2$ ,  $Y_n^2$ , and  $(X_n - Y_n)^2$ . Applying the previous to obtain the desired conclusion.  $\square$

### 3.1.1 Sampling and Statistic

**Definition:** (Consistency) Let  $X$  be a r.v. with cdf  $F(x, \theta)$  with  $\theta \in \Omega$ . Let  $X_1, \dots, X_n$  be a sample from the distribution of  $X$  and let  $T_n$  denote a statistic.  $T_n$  is a **consistent** estimator of  $\theta$  iff

$$T_n \xrightarrow{P} \theta. \quad (3.9)$$

## 3.2 Convergence in Distribution

**Definition:** (Convergence in Distribution) Let  $\{X_n\}$  be a sequence of r.v.s and let  $X$  be a r.v.. Let  $F_{X_n}$  and  $F_X$  be, respectively, the cdfs of  $X_n$  and  $X$ . Let  $C(F_X)$  denote the set of all points where  $F_X$  is continuous. We say that  $X_n$  converges in distribution to  $X$  if

$$\lim_{n \rightarrow \infty} F_{X_n}(x) = F_X(x), \quad \forall x \in C(F_X). \quad (3.10)$$

We denote this convergence by  $X_n \xrightarrow{D} X$ .

**Stirling's Formula:**

$$\Gamma(k+1) \approx \sqrt{2\pi} k^{k+1/2} e^{-k} \quad (3.11)$$

when  $k$  is large .

**Theorem 3.2.1.** If  $X_n$  converges to  $X$  in probability, then  $X_n$  converges to  $X$  in distribution.

**Theorem 3.2.2.** If  $X_n$  converges to the constant  $b$  in distribution, then  $X_n$  converges to  $b$  in probability.

**Theorem 3.2.3.** Suppose  $X_n$  converges to  $X$  in distribution and  $Y_n$  converges in probability to 0, then  $X_n + Y_n$  converges to  $X$  in distribution.

**Theorem 3.2.4.** Suppose  $X_n$  converges to  $X$  in distribution and  $g$  is a continuous function on the support of  $X$ . Then  $g(X_n)$  converges to  $g(X)$  in distribution.

**Theorem 3.2.5.** (Slutsky's Theorem) Let  $X_n, X, A_n$ , and  $B_n$  be random variables and let  $a$  and  $b$  be constants. If  $X_n \xrightarrow{D} X$ ,  $A_n \xrightarrow{P} a$ , and  $B_n \xrightarrow{P} b$  then

$$A_n + B_n X_n \xrightarrow{D} a + bX. \quad (3.12)$$

### 3.2.1 Bounded in Probability

**Definition:** We say that the sequence of random variables  $\{X_n\}$  is bounded in probability if, for all  $\epsilon > 0$ , there exists a constant  $B_\epsilon > 0$  and an integer  $N_\epsilon$  such that

$$n \geq N_\epsilon \implies P[|X_n| \leq B_\epsilon] \geq 1 - \epsilon. \quad (3.13)$$

**Theorem 3.2.6.** Let  $\{X_n\}$  be a sequence of r.v. and let  $X$  be a r.v.. If  $X_n \rightarrow X$  in distribution, then  $\{X_n\}$  is bounded in probability.

**Theorem 3.2.7.** Let  $\{X_n\}$  be a sequence of r.v. bounded in probability and let  $\{Y_n\}$  be a sequence of r.v. that converges to 0 in probability. Then

$$X_n Y_n \xrightarrow{P} 0. \quad (3.14)$$

### 3.2.2 $\Delta$ -method

Little  $o$  notation:  $a = o(b)$  if and only if  $a/b \rightarrow 0$  as  $b \rightarrow 0$ .

**Theorem 3.2.8.** Suppose  $\{Y_n\}$  is a sequence of r.v. that is bounded in probability. Suppose  $X_n = o_P(Y_n)$ , then  $X_n \xrightarrow{P} 0$ , as  $n \rightarrow \infty$ .

**Theorem 3.2.9.** Let  $\{X_n\}$  be a sequence of r.v. such that

$$\sqrt{n}(X_n - \theta) \xrightarrow{D} \mathcal{N}(0, \sigma^2). \quad (3.15)$$

Suppose  $g(x)$  is differentiable at  $\theta$  and  $g'(\theta) \neq 0$ , then

$$\sqrt{n}(g(X_n) - g(\theta)) \xrightarrow{D} \mathcal{N}(0, \sigma^2(g'(\theta))^2) \quad (3.16)$$

### 3.2.3 Moment Generating Function Technique

**Theorem 3.2.10.** Let  $\{X_n\}$  be a sequence of r.v. with mgf  $M_{X_n}(t)$  that exists for  $-h < t < h$  for all  $n$ . Let  $X$  be a r.v. with mdf  $M(t)$ , which exists for  $|t| \leq h_1 \leq h$ . If  $\lim_{n \rightarrow \infty} M_{X_n}(t) = M(t)$  for  $|t| \leq h_1$ , then  $X_n \xrightarrow{D} X$ .

## 3.3 Central Limit Theorem

**Theorem 3.3.1.** (CLT) Let  $X_1, \dots, X_n$  denote the observations of a r.v. from a distribution that has mean  $\mu$  and positive variance  $\sigma^2$ . Then the r.v.

$$Y_n = \frac{\sum_{i=1}^n X_i - n\mu}{\sigma\sqrt{n}} = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow{D} Z \sim \mathcal{N}(0, 1). \quad (3.17)$$

*Proof:* Assume that the mgf  $M(t) = E[e^{tX}]$  exists for  $-h < t < h$ , then the function

$$m(t) = E[e^{t(X-\mu)}] = e^{-\mu t} M(t) \quad (3.18)$$

also exists for  $-h < t < h$ .  $m(t)$  is the mgf for  $X - \mu$ , so  $m(0) = 1, m'(0) = E[X - \mu] = 0$ , and  $m''(0) = E[(X - \mu)^2] = \sigma^2$ . By Taylor theorem, there exists a number  $\xi \in [0, t]$  such that

$$\begin{aligned} m(t) &= m(0) + m'(0)t + \frac{m''(\xi)t^2}{2} \\ &= 1 + \frac{m''(\xi)t^2}{2} \\ &= 1 + \frac{\sigma^2 t^2}{2} + \frac{[m''(\xi) - \sigma^2]t^2}{2}. \end{aligned} \quad (3.19)$$

Now consider  $M(t; n)$ :

$$\begin{aligned} M(t; n) &= E \left[ \exp \left( t \frac{\sum X_i - n\mu}{\sigma\sqrt{n}} \right) \right] \\ &= \dots \\ &= \left\{ E \left[ \exp \left( t \frac{X - \mu}{\sigma\sqrt{n}} \right) \right] \right\}^n \\ &= \left[ m \left( \frac{t}{\sigma\sqrt{n}} \right) \right]^n, \quad -h < \frac{t}{\sigma\sqrt{n}} < h \\ &= \left\{ 1 + \frac{t^2}{2n} + \frac{[m''(\xi) - \sigma^2]t^2}{2} \right\}^n. \end{aligned} \quad (3.20)$$

Taking  $n \rightarrow \infty$ :

$$\lim_{n \rightarrow \infty} [m''(\xi) - \sigma^2] = 0, \quad (3.21)$$

and so

$$\lim_{n \rightarrow \infty} M(t; n) = e^{t^2/2}, \quad t \in \mathbb{R}. \quad (3.22)$$

So  $Y_n \sim \mathcal{N}(0, 1)$ .



## **Part 4**

# **Maximum Likelihood Methods**

## 4.1 Maximum Likelihood Estimation

Recall the likelihood function:

$$\mathcal{L}(\theta; \mathbf{x}) = \prod_{i=1}^n f(x_i; \theta), \quad \theta \in \Omega \quad (4.1)$$

where  $f(x_i; \theta)$  is the pdf which the variables  $X_i$  follow that depends on the parameter  $\theta$  and  $\mathbf{x} = (x_1, x_2, \dots, x_n)^\top$  is the sample. It's often more convenient to use the log likelihood:

$$l(\theta) = \ln \mathcal{L}(\theta) = \sum_{i=1}^n \log f(x_i; \theta). \quad (4.2)$$

$\hat{\theta}$  is the mle of  $\theta$  if  $\hat{\theta}$  maximizes  $l(\theta)$ . Let  $\theta_0$  denote the true value of  $\theta$ . We will look at theorem which shows that the maximum of  $\mathcal{L}(\theta)$  asymptotically separates the true model at  $\theta_0$  from models at  $\theta \neq \theta_0$ . To prove this theorem, we look at *regularity conditions*:

**Regularity Conditions.** Regular conditions are

- The cdfs are distinct, i.e.,  $\theta \neq \theta' \implies F(x_i; \theta) \neq F(x_i; \theta')$ .
- The pdfs have common support for all  $\theta$ .
- The point  $\theta_0$  is an interior point in  $\Omega$ .

**Theorem 4.1.1.** Assume that  $\theta_0$  is the true parameter and that

$$E_{\theta_0}[f(X_i, \theta)/f(X_i, \theta_0)] \quad (4.3)$$

exists. Under the first two regularity conditions

$$\lim_{n \rightarrow \infty} P_{\theta_0}[\mathcal{L}(\theta_0, \mathbf{X}) > \mathcal{L}(\theta, \mathbf{X})] = 1, \quad \forall \theta \neq \theta_0. \quad (4.4)$$

**Definition 4.1.1.** (Maximum Likelihood Estimator). We say that  $\hat{\theta} = \hat{\theta}(\mathbf{X})$  is a maximum likelihood estimator (mle) of  $\theta$  if

$$\hat{\theta} = \text{Argmax} \mathcal{L}(\theta, \mathbf{X}) \quad (4.5)$$

where the notation means  $\mathcal{L}(\theta, \mathbf{X})$  attains maximum at  $\hat{\theta}$ .

**Theorem 4.1.2.** (Invariance Property) Let  $X_1, \dots, X_n$  be iid with pdf  $f(x; \theta), \theta \in \Omega$ . For a specified function  $g$ , let  $\eta = g(\theta)$  be a parameter of interested. Suppose  $\hat{\theta}$  is the mle of  $\theta$ . Then  $g(\hat{\theta})$  is the mle of  $\eta = g(\theta)$ .

**Theorem 4.1.3.** Assume that  $X_1, \dots, X_n$  satisfy the regularity conditions, where  $\theta_0$  is the true parameter, and further that  $f(x; \theta)$  is differentiable w.r.t.  $\theta \in \Omega$ . Then the likelihood equation,

$$\partial_\theta \mathcal{L}(\theta) = 0 \iff \partial_\theta l(\theta) = 0 \quad (4.6)$$

has a solution  $\hat{\theta}_n$  such that  $\hat{\theta}_n \xrightarrow{P} \theta_0$ .

**Theorem 4.1.4.** Assume that  $X_1, \dots, X_n$  satisfy the regularity conditions, where  $\theta_0$  is the true parameter, and that  $f(x; \theta)$  is differentiable w.r.t.  $\theta \in \Omega$ . Suppose the likelihood equation has the **unique** solution  $\hat{\theta}_n$ . Then  $\hat{\theta}_n$  is a consistent estimator of  $\theta_0$ .

So far, we know that two things that make mle good:

- Invariance property
- Consistency of MLEs.

**Example 4.1.1.** Let  $X_1, \dots, X_n$  be iid  $\text{Exp}(\theta)$  r.v.'s. We want to find  $\hat{\theta}_{\text{MLE}}$ . We know that

$$f(x_i; \theta) = \frac{1}{\theta} e^{-x_i/\theta}; \quad x_i > \theta, \theta > 0. \quad (4.7)$$

The likelihood function is

$$\mathcal{L}(\theta) = \frac{1}{\theta^n} e^{-\sum_{i=1}^n x_i/\theta}. \quad (4.8)$$

The log likelihood is

$$l(\theta) = -n \ln \theta - \frac{1}{\theta} \sum_{x_i}. \quad (4.9)$$

from which we can easily solve the mle

$$\hat{\theta} = \frac{1}{n} \sum_i x_i = \bar{x}. \quad (4.10)$$

Since all regular conditions are satisfied, this is a *good* mle.

**Example 4.1.2.** Let  $X_1, \dots, X_n$  be iid  $U(0, \theta)$  r.v.. We want to find the  $\hat{\theta}_{ML}$ . Now,

$$f(x_i; \theta) = \frac{1}{\theta}; 0 \leq x_i \leq \theta; \theta > 0. \quad (4.11)$$

We note that the second and third regularity conditions do not hold. Next,

$$\mathcal{L}(\theta) = \frac{1}{\theta^n} \implies l(\theta) = -n \ln \theta. \quad (4.12)$$

We want  $\hat{\theta}$  to be as small as possible to maximize  $\mathcal{L}(\theta)$ , but it also has to be bigger than all of the observations. Thus,  $\hat{\theta}_{ML} = \max_i(X_i)$ .

**Example 4.1.3.** Let  $X_i \sim \mathcal{N}(\mu, \sigma^2)$  r.v.. We want to find  $\hat{\mu}$  and  $\hat{\sigma}^2$ . We know that

$$f(x_i; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x_i-\mu)^2/2\sigma^2}; x_i, \mu \in \mathbb{R}, \sigma^2 > 0. \quad (4.13)$$

The log likelihood is easy:

$$l(\mu, \sigma^2) = -\frac{n}{2} \ln(2\pi) - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2\sigma^2} \sum (x_i - \mu)^2. \quad (4.14)$$

Then

$$\partial_\mu l(\mu, \sigma^2) = -\frac{1}{\sigma^2} \sum (x_i - \mu) = 0 \implies \hat{\mu} = \bar{x}. \quad (4.15)$$

Also,

$$\partial_{\sigma^2} = -\frac{n}{2} \frac{1}{\sigma^2} + \frac{1}{2\sigma^4} \sum (x_i - \bar{x})^2 = 0 \implies \hat{\sigma}^2 = \frac{1}{n} \sum (x_i - \bar{x})^2. \quad (4.16)$$

Because the regularity conditions are satisfied, these are consistent estimators.

**Example 4.1.4.** Let  $X_i \sim \text{Bernoulli}$  with parameter  $p$ , i.e.,  $P(x_i = 1) = p$  and  $P(x_i = 0) = 1 - p$ . Then

$$P(x_1, \dots, x_n) = p^{\sum x_i} (1-p)^{n-\sum x_i}. \quad (4.17)$$

Then the log likelihood function is just

$$l(p) = \sum x_i \ln p + \left(n - \sum x_i\right) \ln(1-p). \quad (4.18)$$

It follows that

$$\partial_p l(p) = \frac{\sum x_i}{p} - \frac{n - \sum x_i}{1-p} = 0 \implies \hat{p} = \frac{\sum x_i}{n}. \quad (4.19)$$

This is a consistent estimator because the regularity conditions are satisfied.

**Example 4.1.5.** Let  $X_i \sim \text{Poi}(\lambda)$ . We want to find  $\hat{\lambda}$ . We have

$$p(x_i | \lambda) = \frac{e^{-\lambda} \lambda^{x_i}}{x_i!}, \quad x_i = 1, 2, \dots, \quad \lambda > 0. \quad (4.20)$$

The log likelihood function is

$$l(\lambda) = -ns + \sum x_i \ln \lambda - \sum \ln(x_i!). \quad (4.21)$$

So,

$$\partial_\lambda l(\lambda) = 0 \implies \hat{\lambda} = \bar{x}. \quad (4.22)$$

From WLLN,  $\bar{x} \rightarrow \mu$ , so as long as the regularity conditions are satisfied (which they are), then we have a consistent estimator for  $\mu$ .

## 4.2 Rao-Cramér Lower Bound and Efficiency

**Additional Regularity Conditions.**

- The pdf of  $f(x; \theta)$  is twice differentiable as a function of  $\theta$ .
- The integral  $\int f(x; \theta) dx$  can be differentiated twice under the integral sign as a function of  $\theta$ .

All four regularity conditions we have seen so far combined means that the parameter  $\theta$  does not appear in the endpoints of the interval in which  $f(x; \theta) > 0$  and that we can interchange integration and differentiation w.r.t  $\theta$ . The derivation is below is the continuous case, the discrete case can be handled in a similar manner. I'll summarize the derivation in a few steps below:

$$1 = \int_{-\infty}^{\infty} f(x; \theta) dx \xrightarrow{\partial_\theta} 0 = \int_{-\infty}^{\infty} \partial_\theta f(x; \theta) dx. \quad (4.23)$$

Next,

$$\partial_\theta f(x; \theta) = \frac{\partial_\theta f(x; \theta)}{f(x; \theta)} f(x; \theta) \implies 0 = \int_{-\infty}^{\infty} \partial_\theta \ln f(x; \theta) f(x; \theta) dx. \quad (4.24)$$

And so writing this as an expectation:

$$E[\partial_\theta \ln f(X; \theta)] = 0. \quad (4.25)$$

Now, if we take the second derivative of the identity integral we get

$$0 = \int_{-\infty}^{\infty} \partial_\theta^2 \ln f(x; \theta) f(x; \theta) dx + \int_{-\infty}^{\infty} (\partial_\theta \ln f(x; \theta))^2 f(x; \theta) dx. \quad (4.26)$$

The second term on the RHS can be written as an expectation, called the **Fisher information**, denoted  $I(\theta)$ :

$$I(\theta) = E[(\partial_\theta \ln f(X; \theta))^2] = -E[\partial_\theta^2 \ln f(X; \theta)] \quad (4.27)$$

Now, because  $E[\partial_\theta \ln f(x; \theta)] = 0$  we can see that

$$I(\theta) = \text{Var}[\partial_\theta \ln f(X; \theta)] \quad (4.28)$$

The important function

$$\partial_\theta \ln f(x; \theta) \quad (4.29)$$

is called the **score function**. Recall that it determines the estimating equations for the mle, i.e., the mle  $\hat{\theta}$  solves

$$\sum_{i=1}^n \partial_\theta \ln f(x_i; \theta) = 0. \quad (4.30)$$

For an  $n$ -sample of iid r.v., the Fisher information is

$$\mathcal{I}(\theta) = nI(\theta) = \text{Var}(\partial_\theta \ln \mathcal{L}(\theta, \mathbf{X})). \quad (4.31)$$

**Theorem 4.2.1. (Rao-Cramér Lower Bound.)** Let  $X_1, \dots, X_n$  be iid with pdf  $f(x; \theta), \theta \in \Omega$ . Assume that all four regularity conditions hold. Let  $Y = u(X_1, \dots, X_n)$  be a statistic with mean  $E[Y] = k(\theta)$ . Then

$$\text{Var}(Y) \geq \frac{[k'(\theta)]^2}{nI(\theta)} \quad (4.32)$$

*Proof.* Here's a sketch of the proof. Define

$$Z = \sum_{i=1}^n \partial_\theta \ln f(X_i; \theta). \quad (4.33)$$

Then  $E[Z] = 0$  and  $\text{Var}[Z] = nI(\theta)$ . Now, verify that

$$k'(\theta) = E[YZ] = E[Y]E[Z] + \rho\sigma_Y\sqrt{nI(\theta)} \quad (4.34)$$

where  $\rho$  is the correlation coefficient between  $Y$  and  $Z$ . Using  $E[Z] = 0$  and rearrange, we get the desired result.  $\square$

**Theorem 4.2.2.** Under the assumptions of the Rao-Cramér lower bound theorem, if  $Y = u(X_1, \dots, X_n)$  is an unbiased estimator of  $\theta$ , so that  $k(\theta) = \theta$ , then the Rao-Cramér inequality becomes

$$\text{Var}(Y) \geq \frac{1}{nI(\theta)} \quad (4.35)$$

**Example 4.2.1.** Consider a single observation from a Poisson- $\lambda$ . We want to find  $I(\lambda)$ . Well,

$$P(x_i; \lambda) = \frac{e^{-\lambda}\lambda^{x_i}}{x_i!}, \quad x_i \in \mathbb{N}, \lambda > 0 \in \Omega. \quad (4.36)$$

The regularity conditions are satisfied. Now,

$$I(\lambda) = E \left[ (\partial_\lambda \ln P(x_i, \lambda))^2 \right]. \quad (4.37)$$

We can find

$$\partial_\lambda \ln P(x_i, \lambda) = -1 + \frac{x_i}{\lambda}. \quad (4.38)$$

And so

$$\begin{aligned} I(\lambda) &= E \left[ x_i^2/\lambda^2 - 2x_i/\lambda + 1 \right] = \frac{E[x_i^2]}{\lambda^2} - \frac{2}{\lambda}E[x_i] + 1 \\ &= \frac{1}{\lambda^2}(\lambda + \lambda^2) - \frac{2}{\lambda}\lambda + 1 = \frac{1}{\lambda}. \end{aligned} \quad (4.39)$$

We can do it using the other expectation too, but let's not worry about that. What about a sample of  $n$  iid such r.v.'s? The answer is just  $n/\lambda$ .

**Example 4.2.2.** Let  $X_i \sim \text{Poi}(\lambda)$ . Find  $\hat{\mu}_{ML}$ . Well,

$$l(\lambda) = -n\lambda + \sum_{i=1}^n x_i \ln \lambda - \sum \ln x_i!. \quad (4.40)$$

And so,

$$\partial_\lambda l(\lambda) = -n + \frac{1}{\lambda} \sum_{i=1}^n x_i = 0 \implies \hat{\mu}_{ML} = \bar{X}. \quad (4.41)$$

This is unbiased. We also know that

$$\text{Var}(\bar{X}) = \frac{\lambda}{n}. \quad (4.42)$$

How does this compare with the CRLB? We have an unbiased estimator, and so

$$\text{Var}(\bar{X}) \geq \frac{1}{nI(\lambda)} = \frac{\lambda}{n}. \quad (4.43)$$

In this case, our estimator  $\bar{X}$  does achieve the CRLB, which makes it a good mle.

**Example 4.2.3.** Let  $X_i \sim \mathcal{N}(\mu, \sigma^2)$  where  $\mu$  is unknown but  $\sigma^2$  is. What is  $nI(\mu)$ ? Well, for a single observation,

$$f(x_i; \mu) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x_i - \mu)^2/2\sigma^2}; \quad x_i, \mu \in \mathbb{R}. \quad (4.44)$$

Next,

$$l(x_i; \mu) = \frac{-1}{2} \ln(2\pi\sigma^2) - \frac{1}{2\sigma^2}(x_i - \mu)^2. \quad (4.45)$$

This the score function. The information is

$$I(\mu) = E[l(x_i; \mu)^2] = -E[\partial_\mu^2 f(x_i; \mu)] = \frac{1}{\sigma^2}. \quad (4.46)$$

And so,

$$nI(\mu) = \frac{n}{\sigma^2}. \quad (4.47)$$

And so, the CRLB for an unbiased estimator is  $\sigma^2/n$ .

**Definition 4.2.1. (Efficient Estimator.)** Let  $Y$  be an unbiased estimator of a parameter  $\theta$  in the case of point estimator. The statistic  $Y$  is called an **efficient estimator** of  $\theta$  if and only if  $\text{Var}(Y)$  attains the Rao-Cramér lower bound.

**Definition 4.2.2.** (**Efficiency.**) In cases in which we can differentiate w.r.t a parameter under an integral or summation symbol, the ratio of the Rao-Cramér lower bound to the actual variance of any unbiased estimator of a parameter is called the **efficiency** of that estimator.

**Additional Regularity Condition.** (the total is 5 after this)

- The pdf  $f(x; \theta)$  is three times differentiable as a function of  $\theta$ . Further, for all  $\theta \in \Omega$ , there exists a constant  $c$  and a function  $M(x)$  such that

$$|\partial_\theta^3 \ln f(x; \theta)| \leq M(x), \quad (4.48)$$

with  $E_{\theta_0}[M(X)] < \infty$  for all  $\theta_0 - c < \theta < \theta_0 + c$  and all  $x$  in the support of  $X$ .

**Theorem 4.2.3.** Assume  $X_1, \dots, X_n$  are iid with pdf  $f(x; \theta)$  for  $\theta_0 \in \Omega$  such that the 5 regularity conditions are satisfied. Suppose further that the Fisher information satisfies  $0 < I(\theta_0) < \infty$ . Then any consistent sequence of solutions of the mle equations satisfies

$$\sqrt{n}(\hat{\theta} - \theta_0) \xrightarrow{D} \mathcal{N}\left(0, \frac{1}{I(\theta_0)}\right) \quad (4.49)$$

In practical terms, this tells us that the MLEs for large  $n$  have a normal distribution and that they have variance that approaches CRLB.

**Definition 4.2.3.** Let  $X_1, \dots, X_n$  be iid with pdf  $f(x; \theta)$ . Suppose  $\hat{\theta}_{1n} = \hat{\theta}_{1n}(X_1, \dots, X_n)$  is an estimator of  $\theta_0$  such that  $\sqrt{n}(\hat{\theta}_{1n} - \theta_0) \xrightarrow{D} \mathcal{N}(0, \sigma_{\hat{\theta}_{1n}}^2)$ . Then

- The **asymptotic efficiency** of  $\hat{\theta}_{1n}$  is defined to be

$$e(\hat{\theta}_{1n}) = \frac{1/I(\theta_0)}{\sigma_{\hat{\theta}_{1n}}^2} \quad (4.50)$$

- The estimator  $\hat{\theta}_{1n}$  is said to be **asymptotically efficient** if the ratio in the previous item is 1.
- Let  $\hat{\theta}_{2n}$  be another estimator such that  $\sqrt{n}(\hat{\theta}_{2n} - \theta_0) \xrightarrow{D} \mathcal{N}(0, \sigma_{\hat{\theta}_{2n}}^2)$ . Then the **asymptotic relative efficiency** (ARE) of  $\hat{\theta}_{1n}$  to  $\hat{\theta}_{2n}$  is the reciprocal of the ratio of their respective asymptotic variances, i.e.,

$$e(\hat{\theta}_{1n}, \hat{\theta}_{2n}) = \frac{\sigma_{\hat{\theta}_{2n}}^2}{\sigma_{\hat{\theta}_{1n}}^2}. \quad (4.51)$$

**Note:** MLEs are asymptotically efficient estimators.

**Theorem 4.2.4.** Under the assumptions of Theorem 4.2.3., suppose  $g(x)$  is a continuous function of  $x$  that is differentiable at  $\theta_0$  such that  $g'(\theta_0) \neq 0$ . Then

$$\sqrt{n}(g(\hat{\theta}_n) - g(\theta_0)) \xrightarrow{D} \mathcal{N}\left(0, \frac{g'(\theta_0)^2}{I(\theta_0)}\right). \quad (4.52)$$

*Proof.* The proof uses previous theorems and the  $\Delta$ -method.  $\square$

**Theorem 4.2.5.** Under the assumptions of Theorem 4.2.3.,

$$\sqrt{n}(\hat{\theta}_n - \theta_0) = \frac{1}{I(\theta_0)} \frac{1}{\sqrt{n}} \sum_{i=1}^n \partial_\theta \ln f(X_i; \theta_0) + R_n \quad (4.53)$$

where  $R_n \xrightarrow{P} 0$ .

**Example 4.2.4.** Let  $X_1, \dots, X_n$  be iid  $\text{Exp}(\theta)$  r.v., Find the information contained in the sample about  $\theta$ . Well,

$$f(x_i; \theta) = \frac{1}{\theta} e^{-x_i/\theta}, x_i > 0, \theta > 0. \quad (4.54)$$

The information is given by

$$I(\theta) = -E[\partial_\theta^2 l(\theta)] = \dots = -\frac{1}{\theta^2} + \frac{2}{\theta^3} E[x_i] = \frac{1}{\theta^2}. \quad (4.55)$$

So the amount of information contained in the sample is  $nI(\theta) = n/\theta^2$ .

Now consider two estimators for  $\theta$ :  $\bar{Y}$  and  $n \min X_i$ . We know that  $\bar{Y}$  is unbiased. Let's check if  $n \min X_i$  is unbiased:  $E[n \min X_i] = nE[\min X_i]$ . Well,

$$\begin{aligned} E[\min X_i] &= \int_0^\infty y g_{\min}(y) dy \\ &= \int_0^\infty y n [1 - F(y)]^{n-1} f(y) dy \\ &= \dots \\ &= \int_0^\infty y \frac{n}{\theta} e^{-ny/\theta} dy \\ &= \frac{\theta}{n}. \end{aligned} \quad (4.56)$$

So  $E[n \min X_i] = \theta$ , unbiased.

Now, the CRLB is  $1/nI(\theta) = \theta^2/n$ , which is the smallest possible variance for an unbiased estimator. Now,  $\text{Var}(\bar{Y}) = \theta^2/n$ . And so this estimator has efficiency of 1 and is an efficient estimator.  $\text{Var}(n \min X_i)$  turns out to be  $\theta^2 \geq \theta^2/n$ . So, the relative efficiency of  $\hat{\theta}_1$  and  $\hat{\theta}_2$  is  $1/n$ . The asymptotic efficiency is going to be zero.

### 4.3 Maximum Likelihood Tests

Consider the two-sided hypothesis

$$H_0 : \theta = \theta_0 \quad H_a : \theta \neq \theta_0. \quad (4.57)$$

The log likelihood is given by

$$l(\theta) = \sum_{i=1}^n \ln f(X_i; \theta). \quad (4.58)$$

If  $\theta_0$  is the true value of  $\theta$ , then asymptotically  $\mathcal{L}(\theta_0)$  is the maximum value of  $\mathcal{L}(\theta)$ . Consider the ratio of two likelihood functions

$$\Lambda = \frac{\mathcal{L}(\theta_0)}{\mathcal{L}(\hat{\theta})} \leq 1. \quad (4.59)$$

If  $H_0$  is true,  $\Lambda$  large and is close to 1. If  $H_1$  is true,  $\Lambda$  is smaller. For a specified significance level  $\alpha$ , we get a decision rule: Reject  $H_0$  if  $\Lambda \leq c$  where  $c$  is such that  $\alpha = P_{\theta_0}(\Lambda \leq c)$ . This is called the **likelihood ratio test**.

**Theorem 4.3.1.** Assume the same regularity conditions as for Theorem 4.2.3, under the null hypothesis  $H_0 : \theta = \theta_0$ ,

$$-2 \ln \Lambda \xrightarrow{D} \chi^2(1). \quad (4.60)$$

*Proof.* The proof uses Taylor expansion about  $\theta_0$  of  $l(\theta)$  to first order, evaluated at the mle  $\hat{\theta}$ .  $\square$

Define the test statistic  $\chi_L^2 = -2 \ln \Lambda$ . The theorem above suggests the decision rule: Reject  $H_0$  if  $\chi_L^2 \geq \chi_\alpha^2(1)$ .

**Example 4.3.1.** Let  $X_1, \dots, X_n$  be iid  $\mathcal{N}(\mu, \sigma^2)$  r.v. where both  $\mu$  and  $\sigma^2$  are unknown.  $H_0 : \mu = \mu_0$  and  $H_a : \mu \neq \mu_0$ . The null space is  $\{(\mu, \sigma^2) : \mu = \mu_0, \sigma^2 > 0\}$  and the alternative space is  $\{(\mu, \sigma^2) : \mu \neq \mu_0, \sigma^2 > 0\}$ . The union of these two spaces is  $\{(\mu, \sigma^2) : \mu \in \mathbb{R}, \sigma^2 > 0\}$ . Under the null,  $\mu = \mu_0$  and  $\sigma^2$  is unknown.

$$\partial_{\sigma^2} l(\mu_0, \sigma^2) = -\frac{n}{\sigma^2} + \frac{1}{2\sigma^4} \sum (x_i - \mu_0)^2 = 0 \implies \hat{\sigma}_0^2 = \frac{1}{n} \sum (x_i - \mu_0)^2. \quad (4.61)$$

And so

$$\mathcal{L}_0(\hat{\theta}) = (2\pi)^{-n/2} (\hat{\sigma}_0)^{-n/2} \exp \left[ -\frac{1}{2\hat{\sigma}_0^2} \sum (x_i - \mu_0)^2 \right]. \quad (4.62)$$

This is the likelihood in null space. Under the joint null alternative space where  $\mu \in \mathbb{R}$  and  $\sigma^2 > 0$ ,

$$l(\mu, \sigma^2) = -\frac{n}{2} \ln(2\bar{x}) - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2\sigma^2} \sum (x_i - \mu)^2. \quad (4.63)$$

We need to estimate both  $\mu$  and  $\sigma^2$ . We have

$$\partial_\mu l(\mu, \sigma^2) = \frac{1}{\sigma^2} \sum (x_i - \mu) = 0 \implies \hat{\mu} = \bar{x} \quad (4.64)$$

$$\partial_{\sigma^2} l(\mu, \sigma^2) = -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^2} \sum (x_i - \mu)^2 = 0 \implies \hat{\sigma}^2 = \frac{1}{n} \sum (x_i - \mu)^2. \quad (4.65)$$

The likelihood in this case is

$$\mathcal{L}(\hat{\theta}) = (2\pi)^{-n/2} (\hat{\sigma}^2)^{-n/2} \exp \left[ -\frac{1}{2\hat{\sigma}^2} \sum (x_i - \bar{x})^2 \right]. \quad (4.66)$$

Now, we reject if

$$\Lambda = \frac{\mathcal{L}_0}{\mathcal{L}} = \dots = \left( \frac{\hat{\sigma}^2}{\hat{\sigma}_0^2} \right)^{n/2} < K \quad (4.67)$$

which is equivalent to when

$$\frac{\hat{\sigma}^2}{\hat{\sigma}_0^2} = \frac{\sum (x_i - \bar{x})^2}{\sum (x_i - \mu_0)^2} < K^{2/n} \equiv K'. \quad (4.68)$$

Now, note that

$$\sum (x_i - \mu_0)^2 = \sum [(x_i - \bar{x}) + (\bar{x} - \mu_0)]^2 = \sum (x_i - \bar{x})^2 + n(\bar{x} - \mu_0)^2. \quad (4.69)$$

So, we reject whenever

$$\frac{\sum (x_i - \bar{x})^2}{\sum (x_i - \bar{x})^2 + n(\bar{x} - \mu_0)^2} < K^{2/n} = \frac{1}{1 + \frac{n(\bar{x} - \mu_0)^2}{\sum (x_i - \bar{x})^2}} < K' \quad (4.70)$$

which is equivalent to

$$\frac{n(\bar{x} - \mu_0)^2}{\sum (x_i - \bar{x})^2} > \frac{1}{K} - 1 \equiv K'' \iff \frac{n(\bar{x} - \mu_0)^2}{\frac{1}{n-1} \sum (x_i - \bar{x})^2} = \frac{n(\bar{x} - \mu_0)^2}{S^2} > (n-1)K'', \quad (4.71)$$

where  $K''$  is another constant. So we reject whenever

$$\left| \frac{\bar{x} - \mu_0}{S/\sqrt{n}} \right| > \sqrt{(n-1)K''} \quad (4.72)$$

where the LHS is a  $t$ -statistic with  $(n-1)$  df.

In practice, there are two other likelihood-related tests. A natural statistic is based on the asymptotic distribution of  $\hat{\theta}$ . Consider the statistic

$$\chi_W^2 = \left\{ \sqrt{nI(\hat{\theta})}(\hat{\theta} - \theta_0) \right\}^2. \quad (4.73)$$

Because  $I(\theta)$  is a continuous function,  $I(\hat{\theta}) \xrightarrow{P} I(\theta_0)$  under the null hypothesis. So, under the null hypothesis,  $\chi_W^2$  has an asymptotic  $\chi^2$ -distribution with one degree of freedom. So the decision rule is to reject  $H_0$  if  $\chi_W^2 \geq \chi_\alpha^2(1)$ . This is often referred to as a **Wald**-type test.

The third test is called the **scores**-type test. The **scores** are the components of the vector

$$\mathbf{S}(\theta) = (\partial_\theta \ln f(X_1; \theta), \dots, \partial_\theta \ln f(X_n; \theta))'. \quad (4.74)$$

In our notation we have

$$\frac{1}{\sqrt{n}} l'(\theta_0) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \partial_\theta \ln f(X_i; \theta_0). \quad (4.75)$$

Define the statistic

$$\chi_R^2 = \left( \frac{I'(\theta_0)}{\sqrt{nI(\theta_0)}} \right)^2 \quad (4.76)$$

The decision rule is to reject  $H_0$  in favor of  $H_a$  if  $\chi_R^2 \geq \chi_\alpha^2(1)$ .

**Example 4.3.2.** (MLR test) Suppose we want to test whether or not the mean number of goals scored in all English league soccer games is 3. Suppose  $X \sim \text{Poi}(\lambda)$ . We set  $H_0 : \lambda = 3$  and  $H_a : \lambda \neq 3$ . The joint pdf is

$$\mathcal{L}(\lambda) = p(x_i | \lambda) = \frac{e^{-n\lambda} \lambda^{\sum x_i}}{\prod x_i!}, \quad x_i = 0, 1, 2, 3, \dots \quad (4.77)$$

Say  $n = 380$  and  $\bar{x} = 2.534$ . We can find the MLE of  $\lambda$ ,  $\hat{\lambda} = \bar{x}$ . The null space is  $\{\lambda : \lambda = 3\}$ , and the alternative space is  $\{\lambda : \lambda \neq 3\}$ . The joint space is  $\{\lambda : \lambda > 0\}$ . Now,

$$\mathcal{L}(\hat{\lambda}_0) = \frac{e^{-380 \cdot 3} 3^{963}}{\dots} \quad (4.78)$$

and

$$\mathcal{L}(\hat{\lambda}) = \frac{e^{-380 \cdot 2.534} 2.534^{963}}{\dots} \quad (4.79)$$

and so

$$\Lambda = \frac{\mathcal{L}_0}{\mathcal{L}} \quad (4.80)$$

Asymptotically,  $-2 \ln \Lambda \sim \chi^2(1)$ . We reject if  $-2 \ln \Lambda > \chi_{1,0.05}^2 = 3.84$ . We find that  $-2 \ln \Lambda = 29.03 > 3.84$ , so we reject.

**Example 4.3.3.** (Wald test)

$$\chi_W^2 = \left[ \frac{\hat{\lambda} - \lambda_0}{1/\sqrt{nI(\hat{\lambda})}} \right]^2 \sim \chi^2(1)$$

under  $H_0$ . The information can be found to be  $n/\lambda$ . Plugging in the numbers we find

$$\chi_W^2 = \left[ \frac{\bar{x} - \lambda_0}{1/\sqrt{n/\bar{x}}} \right]^2 = \left[ (2.534 - 3)/(1/\sqrt{380 \cdot 1/2.534}) \right]^2 = 32.56 > 3.84,$$

so we still reject.

**Example 4.3.4.** (Score test)

$$\chi_R^2 = \frac{(I'(\theta_0))^2}{nI(\theta_0)} = \dots = \frac{(-380 + 963/3)^2}{380/3} = 27.48 \sim \chi^2.$$

So we reject as well.

So,  $\chi_R^2$  is in general not as conservative as  $\chi_W^2$ . The LRT is somewhere in the middle.

## 4.4 Multiparameter Case: Estimation

In this section we consider the case where  $\boldsymbol{\theta}$  is a vector of  $p$  parameters. Let  $X_1, \dots, X_n$  be iid with common pdf  $f(x; \boldsymbol{\theta})$  where  $\boldsymbol{\theta} \in \Omega \subset \mathbb{R}^p$ . The likelihood function and its log are given by

$$\mathcal{L}(\boldsymbol{\theta}) = \prod_{i=1}^n f(x_i; \boldsymbol{\theta}) \tag{4.81}$$

$$l(\boldsymbol{\theta}) = \ln \mathcal{L}(\boldsymbol{\theta}) = \sum_{i=1}^n \ln f(x_i; \boldsymbol{\theta}). \tag{4.82}$$

There are more regularity conditions to make the theory work, but we won't worry about those. We will also worry only about the continuous case, because the discrete cases can be treated in similar ways.

Here are some facts: just like before,  $\mathcal{L}(\boldsymbol{\theta})$  is maximized at the true value of  $\boldsymbol{\theta}$ . The quantity that maximizes  $\mathcal{L}(\boldsymbol{\theta})$  is called the **maximum likelihood estimator** and we denote it  $\hat{\boldsymbol{\theta}}$ . It also holds that  $\hat{\eta} = g(\hat{\boldsymbol{\theta}})$  if  $\eta = g(\boldsymbol{\theta})$ .

**Theorem 4.4.1.** Let  $X_1, \dots, X_n$  be iid with pdf  $f(x; \boldsymbol{\theta})$  for  $\boldsymbol{\theta} \in \Omega$ . Assume that the regularity conditions hold, then

1. The likelihood equation,

$$\partial_{\boldsymbol{\theta}} l(\boldsymbol{\theta}) = \mathbf{0} \quad (4.83)$$

has a solution  $\hat{\boldsymbol{\theta}}_n$  such that  $\hat{\boldsymbol{\theta}} \xrightarrow{P} \boldsymbol{\theta}$ .

2. For any sequence that satisfies the first item

$$\sqrt{n}(\hat{\boldsymbol{\theta}}_n - \boldsymbol{\theta}) \xrightarrow{D} \mathcal{N}_p(\mathbf{0}, \mathbf{I}^{-1}(\boldsymbol{\theta})). \quad (4.84)$$

**Theorem 4.4.2.** Let  $X_1, \dots, X_n$  be iid with pdf  $f(x; \boldsymbol{\theta})$  for  $\boldsymbol{\theta} \in \Omega$ . Assume the regularity conditions hold. Let  $\hat{\boldsymbol{\theta}}_n$  be a sequence of consistent solutions of the likelihood equation. Then  $\hat{\boldsymbol{\theta}}_n$  are asymptotically efficient estimates; that is, for  $j = 1, \dots, p$ ,

$$\sqrt{n}(\hat{\theta}_{i,j} - \theta_j) \xrightarrow{D} \mathcal{N}(0, [\mathbf{I}^{-1}(\boldsymbol{\theta})]_{jj}). \quad (4.85)$$

**Example 4.4.1.** Let  $X_i$  be iid  $\mathcal{N}(\mu, \sigma^2)$ . Find MLE for  $\mu$  and  $\sigma^2$ . Now,  $\boldsymbol{\theta} = (\mu, \sigma^2)$ , and

$$\mathcal{L}(\boldsymbol{\theta}) = \prod f(x_i | \boldsymbol{\theta}) = \prod \left[ (2\pi\sigma^2)^{1/2} \exp\left(-\frac{1}{2\sigma^2}(x_i - \mu)^2\right) \right]. \quad (4.86)$$

Then

$$\partial_{\mu} l(\boldsymbol{\theta}) = \frac{1}{\sigma^2} \sum (x_i - \mu) = 0 \implies \hat{\mu} = \bar{x} \quad (4.87)$$

and

$$\partial_{\sigma^2} l(\boldsymbol{\theta}) = \frac{-n}{2\sigma^2} + \frac{1}{2\sigma^4} \sum (x_i - \mu)^2 = 0 \implies \hat{\sigma}^2 = \frac{1}{n} \sum (x_i - \mu)^2. \quad (4.88)$$

What is the information in the multiparameter setting? It's a matrix with elements

$$I_{jk} = -E \left[ \frac{\partial^2 \ln f(x_i | \boldsymbol{\theta})}{\partial \theta_j \partial \theta_k} \right] \quad (4.89)$$

Back to the normal situation:

$$I(\boldsymbol{\theta}) = -E \begin{bmatrix} -\frac{1}{\sigma^2} & -\frac{1}{\sigma^4}(x_i - \mu) \\ -\frac{1}{\sigma^4}(x_i - \mu) & \frac{1}{2\sigma^4} - \frac{1}{\sigma^6}(x_i - \mu)^2 \end{bmatrix} \quad (4.90)$$

$$= \begin{bmatrix} \frac{1}{\sigma^2} & 0 \\ 0 & -\frac{1}{2\sigma^4} + \frac{1}{\sigma^6}\sigma^2 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sigma^2} & 0 \\ 0 & \frac{1}{2\sigma^4} \end{bmatrix} \quad (4.91)$$

CRLB for  $\bar{x}$  is

$$\frac{1}{nI_{11}(\boldsymbol{\theta})} = \frac{\sigma^2}{n}. \quad (4.92)$$

The CRLB for  $\hat{\sigma}^2$  is

$$\frac{1}{nI_{22}(\boldsymbol{\theta})} = \frac{2\sigma^4}{n}. \quad (4.93)$$

## 4.5 Multiparameter Case: Testing

Here we talk about likelihood ratio tests in the multiparameter setting. Recall that the test statistic is

$$\Lambda = \frac{\mathcal{L}(\hat{\theta}_0)}{\mathcal{L}(\hat{\theta})} \quad (4.94)$$

where the numerator is maximized in null space and the denominator is maximized in the joint (null + alternative) space. We reject if  $\Lambda < c$ .  $c$  is found from the null distribution.

**Example 4.5.1.** Let  $X_i \sim \exp(\theta)$  and  $Y_i \sim \exp(\mu)$ .  $H_0 : \mu = \theta$  and  $H_a : \mu \neq \theta$ . The null space is  $\{\{\theta, \mu\} : \theta = \mu > 0\}$ . The alternative space is  $\{\{\theta, \mu\} : \theta \neq \mu, \theta, \mu \neq 0\}$ . The joint space is just  $\{\{\theta, \mu\}, \theta > 0, \mu > 0\}$ . It's easy to see that

$$\mathcal{L}(\theta, \mu | X, Y) = \frac{1}{\theta^n} e^{\sum^n x_i / \theta} \frac{1}{\mu^m} e^{\sum^m y_i / \mu}. \quad (4.95)$$

From here one finds that under  $H_0 : \theta = \mu$ :

$$l(\Theta_0) = -(n+m) \ln \theta - \frac{1}{\theta} \left( \sum x_i + \sum y_i \right). \quad (4.96)$$

And so

$$\partial_\theta l(\theta) = 0 \implies \hat{\theta}_0 = \frac{1}{n+m} \left( \sum x_i + \sum y_i \right). \quad (4.97)$$

With this, we can plug back in to calculate the numerator:

$$\mathcal{L}(\hat{\Theta}_0) = \dots = \left( \frac{1}{n+m} \sum x_i + \sum y_i \right)^{-n-m} \exp(-n-m). \quad (4.98)$$

In the joint space,

$$l(\Theta) = -n \ln \theta - \frac{1}{\theta} \sum x_i - m \ln \mu - \frac{1}{\mu} \sum y_j. \quad (4.99)$$

And so

$$\partial_\theta l(\Theta) = 0 \implies \hat{\theta} = \bar{x} \quad (4.100)$$

$$\partial_\mu l(\mu) = 0 \implies \hat{\mu} = \bar{y}. \quad (4.101)$$

With these,

$$\mathcal{L}(\hat{\Theta}) = \dots \quad (4.102)$$

$$= \frac{1}{(\frac{1}{n} \sum x_i)^n} \exp \left[ -\frac{1}{\frac{1}{n} \sum x_i} \sum x_i \right] \frac{1}{(\frac{1}{m} \sum y_i)^m} \exp \left[ -\frac{1}{\frac{1}{m} \sum y_i} \sum y_i \right] \quad (4.103)$$

$$= \frac{1}{(\frac{1}{n} \sum x_i)^n} e^{-n} \frac{1}{(\frac{1}{m} \sum y_i)^m} e^{-m}. \quad (4.104)$$

Putting everything together, we find

$$\Lambda = \frac{\mathcal{L}_0}{\mathcal{L}} = \dots = \frac{\frac{1}{(n+m)^{-n-m}} (\sum x_i + \sum y_j)^{-n-m}}{\frac{1}{n^n} \frac{1}{m^m} (\sum x_i)^{-n} (\sum y_i)^{-m}}. \quad (4.105)$$

We reject if  $\Lambda < c$ , iff

$$\frac{(n+m)^{n+m}}{n^n m^m} \frac{(\sum x_i)^n (\sum y_i)^m}{(\sum x_i + \sum y_j)^{n+m}} < c \quad (4.106)$$

iff (letting  $c$  absorb the constant)

$$\frac{(\sum x_i)^n (\sum y_i)^m}{(\sum x_i + \sum y_j)^{n+m}} < c' \quad (4.107)$$

What does the distribution of  $\Lambda$  look like? Notice that we reject if

$$\Lambda = \frac{(n+m)^{n+m}}{n^n m^m} \frac{(\sum x_i)^n (\sum y_i)^m}{(\sum x_i + \sum y_j)^{n+m}} \quad (4.108)$$

$$= \frac{(n+m)^{n+m}}{n^n m^m} \left( \frac{\sum x_i}{\sum x_i + \sum y_j} \right)^n \left( \frac{\sum y_i}{\sum x_i + \sum y_j} \right)^m \quad (4.109)$$

$$= \frac{(n+m)^{n+m}}{n^n m^m} T^n (1-T)^m < c \quad (4.110)$$

which is akin to saying that we reject if  $T < a$  or  $T > b$  when these constants satisfy

$$a^n (1-a)^m = b^n (1-b)^n. \quad (4.111)$$

What is the distribution of  $T$  under  $H_0$ ? Note that under  $H_0$ ,  $\sum x_i \sim \Gamma(n, \theta)$  and  $\sum y_i \sim \Gamma(m, \mu = \theta)$ . Use transformation method to show that  $T \sim \beta(n, m)$ .

Reminder: in large samples,  $-2 \ln \Lambda \sim \chi_1^2$

## 4.6 The EM algorithm

# Part 5

# Sufficiency

5.1 Measures of Quality of Estimators

5.2 A Sufficient Statistic for a Parameter



# **Part 6**

# **Problems**

## 6.1 Problem Set 1

### 3.6.4

- (a)  $X$  has a standard normal distribution:

```
x=seq(-6,6,.01); plot(dnorm(x)^x)
```

- (b)  $X$  has a  $t$ -distribution with 1 degree of freedom.

```
lines(dt(x,1)^x,lty=2)
```

- (c)  $X$  has a  $t$ -distribution with 3 degrees of freedom.


```
lines(dt(x,3)^x,lty=2)
```

- (d)  $X$  has a  $t$ -distribution with 10 degrees of freedom.


```
lines(dt(x,10)^x,lty=2)
```

- (e)  $X$  has a  $t$ -distribution with 30 degrees of freedom.


```
lines(dt(x,30)^x,lty=2)
```




(a)




(b)



(c)



(d)



(e)

**3.6.5**

(a)  $P(|X| \geq 2) = 2 \times [1 - P(X \leq 2)] = \mathbf{0.046}.$

```
> 2*(1 - pnorm(2))  
[1] 0.04550026
```

(b)  $P(|X| \geq 2) = 2 \times [1 - P(X \leq 2)] = \mathbf{0.295}.$

```
> 2*(1 - pt(2,1))  
[1] 0.2951672
```

(c)  $P(|X| \geq 2) = 2 \times [1 - P(X \leq 2)] = \mathbf{0.139}.$

```
> 2*(1 - pt(2,3))  
[1] 0.139326
```

(d)  $P(|X| \geq 2) = 2 \times [1 - P(X \leq 2)] = \mathbf{0.073}.$

```
> 2*(1 - pt(2,10))  
[1] 0.07338803
```

(e)  $P(|X| \geq 2) = 2 \times [1 - P(X \leq 2)] = \mathbf{0.055}.$

```
> 2*(1 - pt(2,30))  
[1] 0.05462504
```

**3.6.11:** Let  $T = W/\sqrt{V/r}$ , where the independent variables  $W \sim \mathcal{N}(0, 1)$  and  $V \sim \chi^2(r)$ . Show that  $T^2 \sim F(r_1 = 1, r_2 = r)$ . *Hint:* What is the distribution of the numerator of  $T^2$ ?

*Solution:* Let the independent random variables  $U, V$  be given, with  $W \sim \mathcal{N}(0, 1)$  and  $U \sim \chi^2(r)$ . The random variable  $T^2$ , where  $T = W/\sqrt{V/r}$  is given by

$$T^2 = \left( \frac{W}{\sqrt{V/r}} \right)^2 = \frac{W^2}{V/r}. \quad (6.1)$$

Because  $W \sim \mathcal{N}(0, 1)$ , we have that  $W^2 \sim \chi^2(1)$  (by theorem). Now,  $T^2$  has the form

$$T^2 = \frac{W^2}{V/r} = \frac{W^2/1}{V/r} \quad (6.2)$$

where 1 is the df of  $\chi^2(1)$  which  $W$  follows, and  $r$  is the df of  $\chi^2(r)$  which  $U$  follows. Thus,  $T^2 \sim F(1, r)$ , by the definition of the  $F$ -distribution.  $\square$

**3.6.15:** Let  $X_1, X_2$  be iid with common distribution having the pdf

$$f(x) = \begin{cases} e^{-x}, & 0 < x < \infty \\ 0, & \text{else} \end{cases} \quad (6.3)$$

Show that  $Z = X_1/X_2$  has an  $F$ -distribution.

*Solution:* It suffices to show that  $Z$  can be written as a ratio of two  $\chi^2$ -distributed independent random variables. To this end, we can consider the mgf  $M_X(t)$  of  $X_1$ , which is also identically that of  $X_2$  since  $X_1, X_2$  are iid:

$$M_X(t) = E[e^{tx}] = \int_0^\infty e^{tx} e^{-x} dx = (1-t)^{-1}. \quad (6.4)$$

However, this does not quite match the mgf for a  $\chi^2(2)$ . To circumvent this problem, we rewrite

$$Z = \frac{X_1}{X_2} = \frac{2X_1/2}{2X_2/2} = \frac{(X_1 + X_1)/2}{(X_2 + X_2)/2}, \quad (6.5)$$

as we expect  $r = 2$ . Let  $Y_1 = X_1 + X_1$ . Then we have trivially  $Y_1 = 2X_1$ , and so  $|J| = 1/2$ . With this,  $Y_1$  has the pdf

$$\tilde{f}_Y(y) = |J|f(x) = \frac{1}{2}f(x) = \begin{cases} \frac{1}{2}e^{-y/2}, & 0 < y < \infty \\ 0, & \text{else} \end{cases}. \quad (6.6)$$

From here, we find the mgf of  $Y_1$  to be

$$M_{Y_1}(t) = E[e^{ty}] = \frac{1}{2} \int_0^\infty e^{ty} e^{-y/2} dy = (1-2t)^{-1} = (1-2t)^{-2/2}, t < \frac{1}{2}. \quad (6.7)$$

By symmetry,  $M_{Y_2}(t)$  is identically  $M_{Y_1}(t)$ , and both are the mgf for  $\chi^2(r = 2)$ . Because each mgf uniquely determines a pdf,  $Y_1, Y_2 \sim \chi^2(r = 2)$  identically and independently (for each depends exclusively on  $X_1, X_2$ , respectively). Therefore,

$$Z = \frac{(X_1 + X_1)/2}{(X_2 + X_2)/2} = \frac{Y_1/2}{Y_2/2} \quad (6.8)$$

follows the  $F$ -distribution with degrees of freedom  $r_1 = r_2 = 2$ , by definition.  $\square$

**3.6.16:** Let  $X_1, X_2, X_3$  be independent r.v. with  $X_i \sim \chi^2(r_i)$ .

- (a) Show that  $Y_1 = X_1/X_2$  and  $Y_2 = X_1 + X_2$  are independent and that  $Y_2 \sim \chi^2(r_1 + r_2)$ .
- (b) Deduce that

$$\frac{X_1/r_1}{X_2/r_2} \text{ and } \frac{X_3/r_3}{(X_1 + X_2)/(r_1 + r_2)} \quad (6.9)$$

are independent  $F$ -variables.

*Solution:*

- (a) We consider the transformation

$$y_1 = u(x_1, x_2) = \frac{x_1}{x_2} \quad (6.10)$$

$$y_2 = v(x_1, x_2) = x_1 + x_2. \quad (6.11)$$

whose inverse is

$$\begin{aligned} x_1 &= \bar{u}(y_1, y_2) = \frac{y_1 y_2}{1 + y_1} \\ x_2 &= \bar{v}(y_1, y_2) = \frac{y_2}{1 + y_1}. \end{aligned} \quad (6.12)$$

The absolute value of the Jacobian is

$$|J| = \left| \det \begin{pmatrix} \partial_{y_1} \bar{u} & \partial_{y_2} \bar{u} \\ \partial_{y_1} \bar{v} & \partial_{y_2} \bar{v} \end{pmatrix} \right| = \frac{y_2}{(1 + y_1)^2}, \quad (6.13)$$

which maps one-to-one from the space of  $X_1, X_2 \in \mathbb{R}^+ \times \mathbb{R}^+$  onto the space of  $Y_1, Y_2 \in \mathbb{R}^+ \times \mathbb{R}^+$ . Since  $X_1, X_2$  are independent, we consider the joint pdf of  $X_1, X_2$ :

$$h(x_1, x_2) = \begin{cases} \frac{x_1^{r_1/2-1} x_2^{r_2/2-1}}{\Gamma(r_1/2)\Gamma(r_2/2)2^{(r_1+r_2)/2}} e^{-(x_1+x_2)/2}, & 0 < x_1, x_2 < \infty \\ 0, & \text{else} \end{cases} \quad (6.14)$$

from which we can deduce the joint pdf for  $Y_1, Y_2$ :

$$\begin{aligned} \tilde{h}(y_1, y_2) &= |J|h\left(\frac{y_1 y_2}{1 + y_1}, \frac{y_2}{1 + y_1}\right) \\ &= \begin{cases} \frac{y_2(y_1 y_2)^{r_1/2-1} y_2^{r_2/2-1} (1+y_1)^{-r_1/2-r_2/2+\frac{1}{2}}}{(1+y_1)^2 \Gamma(r_1/2)\Gamma(r_2/2)2^{(r_1+r_2)/2}} e^{-y_2/2}, & 0 < y_1, y_2 < \infty \\ 0, & \text{else} \end{cases} \\ &= \begin{cases} \frac{y_2^{r_1/2+r_2/2-1} y_1^{r_1/2-1} (1+y_1)^{-r_1/2-r_2/2}}{\Gamma(r_1/2)\Gamma(r_2/2)2^{(r_1+r_2)/2}} e^{-y_2/2}, & 0 < y_1, y_2 < \infty \\ 0, & \text{else} \end{cases} \end{aligned} \quad (6.15)$$

Without further computation we see that  $\tilde{h}(y_1, y_2)$  can be written as a product of two nonnegative functions of  $y_1$  and  $y_2$ . In view of Theorem 2.4.1,  $Y_1$  and  $Y_2$  are independent.  $\square$

Next, we wish to show  $Y_2 \sim \chi^2(X_1, X_2)$ , to which end we find the marginal pdf  $g_2(y_2)$  of  $Y_2$ :

$$\begin{aligned} g_2(y_2) &= \int_0^\infty \tilde{h}(y_1, y_2) dy_1 \\ &= \mathfrak{C} \int_0^\infty y_1^{r_1/2-1} (1+y_1)^{-r_1/2-r_2/2} dy_1 \\ &= \mathfrak{C} \frac{\Gamma(r_1/2)\Gamma(r_2/2)}{\Gamma[(r_1+r_2)/2]} \end{aligned} \quad (6.16)$$

where  $\mathfrak{C}$  contains all the  $y_1$ -independent elements. From here, via simple back-substitution we obtain the marginal pdf for  $Y_2$ :

$$g_2(y_2) = \begin{cases} \frac{y_2^{(r_1+r_2)/2-1}}{\Gamma[(r_1+r_2)/2]2^{(r_1+r_2)/2}} e^{-y_2/2}, & 0 < y_2 < \infty \\ 0, & \text{else} \end{cases}, \quad (6.17)$$

i.e.,  $Y_2 \sim \chi^2(r_1 + r_2)$ .  $\square$

#### Mathematica code:

```
In[20]:= Integrate[
x^(r1/2 - 1) (1 + x)^(-r1/2 - r2/2), {x, 0, Infinity}]

Out[20]= ConditionalExpression[(Gamma[r1/2] Gamma[r2/2])/
Gamma[(r1 + r2)/2], Re[r2] > 0 && Re[r1] > 0]
```

- (b) By definition, because  $X_1, X_2$  are independent random variables with  $X_i \sim \chi^2(r_i)$ ,

$$\Omega = \frac{X_1/r_1}{X_2/r_2} \sim F(r_1, r_2). \quad (6.18)$$

Also, because  $X_3 \sim \chi^2(r_3)$  and  $(X_1 + X_2) \sim \chi^2(r_1 + r_2)$  (from (a)), we have

$$\Lambda = \frac{X_3/r_3}{(X_1 + X_2)/(r_1 + r_2)} \sim F(r_3, r_1 + r_2) \quad (6.19)$$

as well. Furthermore, because

$$\Omega = \frac{X_1/r_1}{X_2/r_2} = \frac{r_2}{r_1} Y_1 \quad (6.20)$$

$$\Lambda = \frac{r_1 + r_2}{r_3} \frac{X_3}{Y_2} \quad (6.21)$$

and because  $X_1, X_2, X_3$  are independent, we have that  $Y_1, Y_2, X_3$  are independent. Therefore, it is necessary that  $\Omega \sim F(r_1, r_2)$  and  $\Lambda \sim F(r_3, r_1 + r_2)$  are independent as well.  $\square$

**4.1.1** Twenty motors were put on test under a high-temperature setting. The lifetimes in hours of the motors under these conditions are given below. Also, the data are in the file **lifetimemotor.rda** at the site listed in the Preface. Suppose we assume that the lifetime of a motor under these conditions,  $X$ , has a  $\Gamma(1, \theta)$  distribution.

1	4	5	21	22	28	40	42	51	53
58	67	95	124	124	160	202	260	303	363

- (a) Obtain a histogram of the data and overlay it with a density estimate, using the code `hist(x,pr=T); lines(density(x))` where the R vector  $x$  contains the data. Based on this plot, do you think that the  $\Gamma(1, \theta)$  model is credible?
- (b) Assuming a  $\Gamma(1, \theta)$  model, obtain the maximum likelihood estimate  $\hat{\theta}$  of  $\theta$  and locate it on your histogram. Next overlay the pdf of a  $\Gamma(1, \hat{\theta})$  distribution on the histogram. Use the R function `dgamma(x,shape=1,scale=hat_theta)` to evaluate the pdf.
- (c) Obtain the sample median of the data, which is an estimate of the median lifetime of a motor. What parameter is it estimating (i.e., determine the median of  $X$ )?
- (d) Based on the mle, what is another estimate of the median of  $X$ ?

*Solution:*

- (a) For some reason R does not recognize the dataset as of numeric type. Because the dataset is small enough, I recoded and fed it by hand to the data vector  $y$ :

```
> lines(density(y))
> y <- c(1,4,5,21,22,28,40,42,51,53,58,67,
95,124,124,160,202,260,303,363)
> hist(y,pr=T)
> lines(density(y))
```


The  $\Gamma(1, \theta)$ , or  $\text{Exp}(\theta)$ , model seems to be **credible** as far as the histogram is concerned. However, the overlaying density does not look like a  $\Gamma(1, \theta)$ .  $\square$

- (b) Assuming the  $\Gamma(1, \theta)$  model, then the pdf on the support  $\mathbb{R}^+$  is given by

$$f(y) = \frac{1}{\theta} e^{-y/\theta}, \quad (6.22)$$

from which we obtain the logarithm of the likelihood function:

$$l(\theta) = \log \left( \prod_{i=1}^n \frac{1}{\theta} e^{-y_i/\theta} \right) = -n \log \theta - \frac{1}{\theta} \sum_{i=1}^n y_i. \quad (6.23)$$



The first partial derivative wrt  $\theta$  is then

$$\partial_\theta l(\theta) = -\frac{n}{\theta} + \frac{1}{\theta^2} \sum_{i=1}^n y_i. \quad (6.24)$$

Setting  $\partial_\theta l(\theta) = 0$ , we get (by inspection) that  $l(\theta)$  is extremized iff  $\theta = (1/n) \sum_{i=1}^n y_i = \bar{y}$ . We also have that  $\partial_{\theta\theta} < 0 \forall \theta \in \mathbb{R}^+$ , which means  $l(\theta)$  is maximized globally at  $\bar{y}$ . From here, the statistic

$$\hat{\theta} = \bar{Y} = \mathbf{101.15} \quad (6.25)$$

is the mle of  $\theta$ . (Also note that because  $E[Y] = \theta \implies E[\bar{Y}] = \theta$ ,  $\hat{\theta}$  is an unbiased estimator of  $\theta$ .)

```
> mean(y)
[1] 101.15
> abline(v = mean(y), lwd=3, lty=2)
> z=dgamma(y, shape=1, scale=mean(y))
> lines(z~y, lty=2)
```


(c) The sample median of the data is **55.5**

```
> median(y)
[1] 55.5
```

The median of  $Y \sim \Gamma(1, \theta) \equiv \text{Exp}(\theta)$  is the value of  $y'$  at which

$$0.5 = \int_0^{y'} \frac{1}{\theta} e^{-y'/\theta} dy = 1 - e^{-y'/\theta} \implies y' = \theta \ln 2, \quad (6.26)$$

which means that the median of  $Y \sim \Gamma(1, \theta) \equiv \text{Exp}(1, \theta)$  is the half-life,  $\theta \ln 2$ . Since the sample median is just  $\theta$  multiplied by  $\ln 2$ , the sample median also estimates the parameter  $\theta$ .



- (d) From part (a), we know that  $\hat{\theta} = \bar{Y}$ , the sample mean, is the mle of  $\theta$ , the population mean. From part (c), we have shown that the median of  $Y \sim \Gamma(1, \theta)$  is simply  $\theta \ln 2$ . By simple inspection we see that  $\hat{\theta} \ln 2 = \bar{Y} \ln 2$  is the (*unbiased*) mle of  $\theta \ln 2$ , the median of  $Y$ .  $\square$

**4.1.3** Suppose the number of customers  $X$  that enter a store between the hours 9:00 a.m. and 10:00 a.m. follows a Poisson distribution with parameter  $\theta$ . Suppose a random sample of the number of customers that enter the store between 9:00 a.m. and 10:00 a.m. for 10 days results in the values

9    7    9    15    10    13    11    7    2    12

1. Determine the maximum likelihood estimate of  $\theta$ . Show that it is an unbiased estimator.
2. Based on these data, obtain the realization of your estimator in part (a). Explain the meaning of this estimate in terms of the number of customers.

*Solution:*

1. Let  $X \sim \text{Poi}(\theta)$  be given, then the pmf of  $X$  is given by

$$p(x) = \begin{cases} \frac{\theta^x e^{-\theta}}{x!}, & x \in \mathbb{N} \\ 0, & \text{else} \end{cases}. \quad (6.27)$$

Assuming the  $X_i$ 's  $\sim \text{Poi}(\theta)$  are iid, where  $i = 1, \dots, n$ , then the logarithm of the likelihood function is

$$\begin{aligned} l(\theta) &= \log \left( \prod_{i=1}^n \frac{\theta^{x_i} e^{-\theta}}{x_i!} \right) \\ &= \log \left( e^{-n\theta} \theta^{\sum_{i=1}^n x_i} \prod_{i=1}^n \frac{1}{x_i!} \right) \\ &= -n\theta + \left( \sum_{i=1}^n x_i \right) \log \theta - \sum_{i=1}^n \log x_i!. \end{aligned} \quad (6.28)$$

Setting  $\partial_\theta l(\theta) = 0$ , we solve for  $\theta$ :

$$\partial_\theta l(\theta) = -n + \frac{1}{\theta} \sum_{i=1}^n x_i = 0 \iff \theta = \frac{1}{n} \sum_{i=1}^n x_i = \bar{x} \quad (6.29)$$

By inspection,  $\partial_{\theta\theta} l(\theta) < 0 \forall \theta \in \mathbb{R}^+$ , and so the statistic

$$\hat{\theta} = \bar{Y} \quad (6.30)$$

is the mle of  $\theta$ . Further, it is an unbiased estimator of  $\theta$  simply because

$$E[Y] = \theta \implies E[\bar{Y}] = \theta. \quad (6.31)$$

□

2. Part (a) says the sample means is the mle of  $\theta$ . The means of the given sample is **9.5**.

```
> mean(c(9 ,7, 9, 15, 10, 13, 11, 7, 2, 12))  
[1] 9.5
```

This says that on average, 9.5, or about 9-10 customers enter the store between the hours 9:00 a.m. and 10:00 a.m.. □

**4.1.8** Recall that for the parameter  $\eta = g(\theta)$ , the mle of  $\eta$  is  $g(\hat{\theta})$ , where  $\hat{\theta}$  is the mle of  $\theta$ . Assuming that the data in Example 4.1.6 were drawn from a Poisson distribution with mean  $\lambda$ , obtain the mle of  $\lambda$  and then use it to obtain the mle of the pmf. Compare the mle of the pmf to the nonparametric estimate. Note: For the domain value 6, obtain the mle of  $P(X \geq 6)$ .

*Solution:* Based on the previous problem, the mle of  $\lambda$  is the sample means, which has the value **2.13**.

```
> mean(c(2,1,1,1,1,5,1,1,3,0,2,1,1,3,4,2,1,2,2,6,5,2,3,2,4,1,3,1,3,0))
[1] 2.133333
```

Because the sample means  $\bar{x}$  is the mle of  $\lambda$ , and the pmf is given by

$$p(x) = \begin{cases} \frac{\lambda^x e^{-\lambda}}{x!}, & x \in \mathbb{N} \\ 0, & \text{else} \end{cases}, \quad (6.32)$$

the mle of the pmf is given by

$$\tilde{p}(x) = \begin{cases} \frac{\bar{x}^x e^{-\bar{x}}}{x!}, & x \in \mathbb{N} \\ 0, & \text{else} \end{cases}. \quad (6.33)$$

Next, we compare the mle of the pmf to the nonparametric estimate:

j	0	1	2	3	4	5	$\geq 6$
$\hat{p}(j)$	0.067	0.367	0.233	0.167	0.067	0.067	0.033
$\tilde{p}(j)$	0.118	0.253	0.270	0.192	0.102	0.044	0.022

**Mathematica code for  $P(j \geq 6)$  for  $\tilde{p}(j)$ :**

```
P[x_] := (2.1333333)^x*E^(-2.1333333)/x!
N[Sum[P[y], {y, 6, Infinity}]]
0.0218705
```

## 6.2 Problem Set 2

**4.2.2.** Consider the data on the lifetimes of motors given in Exercise 4.1.1. Obtain a large sample 95% confidence interval for the mean lifetime of a motor.

*Solution:* Large sample 95% CI's have the form

$$(\bar{x} - z_{\alpha/2} \frac{S}{\sqrt{n}}, \bar{x} + z_{\alpha/2} \frac{S}{\sqrt{n}}) \quad (6.34)$$

Here,  $\bar{x} = 101.15$ ,  $n = 20$ ,  $s = 105.4091$ ,  $z_{\alpha/2} = 1.96$ . So, the desired CI is

$$(101.15 - 1.96 \frac{105.4091}{\sqrt{20}}, 101.15 + 1.96 \frac{105.4091}{\sqrt{20}}) = \boxed{(54.95, 147.35)} \quad (6.35)$$

□

**4.2.6.**  $\bar{X}$  is the sample mean of a sample of size  $n$  from  $\mathcal{N}(\mu, 9)$ . Find  $n$  such that

$$P(\bar{X} - 1 \leq \mu \leq \bar{X} + 1) = 0.90 \quad (6.36)$$

*Solution:*  $\sigma^2 = 9 \implies \sigma = 3$ . We have

$$\begin{aligned} 0.90 &= P(\bar{X} - 1 \leq \mu \leq \bar{X} + 1) \\ &= P(\mu - 1 \leq \bar{X} \leq \mu + 1) \\ &= P(-1 \leq \bar{X} - \mu \leq 1) \\ &= P\left(\frac{-1}{3/\sqrt{n}} \leq \frac{\bar{X} - \mu}{3/\sqrt{n}} \leq \frac{1}{3/\sqrt{n}}\right). \end{aligned} \quad (6.37)$$

In other words,

$$z_{0.05} = \frac{1}{3/\sqrt{n}} = \frac{\sqrt{n}}{3} = 1.644854 \implies n = 24.35 \approx \boxed{25}. \quad (6.38)$$

□

**4.2.18.**  $X_i$ 's  $\sim \mathcal{N}(\mu, \sigma^2)$ , with  $\mu, \sigma^2$  unknown. A confidence interval for  $\sigma^2$  can be found as follows. We know that  $(n - 1)S^2/\sigma^2$  is a random variable with a  $\chi^2(n-1)$  distribution. Thus we can find constants  $a$  and  $b$  so that  $P((n - 1)S^2/\sigma^2 < b) = 0.975$  and  $P(a < (n - 1)S^2/\sigma^2 < b) = 0.95$ . In R,  $b = qchisq(0.975, n - 1)$ , while  $a = qchisq(0.025, n - 1)$ .

- (a) Show that this second probability statement can be written as

$$P((n - 1)S^2/b < \sigma^2 < (n - 1)S^2/a) = 0.95. \quad (6.39)$$

- (b) If  $n = 9$  and  $S^2 = 7.93$ , find a 95% confidence interval for  $\sigma^2$ .

- (c) If  $\mu$  is known, how would you modify the preceding procedure for finding a confidence interval for  $\sigma^2$ ?

*Solution:*

- (a) We simply re-arrange things in the probability statement:

$$\begin{aligned} 0.95 &= P(a < (n - 1)S^2/\sigma^2 < b) \\ &= P(\sigma^2 < (n - 1)S^2/a \wedge \sigma^2 > (n - 1)S^2/b) \\ &= P((n - 1)S^2/b < \sigma^2 < (n - 1)S^2/a). \end{aligned} \quad (6.40)$$

- (b) When  $n = 9, S^2 = 7.93$ , we have  $a = 2.179731$  and  $b = 17.53455$ . Then the 95% CI for  $\sigma^2$  is

$$\left( \frac{(n - 1)S^2}{b}, \frac{(n - 1)S^2}{a} \right) = \left( \frac{8 \times 7.93}{17.53455}, \frac{8 \times 7.93}{2.179731} \right) = \boxed{(3.618, 29.10451)} \quad (6.41)$$

- (c) If  $\mu$  is known, the unbiased estimator for the population standard deviation becomes proportional to  $1/\sqrt{n}$ , not  $1/\sqrt{n-1}$ . Because of this, we modify some numerics in our procedure from  $n - 1$  to  $n$ . From here, we make the following changes

$$\begin{aligned} (n - 1)S^2/\sigma^2 &\sim \chi^2(n - 1) \rightarrow nS^2/\sigma^2 \sim \chi^2(n) \\ P(nS^2/\sigma^2 < b) &= 0.975 \\ P(a < nS^2/\sigma^2 < b) &= 0.95. \end{aligned} \quad (6.42)$$

The new CI will look like  $(nS^2/b < \sigma^2 < nS^2/a)$ .

□

**4.2.21.** Let two independent random samples, each of size 10, from two normal distributions  $\mathcal{N}(\mu_1, \sigma^2)$  and  $\mathcal{N}(\mu_2, \sigma^2)$  yield  $\bar{x} = 4.8, s_1^2 = 8.64, \bar{y} = 5.6, s_2^2 = 7.88$ . Find a 95% confidence interval for  $\mu_1 - \mu_2$ .

*Solution:* The 95% CI for difference in means in this case looks like

$$\left( (\bar{x} - \bar{y}) - t_{\alpha/2, n_1 + n_2 - 2} s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}, (\bar{x} - \bar{y}) + t_{\alpha/2, n_1 + n_2 - 2} s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} \right). \quad (6.43)$$

The pooled variance is

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2} = \frac{9 \times 8.64 + 9 \times 7.88}{18} = 8.26. \quad (6.44)$$

Plugging in numbers we find the CI, with  $t_{0.025, 18} = 2.100922$ :

$$\begin{aligned} & \left( (4.8 - 5.6) - 2.100922 \sqrt{8.26} \sqrt{\frac{1}{10} + \frac{1}{10}}, (4.8 - 5.6) + 2.100922 \sqrt{8.26} \sqrt{\frac{1}{10} + \frac{1}{10}} \right) \\ &= \boxed{(-3.500, 1.900)} \end{aligned} \quad (6.45)$$

□

**4.2.22.** Let two independent random variables,  $Y_1$  and  $Y_2$ , with binomial distributions that have parameters  $n_1 = n_2 = 100$ ,  $p_1$ , and  $p_2$ , respectively, be observed to be equal to  $y_1 = 50$  and  $y_2 = 40$ . Determine an approximate 90% confidence interval for  $p_1 - p_2$ .

*Solution:* The 90% CI for the difference in proportions looks like

$$\begin{aligned} & \left( (\hat{p}_1 - \hat{p}_2) - z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}}, \right. \\ & \quad \left. (\hat{p}_1 - \hat{p}_2) + z_{\alpha/2} \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}} \right) \end{aligned} \quad (6.46)$$

where  $\alpha_{0.05} = 1.644854$ . Plugging in numbers, we find

$$\begin{aligned} & \left( 0.5 - 0.4 - 1.644854 \sqrt{\frac{(0.5)(0.5)}{100} + \frac{(0.4)(0.6)}{100}}, \right. \\ & \quad \left. 0.5 - 0.4 + 1.644854 \sqrt{\frac{(0.5)(0.5)}{100} + \frac{(0.4)(0.6)}{100}} \right) \\ & = \boxed{(-0.01513978, 0.2151398)} \end{aligned} \quad (6.47)$$

□

**4.2.27.** Let  $X_1, X_2, \dots, X_n$  and  $Y_1, Y_2, \dots, Y_m$  be two independent random samples from the respective normal distributions  $\mathcal{N}(\mu_1, \sigma_1^2)$  and  $\mathcal{N}(\mu_2, \sigma_2^2)$ , where the four parameters are unknown. To construct a confidence interval for the ratio,  $\sigma_1^2/\sigma_2^2$ , of the variances, form the quotient of the two independent 2 variables, each divided by its degrees of freedom, namely,

$$F = \frac{\frac{(m-1)S_2^2}{\sigma_2^2}/(m-1)}{\frac{(n-1)S_1^2}{\sigma_1^2}/(n-1)} = \frac{S_2^2/\sigma_2^2}{S_1^2/\sigma_1^2} \quad (6.48)$$

where  $S_1^2, S_2^2$  are respectively sample variances.

- (a) What kind of distribution does  $F$  have?
- (b) Rewrite the second probability statement as

$$P \left[ a \frac{S_1^2}{S_2^2} < \frac{\sigma_1^2}{\sigma_2^2} < b \frac{S_1^2}{S_2^2} \right] = 0.95. \quad (6.49)$$

The observed values,  $s_1^2$  and  $s_2^2$ , can be inserted in these inequalities to provide a 95% confidence interval for  $\sigma_1^2/\sigma_2^2$ .

*Solution:*

- (a)  $F \sim F(m-1, n-1)$ , by definition.
- (b) We just rearrange the quantities in the probability statement:

$$\begin{aligned} 0.95 &= P(a < F < b) \\ &= P \left( a < \frac{S_2^2/\sigma_2^2}{S_1^2/\sigma_1^2} < b \right) \\ &= P \left( \frac{\sigma_1^2}{\sigma_2^2} < b \frac{S_1^2}{S_2^2} \wedge \frac{\sigma_1^2}{\sigma_2^2} > a \frac{S_1^2}{S_2^2} \right) \\ &= P \left( a \frac{S_1^2}{S_2^2} < \frac{\sigma_1^2}{\sigma_2^2} < b \frac{S_1^2}{S_2^2} \right). \end{aligned} \quad (6.50)$$

□

**4.5.1.** Show that the approximate power function given in expression (4.5.12) of Example 4.5.3 is a strictly increasing function of  $\mu$ . Show then that the test discussed in this example has approximate size  $\alpha$  for testing

$$H_0 : \mu \leq \mu_0 \quad \text{versus} \quad H_1 : \mu > \mu_0. \quad (6.51)$$

*Solution:* The approximate power function is given by

$$\gamma(\mu) \approx \Phi\left(-z_\alpha - \frac{\sqrt{n}(\mu_0 - \mu)}{\sigma}\right) \quad (6.52)$$

$\partial_\mu \gamma(\mu)$  is necessarily positive  $\forall \mu \in \mathbb{R}$  for  $\gamma(\mu)$  to be strictly increasing. So we check:

$$\begin{aligned} \partial_\mu \gamma(\mu) &= \partial_\mu \left[ \Phi\left(-z_\alpha - \frac{\sqrt{n}(\mu_0 - \mu)}{\sigma}\right) \right] \\ &= \Phi'\left(-z_\alpha - \frac{\sqrt{n}(\mu_0 - \mu)}{\sigma}\right) \frac{\sqrt{n}}{\sigma}. \end{aligned} \quad (6.53)$$

We note that  $\sqrt{n}/\sigma > 0$  and  $\Phi'(\dots) > 0$  necessarily because  $\Phi$  is a cdf (for the  $\mathcal{N}(\mu, \sigma^2)$ ). With this, we have shown that  $\gamma(\mu)$  is strictly increasing in  $\mu$ .

Under the hypotheses and the fact that  $\gamma(\mu)$  is strictly increasing in  $\mu$ ,  $\alpha = \max_{\mu \leq \mu_0} \gamma(\mu)$  is maximized whenever  $\mu \leq \mu_0$  is maximized, i.e.  $\mu = \mu_0$ :

$$\max_{\mu \leq \mu_0} \gamma(\mu) = \gamma(\mu_0) = \Phi(-z_\alpha) = \alpha. \quad (6.54)$$

So the test has approximate size  $\alpha$ . □

**4.5.2.** For the Darwin data tabled in Example 4.5.5, verify that the Student t-test statistic is 2.15.

*Solution:*  $\alpha = 0.05$ . The sample mean and standard deviation for the differences are

$$\bar{x} = 2.62 \quad (6.55)$$

$$s_x = 4.71826. \quad (6.56)$$

The t-statistic is then

$$t_{df=14} = \frac{\bar{x} - 0}{s_x} = \frac{2.62}{4.71826/\sqrt{15}} \approx 2.150627 \quad (6.57)$$

**R code:**

```
> mean(darwin$cross)-mean(darwin$self)
[1] 2.62

> sd(darwin$cross - darwin$self)
[1] 4.71826
```

□

**4.5.5.** Let  $X_1, X_2$  be a random sample of size  $n = 2$  from the distribution having pdf  $f(x; \theta) = (1/\theta)e^{-x/\theta}$ ,  $\theta < x < \infty$ , zero elsewhere. We reject  $H_0 : \theta = 2$  and accept  $H_1 : \theta = 1$  if the observed values of  $X_1, X_2$ , say  $x_1, x_2$ , are such that

$$\frac{f(x_1; 2)f(x_2; 2)}{f(x_1; 1)f(x_2; 1)} \leq \frac{1}{2} \quad (6.58)$$

Here  $\Omega = \{\theta : \theta = 1, 2\}$ . Find the significance level of the test and the power of the test when  $H_0$  is false.

*Solution:* We reject  $H_0$  whenever

$$\begin{aligned} \frac{1}{2} &\geq \frac{f(x_1; 2)f(x_2; 2)}{f(x_1; 1)f(x_2; 1)} \\ &= \frac{(1/2)e^{-x_1/2}(1/2)e^{-x_2/2}}{e^{-x_1}e^{-x_2}} \\ &= \frac{1}{4}e^{x_1/2}e^{x_2/2} \\ &= \frac{1}{4}e^{(x_1+x_2)/2} \implies x_1 + x_2 \leq 2\ln(2). \end{aligned} \quad (6.59)$$

The significance level of the test  $\alpha$  is the probability of rejecting  $H_0$  when it is true, i.e.

$$\alpha = P(x_1 + x_2 \leq 2\ln(2) | \theta = 2). \quad (6.60)$$

Recall that the non-zero part of the pdf for  $\Gamma(k, \theta)$  is given by

$$\begin{aligned} f(x) &= \frac{1}{\Gamma(k)\theta^k} x^{k-1} e^{-x/\theta}, \quad x \in \mathbb{R}^+ \\ &= \frac{1}{\theta^k} e^{-x/\theta}, \quad k = 1, \end{aligned} \quad (6.61)$$

we have that  $X_1, X_2 \sim \Gamma(k = 1, \theta = 2)$ , iid, implies  $X_1 + X_2 \sim \Gamma(k = 2, \theta = 2)$ . From here, it is “easy” to calculate  $\alpha$ :

$$\begin{aligned} \alpha &= P(x_1 + x_2 \leq 2\ln(2) | \theta = 2) = \int_0^{2\ln(2)} \frac{1}{\Gamma(2)\theta^2} \xi e^{-\xi/\theta} d\xi \\ &= \int_0^{2\ln(2)} \frac{1}{4} \xi e^{-\xi/2} d\xi \\ &= \frac{1}{2}(1 - \ln(2)) \approx \boxed{0.1534} \end{aligned} \quad (6.62)$$

The power of the test is the probability of rejecting  $H_0$  when  $H_0$  is false. In this case, we make a similar calculation, only setting  $\theta = 1$  (since  $H_0$  false):

$$\begin{aligned} P(x_1 + x_2 \leq 2\ln(2) | \theta = 1) &= \int_0^{2\ln(2)} \xi e^{-\xi} d\xi \\ &= \frac{3}{4} - \frac{\ln(2)}{2} \approx \boxed{0.403426} \end{aligned} \quad (6.63)$$

□

**4.5.12.** Let  $X_1, X_2, \dots, X_8$  be a random sample of size  $n = 8$  from a Poisson distribution with mean  $\mu$ . Reject the simple null hypothesis  $H_0 : \mu = 0.5$  and accept  $H_1 : \mu > 0.5$  if the observed  $\sum_{i=1}^8 x_i \geq 8$ .

- (a) Show that the significance level is  $1 - \text{ppois}(7, 8 * .5)$ .
- (b) Use R to determine  $\gamma(0.75)$ ,  $\gamma(1)$ , and  $\gamma(1.25)$ .
- (c) Modify the code in Exercise 4.5.9 to obtain a plot of the power function.

*Solution:*

- (a) The significance level  $\alpha$  is the probability of rejecting  $H_0$  when  $H_0$  is true. Under the null,  $\mu = 0.5$ , and the r.v.

$$X_1 + \dots + X_8 \sim \text{Poi}(8\mu) \equiv \text{Poi}(8 \times 0.5). \quad (6.64)$$

$\alpha$  is given by

$$\begin{aligned} \alpha = \gamma(\mu) &= P\left(\sum_{i=1}^8 x_i \geq 8 | \mu = 0.5\right) \\ &= 1 - P\left(\sum_{i=1}^8 x_i < 7\right) \\ &= \boxed{1 - \text{ppois}(7, 8 * .5)} \\ &= 0.05113362 \end{aligned} \quad (6.65)$$

(b)

$$\begin{aligned} \gamma(0.75) &= \boxed{0.2560202} \\ \gamma(1) &= \boxed{0.5470392} \\ \gamma(1.25) &= \boxed{0.7797794} \end{aligned} \quad (6.66)$$

**R code:**

```
> 1 - ppois(7, 8 * 0.5)
[1] 0.05113362

> 1 - ppois(7, 8 * 0.75)
[1] 0.2560202

> 1 - ppois(7, 8 * 1)
[1] 0.5470392


> 1 - ppois(7, 8 * 1.25)
[1] 0.7797794
```

(c) **R code:**

```
> theta=seq(.75,1.25,.25); gam=1-ppois(7,theta*8)
> plot(gam~theta,pch=" ",xlab=expression(theta),ylab=expression(gamma))
> lines(gam~theta)
```

We're interested in the range of  $\theta \in [0.75, 1.25]$ . I'm making the step size small to make the power function look smooth.

Plot of  $\gamma(\mu)$ :



□

**4.6.4.** (Note that it is fine to make a heuristic argument here. Just make sure that it is clear with supporting graphs/figures (hand drawn is fine).) Consider the one-sided t-test for  $H_0 : \mu = \mu_0$  versus  $H_{A1} : \mu > \mu_0$  constructed in Example 4.5.4 and the two-sided t-test for t-test for  $H_0 : \mu = \mu_0$  versus  $H_1 : \mu = \mu_0$  given in (4.6.9). Assume that both tests are of size  $\alpha$ . Show that for  $\mu > \mu_0$ , the power function of the one-sided test is larger than the power function of the two-sided test.

*Solution:* We want to show that for  $\mu > \mu_0$ , the power function of the one-sided test is larger than the power function of the two-sided test. To this end, let  $\gamma_1(\mu)$  denote the power function of the one-sided test, and  $\gamma_2(\mu)$  the two-sided test. This gives

$$\gamma_2(\mu) = P(|\text{test-statistic}| \geq t_{\alpha/2,n-1}) = P(\text{test-statistic} \geq t_{\alpha/2,n-1}) \quad (6.67)$$

(test statistic positive because  $\mu > \mu_0$ ), while

$$\gamma_1(\mu) = P(\text{test-statistic} \geq t_{\alpha,n-1}). \quad (6.68)$$

Since  $t_{\alpha/2,n-1} > t_{\alpha,n-1}$ , we have that

$$\gamma_2(\mu) = P(|\cdot| \geq t_{\alpha/2}) \leq P(\cdot \geq t_{\alpha}) = \gamma_1(\mu). \quad (6.69)$$

And so, for a given  $\mu > \mu_0$ , the power function of the one-sided test is larger than the power function of the two-sided test.  $\square$

**4.6.7.** Among the data collected for the World Health Organization air quality monitoring project is a measure of suspended particles in  $\mu\text{g}/\text{m}^3$ . Let  $X$  and  $Y$  equal the concentration of suspended particles in  $\mu\text{g}/\text{m}^3$  in the city center (commercial district) for Melbourne and Houston, respectively. Using  $n = 13$  observations of  $X$  and  $m = 16$  observations of  $Y$ , we test  $H_0 : \mu_X = \mu_Y$  against  $H_1 : \mu_X < \mu_Y$ .

- (a) Define the test statistic and critical region, assuming that the unknown variances are equal. Let  $\alpha = 0.05$ .
- (b) If  $\bar{x} = 72.9$ ,  $s_x = 25.6$ ,  $\bar{y} = 81.7$ , and  $s_y = 28.3$ , calculate the value of the test statistic and state your conclusion.

*Solution:*

- (a) Assuming the unknown variances are equal, we have

$$\tau = \frac{(\bar{y} - \bar{x})}{s_p \sqrt{\frac{1}{13} + \frac{1}{16}}} \quad (6.70)$$

The critical region is given by

$$C := \{(X_1, \dots, X_{13}, Y_1, \dots, Y_{16}) | \tau \geq t_{0.05, 13+16-2} = [1.703288]\}. \quad (6.71)$$

- (b) With the given numbers, we calculate the pooled variance is

$$S_p^2 = \frac{(13-1)(25.6)^2 + (16-1)(28.3)^2}{13+16-2} = 736.21 \quad (6.72)$$

With this,

$$\tau = \frac{(81.7 - 72.9) - 0}{\sqrt{736.21} \sqrt{\frac{1}{13} + \frac{1}{16}}} = [0.8685893] \quad (6.73)$$

Since  $0.8685893 < 1.703288$ , there is not enough evidence to reject  $H_0$  in favor of  $H_a$ .

□

**4.6.8.** Let  $p$  equal the proportion of drivers who use a seat belt in a country that does not have a mandatory seat belt law. It was claimed that  $p = 0.14$ . An advertising campaign was conducted to increase this proportion. Two months after the campaign,  $y = 104$  out of a random sample of  $n = 590$  drivers were wearing their seat belts. Was the campaign successful?

1. Define the null and alternative hypotheses.
2. Define a critical region with an  $\alpha = 0.01$  significance level.
3. Determine the approximate  $p$ -value and state your conclusion.

*Solution:*

(a)  $H_0 : p = 0.14 \quad H_a : p > 0.14$ .

(b) The critical region is given by

$$C := \left\{ y \left| \frac{y/n - 0.14}{\sqrt{\frac{0.14(1-0.14)}{590}}} \geq z_{\alpha=0.01} = 2.326348 \right. \right\}. \quad (6.74)$$

(c) The value of the test statistic is

$$z^* = \frac{104/590 - 0.14}{\sqrt{\frac{0.14(1-0.14)}{590}}} = \boxed{2.539069 > 2.326348} \quad (6.75)$$

Since  $z^* > z$ , there is enough evidence to reject  $H_0$  in favor of  $H_a$  ( $p$ -value:  $\boxed{0.006 < 0.01 = \alpha}$ ), i.e., there is enough evidence to suggest that the campaign was successful.

**R code:**

```
> 1-pnorm(2.539069)
[1] 0.005557395
```

□

### 6.3 Problem set 3

#### 4.7.4

A die was cast  $n = 120$  independent times and the following data resulted:

Spots Up	1	2	3	4	5	6
Frequency	$b$	20	20	20	20	$40 - b$

If we use a chi-square test, for what values of  $b$  would the hypothesis that the die is unbiased be rejected at the 0.025 significance level?

*Solution:* The test statistic under the null hypothesis  $H_0 : \partial_i = 1/6 \forall i$  is

$$\chi = \sum_{i=1}^6 \frac{(\text{Freq}_i - 20)^2}{20} = \frac{(b - 20)^2}{20} + 4 \cdot 0 + \frac{(40 - b - 20)^2}{20} = \frac{(b - 20)^2}{10}. \quad (6.76)$$

Under the null hypothesis,  $\chi \sim \chi^2(df = 5)$ . At  $\alpha = 0.025$ , we reject the null hypothesis whenever  $\chi \geq \text{qchisq}(1-0.025, 5) = 12.8325$ , i.e.,

$$\frac{(b - 20)^2}{10} \geq 12.8325 \implies [b \leq 8 \vee b \geq 32] \quad (6.77)$$

□

**4.7.5**

Consider the problem from genetics of crossing two types of peas. The Mendelian theory states that the probabilities of the classifications (a) round and yellow, (b) wrinkled and yellow, (c) round and green, and (d) wrinkled and green are  $\frac{9}{16}$ ,  $\frac{3}{16}$ ,  $\frac{3}{16}$ , and  $\frac{1}{16}$ , respectively. If, from 160 independent observations, the observed frequencies of these respective classifications are 86, 35, 26, and 13, are these data consistent with the Mendelian theory? That is, test, with  $\alpha = 0.01$ , the hypothesis that the respective probabilities are  $\frac{9}{16}$ ,  $\frac{3}{16}$ ,  $\frac{3}{16}$ , and  $\frac{1}{16}$ .

*Solution:* The test statistic under  $H_0 : p_a = 9/16, p_b = 3/16, p_c = 3/16, p_d = 1/16$ , the test statistic is

$$\chi^2 = \frac{(86 - 90)^2}{90} + \frac{(35 - 30)^2}{30} + \frac{(26 - 30)^2}{30} + \frac{(13 - 10)^2}{10} = \frac{22}{9}. \quad (6.78)$$

Under  $H_0$ ,  $\chi^2 \sim \chi^2(df = 4-1 = 3)$ . The p-value for this  $\chi^2$  is  $1 - \text{pchisq}(22/9, 3) = 0.4854149 > 0.01$ . So, we don't have enough evidence to reject  $H_0$ , i.e. the data is consistent with the Mendelian theory.  $\square$

**4.7.6**

Two different teaching procedures were used on two different groups of students. Each group contained 100 students of about the same ability. At the end of the term, an evaluating team assigned a letter grade to each student. The results were tabulated as follows.

Group	Grade					Total
	A	B	C	D	F	
I	15	25	32	17	11	100
II	9	18	29	28	16	100

If we consider these data to be independent observations from two respective multinomial distributions with  $k = 5$ , test at the 5% significance level the hypothesis that the two distributions are the same (and hence the two teaching procedures are equally effective). For computation in R, use

```
r1=c(15,25,32,17,11);r2=c(9,18,29,28,16);mat=rbind(r1,r2)
chisq.test(mat)
```

*Solution:* Using the code provided by the problem we get

```
> r1=c(15,25,32,17,11);r2=c(9,18,29,28,16);mat=rbind(r1,r2)
> chisq.test(mat)

Pearson's Chi-squared test

data: mat
X-squared = 6.4019, df = 4, p-value = 0.1711
```

Since  $p = 0.1711 > 0.05$ , we don't have enough evidence to reject  $H_0$ , i.e. the two teaching procedures are (statistically) equally effective.  $\square$

**4.8.9** (use acceptance sampling)

Determine a method to generate random observations for the Cauchy distribution with pdf

$$f(x) = \frac{1}{\pi(1+x^2)}, \quad -\infty < x < \infty. \quad (4.8.16)$$

Write an R function that returns a random sample of observations from this Cauchy distribution.

*Solution: R code:*

```
# acceptance sampling
# approximate the Cauchy distribution

x <- runif(100000,-7,7)
y <- runif(100000,0,1)

z <- cbind(x,y)
accept <- NULL
reject <- NULL

dens <- function(x){
  d <- 1/((pi^2)*(x^2+1))
  return(d)}


for (i in 1:length(x)){
  d <- dens(x[i])
  if (y[i] < d) {
    accept <- rbind(accept,z[i,])}
  else{
    reject <- rbind(reject,z[i,])}
}

# plot accepted values in red
plot(accept[,1],accept[,2],
main = "Approximated Distribution", ylim=c(0,1), col="red")

# plot rejected values in blue
points(reject[,1], reject[,2], col = "dark blue")
```

Here's a small sample:

```
> accept
x           y
[1,] -0.2257702402 8.676955e-02
[2,] -0.3536258731 6.871887e-02
[3,]  4.1498403600 4.068073e-03
[4,] -0.3986396971 3.033239e-02
[5,]  0.7077048477 1.492551e-02
[6,] -0.0422784868 1.612575e-02
[7,] -0.6391186416 1.029603e-02
[8,]  0.4845524384 6.119987e-02
[9,]  0.3572320105 2.479371e-02
.
.
```



□

**4.8.10** (use inverse transformation sampling)

Suppose we are interested in a particular Weibull distribution with pdf

$$f(x) = \begin{cases} \frac{1}{\theta^3} 3x^2 e^{-x^3/\theta^3} & 0 < x < \infty \\ 0 & \text{elsewhere.} \end{cases}$$

Determine a method to generate random observations from this Weibull distribution. Write an R function that returns such a sample.

*Hint:* Find  $F^{-1}(u)$ .

*Solution:* We first want to find the cdf  $F$ , given  $f(x)$ . Well,

$$F(x) = \int_{-\infty}^x f(x') dx' = \int_0^x \frac{1}{\theta^3} 3x'^2 e^{-x'^3/\theta^3} dx' = \dots = 1 - e^{-x^3/\theta^3}. \quad (6.79)$$

With this, let  $u \sim U(0, 1)$  then

$$F^{-1}(u) = \sqrt[3]{-\theta^3 \ln(1-u)} \sim \text{Wei}(k=3, \theta), \quad 0 \leq u \leq 1. \quad (6.80)$$

Suppose  $\theta = 2$  then in R, we do the following:



```
# inverse transformation sampling
# generate vector of random uniform(0,1)
u <- runif(100000)

# set beta and transform to random weibull(shape=3, scale=theta)
beta = 2
x <- (-(beta^3)*log(1 - u))^(1/3)
hist(x, main="Histogram of Tranformed Variable", pr = TRUE)

# overlay weibull
curve(dweibull(x, shape = 3, scale=2),
col="darkgreen", lwd=3, add=TRUE)
```

Here's a sample:

```
> x <- (-(beta^3)*log(1 - u))^(1/3)
> x
1.57511912 1.81610458 2.51359811 1.12564398 1.21991006
1.10365173 0.71966428 2.06591082 3.08423326 1.67620145
1.02533405 1.24429387 0.94192313 2.19097666 1.98069305
0.43007645 1.18581709 1.11303036 1.63002049 2.37205273
2.45962833 1.67162606 1.61624974 1.46149955 1.92907758
1.12735030 1.02997348 2.25870529 1.58137215 2.71199084
1.05306668 1.43121600 1.17977939 1.24985669 2.22292333
1.86914252 3.01206737 1.11948037 1.64757666 1.35904188
1.39351092 2.60002179 1.12125772 2.15576011 2.41963279
```

**Histogram of Tranformed Variable, beta=0.5****Histogram of Tranformed Variable, beta=2**

□

## 6.4 Problem set 4

**4.9.4** (for part B, write your own R function to generate the bootstrap distribution of the median - using the code posted on Moodle is an okay starting point)

- Consider the situation discussed in Example 4.9.1. Suppose we want to estimate the median of  $X_i$  using the sample median.

- (a) Determine the median for a  $\Gamma(1, \beta)$  distribution.
- (b) The algorithm for the bootstrap percentile confidence intervals is general and hence can be used for the median. Rewrite the R code in the function `percentciboot.s` so that the median is the estimator. Using the sample given in the example, obtain a 90% bootstrap percentile confidence interval for the median. Did it trap the true median in this case?

*Solution:*

- (a) The median  $x_{1/2}$  for a  $\Gamma(1, \beta)$  is such that

$$\frac{1}{2} = \int_0^{x_{1/2}} \frac{1}{\beta} e^{-x/\beta} = 1 - e^{-x_{1/2}/\beta} \implies \boxed{x_{1/2} = \beta \ln(2)} \quad (6.81)$$

- (b) Here's the R code (I'm not using Prof. O'Brien's code here)

```
> x <- c(131.7, 182.7, 73.3, 10.7, 150.4, 42.3, 22.2, 17.9, 264.0,
154.4, 4.3, 265.6, 61.9, 10.8, 48.8, 22.5, 8.8, 150.6, 103.0, 85.9)
> percentciboot <- function(x,b,alpha){
+   theta=median(x); thetastar=rep(0,b); n=length(x)
+   for(i in 1:b){xstar=sample(x,n,replace=T)
+     thetastar[i]=median(xstar)}
+   thetastar=sort(thetastar); pick=round((alpha/2)*(b+1))
+   lower=thetastar[pick]; upper=thetastar[b-pick+1]
+   list(theta=theta,lower=lower,upper=upper)}
> percentciboot(x,3000,.10)

$theta
[1] 67.6

$lower
[1] 30.1

$upper
[1] 131.7

> median(x)
[1] 67.6

> 100*log(2)
[1] 69.31472
```

The 90% bootstrap percentile CI for the median is given by  $(30.1, 131.7)$ . The true median is given by  $\beta \ln(2) = 100 \ln(2) \approx 69.31472$ . So, yes, the 90% CI traps the true median in this case.

□

**4.9.11**

Let  $z^*$  be drawn at random from the discrete distribution that has mass  $n^{-1}$  at each point  $z_i = x_i - \bar{x} + \mu_0$ , where  $(x_1, x_2, \dots, x_n)$  is the realization of a random sample. Determine  $E(z^*)$  and  $V(z^*)$ .

*Solution:*

(a)  $E(z^*)$ :

$$E(z^*) = \sum_{i=1}^n \frac{x_i - \bar{x} + \mu_0}{n} = \bar{x} - \bar{x} + \mu_0 = \boxed{\mu_0} \quad (6.82)$$

(b)  $V(z^*)$ :

$$V(z^*) = \sum_{i=1}^n (z_i - E(z^*))^2 = \sum_{i=1}^n (x_i - \bar{x} + \mu_0 - \mu_0)^2 = \boxed{\sum_{i=1}^n (x_i - \bar{x})^2} \quad (6.83)$$

□

### 4.9.13

For the situation described in Example 4.9.3, obtain the bootstrap test based on medians. Use the same hypotheses; i.e.,

$$H_0 : \mu = 90 \text{ versus } H_1 : \mu > 90.$$

*Solution:* Here's the R code:

```
> X <- c(119.7, 104.1, 92.8, 85.4, 108.6, 93.4, 67.1, 88.4, 101.0, 97.2,
+      95.4, 77.2, 100.0, 114.2, 150.3, 102.3, 105.8, 107.5, 0.9, 94.1)
> boottestonemed <-
+   function(x,theta0,b){
+     #
+     # x = sample
+     # theta0 is the null value of the mean
+     # b is the number of bootstrap resamples
+     #
+     # origtest contains the value of the test statistics
+     # for the original sample
+     # pvalue is the bootstrap p-value
+     # teststatall contains the b bootstrap tests
+     #
+     n<-length(x)
+     v<-median(x)
+     z<-x-median(x)+theta0
+     counter<-0
+     teststatall<-rep(0,b)
+     for(i in 1:b){xstar<-sample(z,n,replace=T)
+     vstar<-median(xstar)
+     if(vstar >= v){counter<-counter+1}
+     teststatall[i]<-vstar}
+     pvalue<-counter/b
+     list(origtest=v,pvalue=pvalue,teststatall=teststatall)
+     #list(origtest=v,pvalue=pvalue)
+   }
> boottestonemed(X,90,3000)
$origtest
[1] 98.6

$pvalue
[1] 0.006
```

At such a low p-value ( $p = 0.006$ ), we reject the null hypothesis  $H_0$ . So even though we don't reject  $H_0$  with the test based on sampling mean, we *do* reject  $H_0$  with the test based on medians.  $\square$

## 5.1.7

**5.1.7.** Let  $X_1, \dots, X_n$  be iid random variables with common pdf

$$f(x) = \begin{cases} e^{-(x-\theta)} & x > \theta, -\infty < \theta < \infty \\ 0 & \text{elsewhere.} \end{cases} \quad (5.1.3)$$

This pdf is called the **shifted exponential**. Let  $Y_n = \min\{X_1, \dots, X_n\}$ . Prove that  $Y_n \rightarrow \theta$  in probability by first obtaining the cdf of  $Y_n$ .

*Solution:* By definition,

$$\begin{aligned} Y_n \xrightarrow{P} \theta &\iff \forall \epsilon > 0, \lim_{n \rightarrow \infty} P[|Y_n - \theta| \geq \epsilon] = 0 \\ &\iff \forall \epsilon > 0, \lim_{n \rightarrow \infty} P[|\min\{X_1, \dots, X_n\} - \theta| \geq \epsilon] = 0 \\ &\iff \forall \epsilon > 0, \lim_{n \rightarrow \infty} P[\min\{X_1, \dots, X_n\} - \theta \geq \epsilon] = 0 \\ &\iff \forall \epsilon > 0, \lim_{n \rightarrow \infty} P[Y_n \geq \epsilon + \theta] = 0 \end{aligned} \quad (6.84)$$

where the second to last equivalence statement comes from the fact that  $x_i > \theta \forall i = 1, 2, \dots, n$ . To show  $Y_n \xrightarrow{P} \theta$ , we find the cdf for  $Y_n$ :

$$\begin{aligned} F_{Y_n}(y) &= P(Y_n < y) \\ &= 1 - P(Y_n \geq y) \\ &= 1 - \prod_{i=1}^n P(X_i \geq y) \\ &= 1 - \prod_{i=1}^n \int_y^\infty e^{-(x-\theta)} dx \\ &= 1 - \prod_{i=1}^n e^{-(y-\theta)} \\ &= 1 - e^{-n(y-\theta)} \\ P(Y_n \geq y) &= e^{-n(y-\theta)}. \end{aligned} \quad (6.85)$$

Let  $\epsilon > 0$  be given. Then

$$P(Y_n \geq \epsilon + \theta) = e^{-n(\epsilon+\theta-\theta)} = e^{-n\epsilon}, \quad (6.86)$$

and so

$$\lim_{n \rightarrow \infty} P[|Y_n - \theta| \geq \epsilon] = \lim_{n \rightarrow \infty} P[Y_n \geq \epsilon + \theta] = \lim_{n \rightarrow \infty} e^{-n\epsilon} = 0 \implies Y_n \xrightarrow{P} \theta. \quad (6.87)$$

□

**5.2.2** (Investigate = find)

**5.2.2.** Let  $Y_1$  denote the minimum of a random sample of size  $n$  from a distribution that has pdf  $f(x) = e^{-(x-\theta)}$ ,  $\theta < x < \infty$ , zero elsewhere. Let  $Z_n = n(Y_1 - \theta)$ . Investigate the limiting distribution of  $Z_n$ .

*Solution:* Let  $F_{Z_n}$  and  $F_Z$  be, respectively, the cdfs of  $Z_n$  and  $Z$ . Then by definition

$$Z_n \xrightarrow{D} Z \iff \lim_{n \rightarrow \infty} F_{Z_n}(z) = F_Z(z) \quad (6.88)$$

for all  $z$  at which  $F_Z(z)$  is continuous. Now, we don't know what  $F_Z$  is in this case, but we can find what  $F_{Z_n}$  converges to when  $n \rightarrow \infty$ . To show this, we find  $F_{Z_n}$ :

$$\begin{aligned} F_{Z_n}(z) &= P(Z_n \leq z) \\ &= P(n(Y_1 - \theta) \leq z) \\ &= P(Y_1 \leq z/n + \theta) \\ &= 1 - e^{-n(z/n + \theta - \theta)}, \quad \text{calculated in Problem 5.1.7} \\ &= 1 - e^{-z}. \end{aligned} \quad (6.89)$$

And so (obviously)

$$\lim_{n \rightarrow \infty} F_{Z_n}(z) = 1 - e^{-z} \equiv \text{cdf}(\text{Exp}(1)) \quad (6.90)$$

Therefore,  $Z_n \xrightarrow{D} Z \sim \text{Exp}(1)$ . □

### 5.2.7

**5.2.7.** Let  $X_n$  have a gamma distribution with parameter  $\alpha = n$  and  $\beta$ , where  $\beta$  is not a function of  $n$ . Let  $Y_n = X_n/n$ . Find the limiting distribution of  $Y_n$ .

*Solution:* Let  $X_n \sim \Gamma(\alpha = n, \beta)$  be given, where  $\beta$  is not a function of  $n$ . Let  $Y_n = X_n/n$ . To find the limiting distribution of  $Y_n$ , we find  $\lim_{n \rightarrow \infty} M_{Y_n}(t)$ . The reason we don't want to find the cdf  $F_{Y_n}$  is that integrals involving the Gamma distribution are often ugly.

$$\begin{aligned} M_{Y_n}(t) &= E[e^{tY_n}] = E[e^{tX_n/n}] \equiv E[e^{t_n X_n}] \\ &= (1 - \beta t_n)^{-n} \\ &= \left(1 - \frac{\beta t}{n}\right)^{-n}. \end{aligned} \quad (6.91)$$

And so using the identity

$$\lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^{-n} = e, \quad (6.92)$$

by change of variables we obtain

$$\lim_{n \rightarrow \infty} M_{Y_n}(t) = \lim_{n \rightarrow \infty} \left(1 - \frac{\beta t}{n}\right)^{-n} = e^{\beta t} \quad (6.93)$$

So what is the limiting distribution of  $Y_n$ ? By definition,

$$M_Y(t) = E[e^{tY}] = \int_{-\infty}^{\infty} f_Y(y) e^{yt} dy = e^{\beta t}. \quad (6.94)$$

Upon inspection, this equality holds if and only if  $f_Y(y) \equiv \delta(y - \beta)$ , the delta function centered at  $\beta$ .

$$\int_{-\infty}^{\infty} \delta(y - \beta) e^{yt} dy = e^{\beta t}. \quad (6.95)$$

And so, the limiting distribution of  $Y_n$  is the degenerate distribution with parameter  $\beta$ :

$$Y_n \xrightarrow{D} Y \sim \delta(y - \beta) \equiv \begin{cases} 1, & y = \beta \\ 0, & \text{else} \end{cases}. \quad (6.96)$$

□

**5.3.9**

**5.3.9.** Let  $f(x) = 1/x^2$ ,  $1 < x < \infty$ , zero elsewhere, be the pdf of a random variable  $X$ . Consider a random sample of size 72 from the distribution having this pdf. Compute approximately the probability that more than 50 of the observations of the random sample are less than 3.

*Solution:* We have  $X_1, \dots, X_{72}$ , with

$$X_i \sim f(x) = \begin{cases} \frac{1}{x^2}, & 1 < x < \infty \\ 0, & \text{else} \end{cases}. \quad (6.97)$$

We first find the probability that any given observation is less than 3:

$$P(X < 3) = \int_1^3 \frac{1}{x^2} dx = \frac{2}{3}. \quad (6.98)$$

So we have a “binomial situation” where the probability of success is  $p = 2/3$ . Given  $n = 72$  trials, we have  $\mu = np = 72(2/3) = 48$  and  $\sigma = \sqrt{np(1-p)} = 4$ . We wish to find the probability of having more than 50 successes. To this end, we use the normal approximation (CLT), which says

$$\frac{Y_{72} - \mu}{\sigma} \sim \mathcal{N}(0, 1). \quad (6.99)$$

And so

$$\begin{aligned} P(Y_{72} > 50) &\approx P\left(Z \geq \frac{51 - 48}{4}\right) \\ &= 1 - \text{pnorm}(3/4) \\ &= \boxed{0.2266274} \end{aligned} \quad (6.100)$$

Or we can also use the continuity correction to get

$$\begin{aligned} P(Y_{72} > 50) &\approx P\left(Z \geq \frac{50.5 - 48}{4}\right) \\ &= 1 - \text{pnorm}((50.5 - 48)/4) \\ &= \boxed{0.2659855} \end{aligned} \quad (6.101)$$

□

**5.3.11**

**5.3.11.** We know that  $\bar{X}$  is approximately  $N(\mu, \sigma^2/n)$  for large  $n$ . Find the approximate distribution of  $u(\bar{X}) = \bar{X}^3$ , provided that  $\mu \neq 0$ .

*Solution:* We want to use the  $\Delta$ -method for this problem. Since  $\bar{X} \sim N(\mu, \sigma^2/n)$ , we have

$$\sqrt{n}(\bar{X} - \mu) \xrightarrow{D} N(0, \sigma^2), \quad (6.102)$$

by a simple change of variables. The function  $u(\bar{X}) = \bar{X}^3$  is differential at for all  $\bar{X}$  and  $u'(\bar{X}) \neq 0$  in general, so by the  $\Delta$ -method,

$$\begin{aligned} & \sqrt{n}(u(\bar{X}) - u(\mu)) \xrightarrow{D} N(0, \sigma^2(u'(\mu))^2) \\ \iff & \sqrt{n}(u(\bar{X}) - u(\mu)) \xrightarrow{D} N(0, \sigma^2(3\mu^2)^2) \\ \iff & \sqrt{n}(u(\bar{X}) - \mu^3) \xrightarrow{D} N(0, 9\sigma^2\mu^4). \end{aligned} \quad (6.103)$$

But of course, the convergence in distribution above is equivalent to

$$u(\bar{X}) = \bar{X}^3 \xrightarrow{D} N\left(\mu^3, \frac{9\sigma^2\mu^4}{n}\right), \quad (6.104)$$

again by change of variables.  $\square$

## 6.5 Problem set 5

### 6.1.1

**6.1.1.** Let  $X_1, X_2, \dots, X_n$  be a random sample on  $X$  that has a  $\Gamma(\alpha = 4, \beta = \theta)$  distribution,  $0 < \theta < \infty$ .

- (a) Determine the mle of  $\theta$ .
- (b) Suppose the following data is a realization (rounded) of a random sample on  $X$ . Obtain a histogram with the argument `pr=T` (data are in `ex6111.rda`).

```
9 39 38 23 8 47 21 22 18 10 17 22 14
9 5 26 11 31 15 25 9 29 28 19 8
```

- (c) For this sample, obtain  $\hat{\theta}$  the realized value of the mle and locate  $4\hat{\theta}$  on the histogram. Overlay the  $\Gamma(\alpha = 4, \beta = \hat{\theta})$  pdf on the histogram. Does the data agree with this pdf? Code for overlay:

```
xs=sort(x);y=dgamma(xs,4,1/betahat);hist(x,pr=T);lines(y~xs).
```

*Solution:*

- (a) The likelihood function is

$$\mathcal{L}(\theta) = \prod_{i=1}^n \frac{1}{\Gamma(4)\theta^4} x_i^{4-1} e^{-\theta x_i} = \frac{1}{\Gamma(4)^n} \frac{1}{\theta^{4n}} e^{-\sum_{i=1}^n x_i/\theta} \prod_{i=1}^n x_i^3. \quad (6.105)$$

The log likelihood function is then

$$l(\theta) = -n \ln(\Gamma(4)) - 4n \ln \theta - \frac{1}{\theta} \sum_{i=1}^n x_i + \ln \left( \prod_{i=1}^n x_i^3 \right). \quad (6.106)$$

Next, solve for  $\partial_\theta l(\theta) = 0$ :

$$\partial_\theta l(\theta) = -\frac{4n}{\theta} + \frac{1}{\theta^2} \sum_{i=1}^n x_i = 0 \iff \boxed{\hat{\theta}_{ML} = \frac{1}{4n} \sum_{i=1}^n x_i = \bar{x}} \quad (6.107)$$

- (b) **R code:**

```
> dat = c(9, 39, 38, 23, 8, 47, 21, 22, 18, 10, 17, 22, 14,
+       9, 5, 26, 11, 31, 15, 25, 9, 29, 28, 19, 8)
> hist(dat, pr=T)
```




Figure 6.1: (b)


(c)  $\hat{\theta} = \bar{x}/4 = 5.03$  **R code:**

```
> mean(dat)/4
[1] 5.03

> xs=sort(dat)
> y=dgamma(xs,4,4/mean(dat))
> hist(dat,pr=T)
> lines(y~xs)
```

Locate  $4\hat{\theta}$  on the histogram and overlay with  $\Gamma(\alpha = 4, \beta = \hat{\theta} = 5.03)$ :

```
> xs=sort(dat)
> y=dgamma(xs,4,4/mean(dat))
> hist(dat,pr=T)
> lines(y~xs)
> abline(v=mean(dat))
```



The data somewhat agrees with this pdf.

□

### 6.1.2

**6.1.2.** Let  $X_1, X_2, \dots, X_n$  represent a random sample from each of the distributions having the following pdfs:

- (a)  $f(x; \theta) = \theta x^{\theta-1}$ ,  $0 < x < 1$ ,  $0 < \theta < \infty$ , zero elsewhere.
- (b)  $f(x; \theta) = e^{-(x-\theta)}$ ,  $\theta \leq x < \infty$ ,  $-\infty < \theta < \infty$ , zero elsewhere. Note that this is a nonregular case.

In each case find the mle  $\hat{\theta}$  of  $\theta$ .

*Solution:*

- (a) The likelihood function is

$$\mathcal{L}(\theta) = \theta^n \prod_{i=1}^n x_i^{\theta-1}, \quad x_i \in (0, 1), 0 < \theta < \infty. \quad (6.108)$$

The log likelihood is

$$l(\theta) = \ln \mathcal{L}(\theta) = n \ln \theta + (\theta - 1) \sum_{i=1}^n \ln x_i. \quad (6.109)$$

Then we solve for  $\theta$  in  $\partial_\theta l(\theta) = 0$ :

$$\partial_\theta l(\theta) = \frac{n}{\theta} + \sum_{i=1}^n \ln(x_i) = 0 \implies \boxed{\hat{\theta} = \frac{-n}{\sum_{i=1}^n \ln x_i}} \quad (6.110)$$

- (b) The likelihood function is

$$\mathcal{L}(\theta) = e^{-\sum_{i=1}^n (x_i - \theta)}, \quad \theta < x_i < \infty, -\infty < \theta < \infty. \quad (6.111)$$

Then the log likelihood is

$$l(\theta) = \ln \mathcal{L}(\theta) = -\sum_{i=1}^n x_i + n\theta. \quad (6.112)$$

Next,

$$\partial_\theta l(\theta) = n > 0. \quad (6.113)$$

So,  $\hat{\theta}$  must be as large as possible to maximize  $\mathcal{L}(\theta)$ . But at the same time,  $\theta \leq x_i$  for all  $i$ , so

$$\boxed{\hat{\theta} = \min_i(X_i)} \quad (6.114)$$

□

**6.1.4**

**6.1.4.** Suppose  $X_1, \dots, X_n$  are iid with pdf  $f(x; \theta) = 2x/\theta^2$ ,  $0 < x \leq \theta$ , zero elsewhere. Note this is a nonregular case. Find:

- (a) The mle  $\hat{\theta}$  for  $\theta$ .
- (b) The constant  $c$  so that  $E(c\hat{\theta}) = \theta$ .
- (c) The mle for the median of the distribution. Show that it is a consistent estimator.

*Solution:*

- (a) The likelihood function is

$$\mathcal{L}(\theta) = \frac{2^n}{\theta^{2n}} \prod_{i=1}^n x_i, \quad 0 < x_i \leq \theta. \quad (6.115)$$

The log likelihood is then

$$l(\theta) = n \ln 2 - 2n \ln \theta + \ln \left( \prod_{i=1}^n x_i \right). \quad (6.116)$$

Next,

$$\partial_\theta l(\theta) = -\frac{2n}{\theta}. \quad (6.117)$$

We cannot set this to zero. However, by inspection,  $\mathcal{L}(\theta)$  is maximized whenever  $\theta$  is minimized while  $x_i \leq \theta$  for all  $i$ , so

$$\boxed{\hat{\theta} = \max(X_i) = Y_n} \quad (6.118)$$

- (b) To find  $c$  we first find  $E(\hat{\theta})$ . To get this, we must first find its cdf.

$$F_{Y_n}(x) = P(Y_n \leq x) = \prod_{i=1}^n P(x_i \leq x) = \prod_{i=1}^n F_X(x) = \frac{x^{2n}}{\theta^{2n}}. \quad (6.119)$$

Differentiating this w.r.t.  $x$  we get the pdf of  $Y_n$ :

$$f_{Y_n}(x) = \partial_x F_{Y_n}(x) = 2n \frac{x^{2n-1}}{\theta^{2n}}. \quad (6.120)$$

From here, calculating the expectation is easy:

$$E(Y_n) = \int_0^\theta 2n \frac{x^{2n-1} \cdot x}{\theta^{2n}} dx = \frac{2n}{2n+1} \theta. \quad (6.121)$$

Because we want  $E(c\hat{\theta}) = cE(\hat{\theta}) = \theta$ , we can just make

$$\boxed{c = \frac{2n+1}{2n}} \quad (6.122)$$

(c) The median  $x_{1/2}$  is such that:

$$\frac{1}{2} = \int_0^{x_{1/2}} \frac{2x}{\theta^2} dx = \frac{x_{1/2}^2}{\theta^2}. \quad (6.123)$$

So,  $x_{1/2} = \theta/\sqrt{2}$ . By the invariance property, we have

$$\hat{x}_{1/2} = \frac{\hat{\theta}}{\sqrt{2}}$$

(6.124)

Since  $E(\hat{\theta}) = (2n+1)/2n \cdot \theta$ , we see that  $\lim_{n \rightarrow \infty} E(\hat{\theta}) = \theta$ . Next, look at the variance of  $\hat{\theta}$ :

$$\begin{aligned} \text{Var}(\hat{\theta}) &= \text{Var}(Y_n) = E[\hat{\theta}^2] - E[\hat{\theta}]^2 \\ &= \int_0^\theta 2n \frac{x^{2n-1} \cdot x^2}{\theta^{2n}} dx - \left( \frac{2n}{2n+1} \right)^2 \theta^2 \\ &= \frac{2n}{2+2n} \theta^2 - \left( \frac{2n}{2n+1} \right)^2 \theta^2. \end{aligned} \quad (6.125)$$

Obviously,  $\lim_{n \rightarrow \infty} 2n/(2+2n) - (2n/(2n+1))^2 = 0$ , so  $\lim_{n \rightarrow \infty} \text{Var}(\hat{\theta}) = 0$ . With this, we conclude  $\hat{\theta}$  is a consistent estimator of  $\theta$ .

□  
□

**6.1.10**

**6.1.10.** Let  $X_1, X_2, \dots, X_n$  be a random sample from a Bernoulli distribution with parameter  $p$ . If  $p$  is restricted so that we know that  $\frac{1}{2} \leq p \leq 1$ , find the mle of this parameter.

*Solution:* The likelihood function is

$$\mathcal{L}(x; p) = \prod_{i=1}^n f(x_i; p) = p^{\sum_{i=1}^n x_i} (1-p)^{n - \sum_{i=1}^n x_i}. \quad (6.126)$$

The log likelihood is then

$$l(p) = \sum_{i=1}^n x_i \ln p + \left( n - \sum_{i=1}^n x_i \right) \ln(1-p). \quad (6.127)$$

Taking  $\partial_p$  of  $l(p)$  gives

$$\partial_p l(p) = \frac{\sum_{i=1}^n x_i}{p} + \frac{n - \sum_{i=1}^n x_i}{1-p}. \quad (6.128)$$

Letting  $\partial_p l(p) = 0$ , we get

$$\hat{p} = \frac{1}{n} \sum_{i=1}^n x_i = \bar{X} \quad (6.129)$$

as expected. Now, since  $1/2 \leq p \leq 1$ , we must consider the case where  $\bar{X} < 1/2$ . If  $\bar{X} < 1/2$ , then because  $p$  cannot take the value of  $\bar{X}$ , it must take the boundary value of  $1/2$  at which  $l(p)$  is maximized. So,

$$\hat{p} = \max\left(\frac{1}{2}, \bar{X}\right)$$

(6.130)

□

### 6.2.2

**6.2.2.** Given  $f(x; \theta) = 1/\theta$ ,  $0 < x < \theta$ , zero elsewhere, with  $\theta > 0$ , formally compute the reciprocal of

$$nE\left\{\left[\frac{\partial \log f(X : \theta)}{\partial \theta}\right]^2\right\}.$$

Compare this with the variance of  $(n+1)Y_n/n$ , where  $Y_n$  is the largest observation of a random sample of size  $n$  from this distribution. Comment.

*Solution:*

$$nE[(\partial_\theta \ln f(x; \theta))^2] = nE[(\partial_\theta(-\ln \theta))^2] = nE\left[\frac{1}{\theta^2}\right] = \frac{n}{\theta^2}. \quad (6.131)$$

The reciprocal of this is therefore

$$\frac{\theta^2}{n}. \quad (6.132)$$

On the other hand,

$$\text{Var}\left[\frac{(n+1)}{n} Y_n\right] = \left(\frac{n+1}{n}\right)^2 (E[Y_n^2] - E[Y_n]^2). \quad (6.133)$$

The cdf of  $x$  is given by  $F_X(x) = x/\theta$ . The cdf of  $Y_n$  is therefore

$$F_Y(x) = \prod_{i=1}^n F_x(x) = \frac{x^n}{\theta^n} \implies f_Y(x) = n \frac{x^{n-1}}{\theta^n}. \quad (6.134)$$

And so

$$E[Y_n] = \int_0^\theta n \frac{x^{n-1} \cdot x}{\theta^n} dx = \frac{n\theta}{n+1}. \quad (6.135)$$

$$E[Y_n^2] = \int_0^\theta n \frac{x^{n-1} \cdot x^2}{\theta^n} dx = \frac{n\theta^2}{n+2}. \quad (6.136)$$

So,

$$\text{Var}\left[\frac{(n+1)}{n} Y_n\right] = \left(\frac{n+1}{n}\right)^2 \left[ \frac{n\theta^2}{n+2} - \left(\frac{n\theta}{n+1}\right)^2 \right] = \frac{\theta^2}{n(n+2)}. \quad (6.137)$$

So, obviously,

$$\text{Var}[Y_n] = \frac{\theta^2}{n(n+2)} < \frac{\theta^2}{n}. \quad (6.138)$$

However, since the support of  $f(x; \theta)$  depends on  $\theta$ , CRLB does not apply.  $\square$

**6.2.7**

**6.2.7.** Recall Exercise 6.1.1 where  $X_1, X_2, \dots, X_n$  is a random sample on  $X$  that has a  $\Gamma(\alpha = 4, \beta = \theta)$  distribution,  $0 < \theta < \infty$ .

- (a) Find the Fisher information  $I(\theta)$ .
- (b) Show that the mle of  $\theta$ , which was derived in Exercise 6.1.1, is an efficient estimator of  $\theta$ .
- (c) Using Theorem 6.2.2, obtain the asymptotic distribution of  $\sqrt{n}(\hat{\theta} - \theta)$ .
- (d) For the data of Example 6.1.1, find the asymptotic 95% confidence interval for  $\theta$ .

*Solution:*

- (a) From 6.1.1. we know that

$$\partial_\theta l(\theta) = -\frac{4n}{\theta} + \frac{1}{\theta^2} \sum_{i=1}^n x_i \implies \partial_\theta^2 l(\theta) = \frac{4n}{\theta^2} - \frac{2}{\theta^3} \sum_{i=1}^n x_i \quad (6.139)$$

And so

$$\begin{aligned} nI(\theta) &= -E \left[ \frac{4n}{\theta^2} - \frac{2}{\theta^3} \sum_{i=1}^n x_i \right] \\ &= -\frac{4n}{\theta^2} + \frac{2}{\theta^3} \sum_{i=1}^n E[x_i] \\ &= -\frac{4n}{\theta^2} + \frac{2}{\theta^3} n4\theta \\ &= -\frac{4n}{\theta^2} + \frac{8n}{\theta^2} \\ &= \boxed{\frac{4n}{\theta^2}} \implies \boxed{I(\theta) = \frac{4}{\theta^2}} \end{aligned} \quad (6.140)$$

- (b) From 6.1.1.,  $\hat{\theta} = \bar{X}/4$ , which is unbiased because

$$E[\hat{\theta}] = \frac{1}{4} E[\bar{X}] = \frac{1}{4} \frac{4n\theta}{n} = \theta. \quad (6.141)$$

This is an efficient estimator of  $\theta$  because

$$\begin{aligned}
 \text{Var}[\hat{\theta}] &= \frac{1}{16} \text{Var}[\bar{X}] = \frac{1}{16n^2} \text{Var} \left[ \sum_{i=1}^n x_i \right] \\
 &= \frac{1}{16n^2} \sum_{i=1}^n \text{Var}[x_i] \\
 &= \frac{1}{16n^2} n 4\theta^2 \\
 &= \frac{\theta^2}{4n}.
 \end{aligned} \tag{6.142}$$

So, we see that

$$\text{Var}[\hat{\theta}] = \frac{1}{nI(\theta)}, \tag{6.143}$$

which means  $\text{Var}(Y)$  attains the Rao-Cramér lower bound.

- (c) The observations are iid with  $\Gamma(\alpha = 4, \beta = \theta)$ . The regularity conditions are also satisfied. Since the Fisher information  $I(\theta) = 4n/\theta^2$  is positive and finite, Theorem 6.2.2. says that

$$\sqrt{n}(\hat{\theta} - \theta) \xrightarrow{D} \mathcal{N} \left( 0, \frac{1}{I(\theta)} \right) = \boxed{\mathcal{N} \left( 0, \frac{\theta^2}{4} \right)} \tag{6.144}$$

- (d) The 95% CI for  $\theta$ , with  $\alpha = 0.025, n = 25, \sigma = \hat{\theta}/2$ , is given by

$$\begin{aligned}
 \left( \hat{\theta} - z_{\alpha/2} \frac{\hat{\theta}/2}{\sqrt{n}}, \hat{\theta} + z_{\alpha/2} \frac{\hat{\theta}/2}{\sqrt{n}} \right) &= \left( 5.03 - 1.96 \frac{5.03/2}{5}, 5.03 + 1.96 \frac{5.03/2}{5} \right) \\
 &= \boxed{(4.04, 6.02)}
 \end{aligned} \tag{6.145}$$

□

### 6.2.8

**6.2.8.** Let  $X$  be  $N(0, \theta)$ ,  $0 < \theta < \infty$ .

- (a) Find the Fisher information  $I(\theta)$ .
- (b) If  $X_1, X_2, \dots, X_n$  is a random sample from this distribution, show that the mle of  $\theta$  is an efficient estimator of  $\theta$ .
- (c) What is the asymptotic distribution of  $\sqrt{n}(\hat{\theta} - \theta)$ ?

*Solution:*

- (a) Well, the likelihood is just the pdf:

$$f(x; \theta) = \frac{1}{\sqrt{2\pi\theta}} e^{-x^2/2\theta}, \theta \in \mathbb{R}^+ \setminus \{\infty\}. \quad (6.146)$$

Then, the log likelihood is

$$l(\theta) = -\frac{1}{2} \ln(2\pi\theta) - \frac{x^2}{2\theta} \implies \partial_\theta l(\theta) = -\frac{1}{2\theta} + \frac{x^2}{2\theta^2}. \quad (6.147)$$

The Fisher information is then

$$\begin{aligned} I(\theta) &= -E[\partial_\theta^2 l(\theta)] \\ &= -E\left[\frac{1}{2\theta^2} - \frac{X^2}{\theta^3}\right] \\ &= -\frac{1}{2\theta^2} + \frac{1}{\theta^3} E[X^2] \\ &= -\frac{1}{2\theta^2} + \frac{1}{\theta^3} (\text{Var}[X] + \cancel{E[X]^2}) \\ &= -\frac{1}{2\theta^2} + \frac{1}{\theta^3} \theta \\ &= \boxed{\frac{1}{2\theta^2}} \end{aligned} \quad (6.148)$$

- (b) We know that the mle of  $\theta$  is given by

$$\hat{\theta} = \frac{1}{n} \sum_{i=1}^n (x_i - 0)^2, \quad (6.149)$$

which is just the sample variance with denominator  $n$ . Now,  $\hat{\theta}$  is an unbiased estimator of  $\theta$  because

$$E[\hat{\theta}] = \frac{1}{n} \sum_{i=1}^n E[X_i^2 + 2X_i \cdot 0 + 0] = \frac{1}{n} \sum_{i=1}^n E[X_i^2] = \frac{n}{n} \text{Var}[X] = \theta. \quad (6.150)$$

Next we want to find  $\text{Var}[\hat{\theta}]$ . Well,

$$\begin{aligned}
 \text{Var}[\hat{\theta}] &= \text{Var}\left[\frac{1}{n} \sum_{i=1}^n X_i^2\right] \\
 &= \frac{1}{n^2} \sum_{i=1}^n \text{Var}[X_i^2] \\
 &= \frac{n}{n^2} (E[X_i^4] - E[X_i^2]^2) \\
 &= \frac{n}{n^2} (E[X_i^4] - \text{Var}[X_i]^2) \\
 &= \frac{n}{n^2} (E[X_i^4] - \theta^2). \tag{6.151}
 \end{aligned}$$

To find  $E[X_i^4]$  we can either brute force:

$$E[X^4] = \int_{\mathbb{R}} x^4 \frac{1}{\sqrt{2\pi\theta}} e^{-x^2/2\theta} dx = 3\theta^2. \tag{6.152}$$

Or use the MGF:

$$\begin{aligned}
 M(t) &= e^{\mu\ell + \theta t^2/2} \\
 \implies E[X^4] &= M^{(4)}(0) = (3\theta^2 + 6t^2\theta^3 + t^4\theta^4)e^{\theta t^2/2} \Big|_{t=0} = 3\theta^2. \tag{6.153}
 \end{aligned}$$

And so,

$$\text{Var}[\hat{\theta}] = \frac{1}{n} (3\theta^2 - \theta^2) = \boxed{\frac{2\theta^2}{n}} \tag{6.154}$$

Because  $\text{Var}[\hat{\theta}] = 2\theta^2/n = 1/nI(\theta)$ , the CRLB is attained. Further, because  $\hat{\theta}$  is an unbiased estimator of  $\theta$ ,  $\hat{\theta}$  is an efficient estimator of  $\theta$ .

- (c) Since all regularity conditions are met, we can use Theorem 6.2.2., which says that

$$\sqrt{n}(\hat{\theta} - \theta) \xrightarrow{D} \mathcal{N}(0, 1/I(\theta)) = \boxed{\mathcal{N}(0, 2\theta^2)} \tag{6.155}$$

□

**6.2.9**

**6.2.9.** If  $X_1, X_2, \dots, X_n$  is a random sample from a distribution with pdf

$$f(x; \theta) = \begin{cases} \frac{3\theta^3}{(x+\theta)^4} & 0 < x < \infty, 0 < \theta < \infty \\ 0 & \text{elsewhere,} \end{cases}$$

show that  $Y = 2\bar{X}$  is an unbiased estimator of  $\theta$  and determine its efficiency.

*Solution:* Well, we first show that  $Y$  is an unbiased estimator of  $\theta$ :

$$E[Y] = 2E[\bar{X}] = \frac{2}{n}nE[X_i] = 2E[X_i] = 2 \int_0^\infty \frac{3x\theta^3}{(x+\theta)^4} dx = 2 \frac{\theta}{2} = \theta. \quad (6.156)$$

So, yes,  $Y$  is an unbiased estimator of  $\theta$ .

The efficiency of  $Y$  is given by

$$\epsilon = \frac{CRLB}{\text{Var}[Y]} = \frac{1}{\text{Var}[Y] \cdot nI(\theta)} \quad (6.157)$$

To find this we next find the Fisher information:

$$\begin{aligned} I(\theta) &= -E[\partial_\theta^2 l(\theta)] \\ &= -E\left[\partial_\theta^2 \ln \frac{3\theta^3}{(X+\theta)^4}\right] \\ &= -E\left[\partial_\theta^2 \ln \frac{3\theta^3}{(X+\theta)^4}\right] \\ &= -E\left[\frac{\theta^2 - 6\theta X - 3X^2}{\theta^2(\theta+X)^2}\right] \\ &= -\int_0^\infty \frac{\theta^2 - 6\theta x - 3x^2}{\theta^2(\theta+x)^2} \cdot \frac{3\theta^3}{(x+\theta)^4} dx = \frac{3}{5\theta^2}. \end{aligned} \quad (6.158)$$

Next, we find the variance of  $Y$ :

$$\begin{aligned} \text{Var}[Y] &= \text{Var}[2\bar{X}] = \frac{4}{n^2} \text{Var}\left[\sum_{i=1}^n X_i\right] = \frac{4n}{n^2} \text{Var}[X_i] = \frac{4}{n} (E[X_i^2] - E[X_i]^2) \\ &= \frac{4}{n} \left\{ \int_0^\infty \frac{3x^2\theta^3}{(x+\theta)^4} dx - \left[ \int_0^\infty \frac{3x\theta^2}{(x+\theta)^4} dx \right]^2 \right\} \\ &= \frac{4}{n} \left( \theta^2 - \frac{\theta^2}{4} \right) = \frac{3\theta^2}{n}. \end{aligned} \quad (6.159)$$

So,

$$\epsilon = \frac{5\theta^2}{3\theta^2 \cdot 3} = \boxed{\frac{5}{9}} \quad (6.160)$$

## 6.6 Problem set 6

### 6.3.1

**6.3.1.** The following data were generated from an exponential distribution with pdf  $f(x; \theta) = (1/\theta)e^{-x/\theta}$ , for  $x > 0$ , where  $\theta = 40$ .


- (a) Histogram the data and locate  $\theta_0 = 50$  on the plot.
- (b) Use the test described in Example 6.3.1 to test  $H_0 : \theta = 50$  versus  $H_1 : \theta \neq 50$ . Determine the decision at level  $\alpha = 0.10$ .

19 15 76 23 24 66 27 12 25 7 6 16 51 26 39

*Solution:*

(a) **R code:**

```
> dt = c(19, 15, 76, 23, 24, 66, 27, 12, 25, 7, 6, 16, 51, 26, 39)
> hist(dt)
> abline(v=50)
```



- (b)  $H_0 : \theta = 50$ ,  $H_a : \theta \neq 50$ . Example 6.3.1 says that we reject  $H_0$  if

$$\frac{2}{\theta_0} \sum_{i=1}^n X_i \geq \chi^2_{1-\alpha/2}(2n) \text{ or } \frac{2}{\theta_0} \sum_{i=1}^n X_i \leq \chi^2_{\alpha/2}(2n) \quad (6.161)$$

In our case,

$$\tau = \frac{2}{\theta_0} \sum_{i=1}^n X_i = 17.28, \quad (6.162)$$

while

$$\begin{aligned}\chi^2_{0.10/2}(30) &= 18.49266 \\ \chi^2_{1-0.10/2}(30) &= 43.77297.\end{aligned}\tag{6.163}$$

Since  $\tau < \chi^2_{0.10/2}(30)$ , we **reject**  $H_0$ .

**R code:**

```
> sum(dt)
[1] 432
> (2/50)*sum(dt)
[1] 17.28
> length(dt)
[1] 15
> qchisq(0.10/2,30)
[1] 18.49266
> qchisq(1-0.10/2,30)
[1] 43.77297
```

□

### 6.3.6

**6.3.6.** Let  $X_1, X_2, \dots, X_n$  be a random sample from a  $N(\mu_0, \sigma^2 = \theta)$  distribution, where  $0 < \theta < \infty$  and  $\mu_0$  is known. Show that the likelihood ratio test of  $H_0 : \theta = \theta_0$  versus  $H_1 : \theta \neq \theta_0$  can be based upon the statistic  $W = \sum_{i=1}^n (X_i - \mu_0)^2 / \theta_0$ . Determine the null distribution of  $W$  and give, explicitly, the rejection rule for a level  $\alpha$  test.

*Solution:* The mle for  $\sigma^2$  is

$$\hat{\sigma}^2 = \frac{1}{n} \sum (X_i - \mu_0)^2. \quad (6.164)$$

With this,

$$\Lambda = \frac{\mathcal{L}(\theta_0)}{\mathcal{L}(\hat{\theta})} = \left( \frac{\hat{\sigma}^2}{\theta_0} \right)^{n/2} \exp \left[ \frac{-1}{2\theta_0} \sum_{i=1}^n (x_i - \mu_0)^2 + \frac{n}{2} \frac{\sum (x_i - \mu_0)^2}{\sum (x_i - \mu_0)^2} \right]. \quad (6.165)$$

This has the form  $\Lambda = n^{-n/2} W^{n/2} e^{-W/2} e^{n/2}$ , which is unimodal. So,  $\Lambda \leq C \implies W \in (c_1, c_2)$  for some  $c_1, c_2$ . This says we can use  $W$  as a testing statistic. Now, under  $H_0 : \theta = \theta_0$ , we have that  $X_i \sim N(\mu_0, \theta = \theta_0)$ , and so  $(X_i - \mu_0) / \sqrt{\theta_0} \sim N(0, 1)$ . This means  $W = \sum^n (X_i - \mu_0)^2 / \theta_0 \sim \chi^2(n)$ .

We reject whenever  $W \geq \chi^2_{1-\alpha/2}(n)$  or  $W \leq \chi^2_{\alpha/2}(n)$ . □

### 6.3.11

**6.3.11.** Let  $X_1, X_2, \dots, X_n$  be a random sample from a  $\Gamma(\alpha = 4, \beta = \theta)$  distribution, where  $0 < \theta < \infty$ .

- (a) Show that the likelihood ratio test of  $H_0 : \theta = \theta_0$  versus  $H_1 : \theta \neq \theta_0$  is based upon the statistic  $W = \sum_{i=1}^n X_i$ . Obtain the null distribution of  $2W/\theta_0$ .
- (b) For  $\theta_0 = 3$  and  $n = 5$ , find  $c_1$  and  $c_2$  so that the test that rejects  $H_0$  when  $W \leq c_1$  or  $W \geq c_2$  has significance level 0.05.

*Solution:*

- (a) We know that

$$\hat{\beta} = \frac{1}{n\alpha} \sum_{i=1}^n X_i = \frac{\bar{X}}{\alpha} = \frac{\bar{X}}{4}. \quad (6.166)$$

$$\begin{aligned} \Lambda &= \frac{\mathcal{L}(\hat{\theta})}{\mathcal{L}(\theta_0)} = \frac{1/\hat{\theta}^{4n}}{1/\theta_0^{4n}} \exp \left[ -\left( \frac{1}{\hat{\theta}} - \frac{1}{\theta_0} \right) \sum x_i \right] \\ &\sim (4n)^{-4n} \left( \sum x_i / \theta_0 \right)^{-4n} \exp \left[ -\sum X_i / \theta_0 \right] \exp [4n] \\ &= (4n)^{-4n} W^{-4n} \theta_0^{4n} \exp [-W/\theta_0] \exp [4n]. \end{aligned} \quad (6.167)$$

So,  $\Lambda$  depends on  $W$  as desired. Now, since  $X_i \sim \Gamma(4, \theta_0)$  under  $H_0$ , we have

$$\frac{2W}{\theta_0} = \frac{2}{\theta_0} \sum X_i \sim \Gamma \left( n\alpha, \frac{2\theta_0}{\theta_0} \right) = \Gamma(4n, 2) = \boxed{\chi^2(4n)} \quad (6.168)$$

(b)

$$\begin{aligned} c_1 &= \frac{\theta_0}{2} \chi_{\alpha/2}^2(4n) = \frac{3}{2} \times 9.590777 = \boxed{14.38617} \\ c_2 &= \frac{\theta_0}{2} \chi_{1-\alpha/2}^2(4n) = \frac{3}{2} \times 34.16961 = \boxed{51.25441} \end{aligned} \quad (6.169)$$

**R code:**

```
> qchisq(1-0.05/2, 4*5)
[1] 34.16961
> qchisq(0.05/2, 4*5)
[1] 9.590777
> 34.16961*1.5
[1] 51.25441
> 9.590777*1.5
[1] 14.38617
```

□

**6.3.18**

**6.3.18.** Let  $X_1, X_2, \dots, X_n$  be a random sample from a  $\Gamma(\alpha, \beta)$  distribution where  $\alpha$  is known and  $\beta > 0$ . Determine the likelihood ratio test for  $H_0 : \beta = \beta_0$  against  $H_1 : \beta \neq \beta_0$ .

*Solution:* The setup of this problem is exactly like that of 6.3.11. The likelihood ratio test of  $H_0 : \beta = \beta_0$  versus  $H_a : \beta \neq \beta_0$  is based upon the statistic  $W = \sum X_i$ . The null distribution of  $2W/\beta_0$  is  $\chi^2(n\alpha)$ , under  $H_0$ . We reject whenever  $W \leq c_1$  or  $W \geq c_2$ , where

$$\begin{aligned} c_1 &= \frac{\beta_0}{2} \chi_{\alpha/2}^2(\alpha n) \\ c_2 &= \frac{\beta_0}{2} \chi_{1-\alpha/2}^2(\alpha n). \end{aligned} \tag{6.170}$$

□

### 6.4.1

**6.4.1.** A survey is taken of the citizens in a city as to whether or not they support the zoning plan that the city council is considering. The responses are: Yes, No, Indifferent, and Otherwise. Let  $p_1, p_2, p_3$ , and  $p_4$  denote the respective true probabilities of these responses. The results of the survey are:

Yes	No	Indifferent	Otherwise
60	45	70	25

- (a) Obtain the mles of  $p_i$ ,  $i = 1, \dots, 4$ .
- (b) Obtain 95% confidence intervals, (4.2.7), for  $p_i$ ,  $i = 1, \dots, 4$ .

*Solution:*

- (a) The mle for each  $p_i$  is

$$\hat{p}_i = \frac{\# \text{ Observations of } i}{n}. \quad (6.171)$$

- (b)

$$\begin{aligned} CI_1 &= \left( \hat{p}_1 - z_{\alpha/2} \sqrt{\hat{p}_1(1 - \hat{p}_1)/n}, \hat{p}_1 + z_{\alpha/2} \sqrt{\hat{p}_1(1 - \hat{p}_1)/n} \right) \\ &= \left( \frac{60}{200} - 1.96 \times 0.0324037, \frac{60}{200} + 1.96 \times 0.0324037 \right) \\ &= \boxed{(0.2364887, 0.3635113)} \end{aligned} \quad (6.172)$$

$$\begin{aligned} CI_2 &= \left( \frac{45}{200} - 1.96 \times 0.02952753, \frac{45}{200} + 1.96 \times 0.02952753 \right) \\ &= \boxed{(0.167126, 0.282874)} \end{aligned} \quad (6.173)$$

$$\begin{aligned} CI_3 &= \left( \frac{70}{200} - 1.96 \times 0.03372684, \frac{70}{200} + 1.96 \times 0.03372684 \right) \\ &= \boxed{(0.2838954, 0.4161046)} \end{aligned} \quad (6.174)$$

$$\begin{aligned} CI_4 &= \left( \frac{25}{200} - 1.96 \times 0.02338536, \frac{25}{200} + 1.96 \times 0.02338536 \right) \\ &= \boxed{(0.0791647, 0.7732091)} \end{aligned} \quad (6.175)$$

□

**6.4.3**

**6.4.3.** Let  $X_1, X_2, \dots, X_n$  be iid, each with the distribution having pdf  $f(x; \theta_1, \theta_2) = (1/\theta_2)e^{-(x-\theta_1)/\theta_2}$ ,  $\theta_1 \leq x < \infty$ ,  $-\infty < \theta_2 < \infty$ , zero elsewhere. Find the maximum likelihood estimators of  $\theta_1$  and  $\theta_2$ .

*Solution:* The likelihood function is

$$\mathcal{L}(\theta_1, \theta_2) = \frac{1}{\theta_2^n} \exp \left[ -\frac{1}{\theta_2} \sum_{i=1}^n (x_i - \theta_1) \right]. \quad (6.176)$$

with the support given in the problem. We find

$$\ln \mathcal{L} = -n \ln \theta_2 - \frac{1}{\theta_2} \sum_{i=1}^n (x_i - \theta_1). \quad (6.177)$$

And so,

$$\begin{aligned} \partial_{\theta_1} \ln \mathcal{L}(\theta_1, \theta_2) &= \frac{n}{\theta_2} > 0 \implies \boxed{\hat{\theta}_1 = \text{Argmax } \ln \mathcal{L}(\theta_1, \theta_2) = \min_i X_i} \\ \partial_{\theta_2} \ln \mathcal{L}(\hat{\theta}_1, \theta_2) &= -\frac{n}{\theta_2} + \frac{1}{\theta_2^2} \sum_{i=1}^n (x_i - \hat{\theta}_1) = 0 \implies \boxed{\hat{\theta}_2 = \frac{1}{n} \sum_{i=1}^n (X_i - \min_i X_i)} \end{aligned} \quad (6.178)$$

where we have used  $\hat{\theta}_1 = \min_i X_i$  in writing the mle for  $\theta_2$ .  $\square$

**6.4.5**

**6.4.5.** Let  $Y_1 < Y_2 < \dots < Y_n$  be the order statistics of a random sample of size  $n$  from the uniform distribution of the continuous type over the closed interval  $[\theta - \rho, \theta + \rho]$ . Find the maximum likelihood estimators for  $\theta$  and  $\rho$ . Are these two unbiased estimators?

*Solution:* It's not difficult to see that

$$\begin{aligned}\hat{\theta} - \hat{\rho} &= \min Y_i = Y_1 \\ \hat{\theta} + \hat{\rho} &= \max Y_i = Y_n.\end{aligned}\tag{6.179}$$

Solving this for  $\hat{\theta}, \hat{\rho}$  gives

$$\begin{aligned}\hat{\theta} &= \frac{Y_n + Y_1}{2} \\ \hat{\rho} &= \frac{Y_n - Y_1}{2}.\end{aligned}\tag{6.180}$$

Next,

$$\begin{aligned}F_{Y_1}(x) &= P(Y_1 \leq x) = P(\min_i Y_i \leq x) \\ &= 1 - P(\min_i Y_i > x) \\ &= 1 - \prod_{i=1}^n P(Y_i > x) \\ &= 1 - (1 - F_{Y_i}(x))^n = 1 - \left(1 - \int_{\theta-\rho}^x \frac{1}{2\rho} dx\right)^n \\ &= 1 - \left(1 - \frac{x - \theta + \rho}{2\rho}\right)^n \\ \implies f_{Y_1}(x) &= \frac{n(\theta + \rho - x)^{n-1}}{(2\rho)^n}\end{aligned}\tag{6.181}$$

And so,

$$E[Y_1] = \int_{\theta-\rho}^{\theta+\rho} x \frac{n(\theta + \rho - x)^{n-1}}{(2\rho)^n} dx = \frac{1-n}{1+n} \rho + \theta\tag{6.182}$$

Next,

$$\begin{aligned}
 F_{Y_n}(x) &= P(Y_n \leq x) = P(\max_i Y_i \leq x) \\
 &= \prod_{i=1}^n P(Y_i \leq x) = \prod_{i=1}^n F_{Y_i} \\
 &= \frac{(x - \theta + \rho)^n}{(2\rho)^n} \\
 \implies f_{Y_n}(x) &= \frac{n(x - \theta + \rho)^{n-1}}{(2\rho)^n}.
 \end{aligned} \tag{6.183}$$

And so,

$$E[Y_n] = \int_{\theta-\rho}^{\theta+\rho} x \frac{n(x - \theta + \rho)^{n-1}}{(2\rho)^n} dx = \frac{n-1}{n+1} \rho + \theta. \tag{6.184}$$

With these,

$$\begin{aligned}
 E[\hat{\theta}] &= E\left[\frac{Y_n + Y_1}{2}\right] = \frac{1}{2}E[2\theta] = \theta \\
 E[\hat{\rho}] &= E\left[\frac{Y_n - Y_1}{2}\right] = \frac{1}{2}E\left[\frac{n-1-1+n}{n+1}\rho\right] = \rho.
 \end{aligned} \tag{6.185}$$

So,  $\hat{\theta}$  and  $\hat{\rho}$  are unbiased.  $\square$

### 6.5.3

**6.5.3.** Consider the survey of citizens discussed in Exercise 6.4.1. Suppose that the hypotheses of interest are  $H_0 : p_1 = p_2$  versus  $H_1 : p_1 \neq p_2$ . Note that computation can be carried out using the R function `p2pair.R`, which can be downloaded at the site mentioned in the Preface.

- (a) Test these hypotheses at level  $\alpha = 0.05$  using the test (6.5.20). Conclude in terms of the problem.
- (b) Obtain the 95% confidence interval, (6.5.21), for  $p_1 - p_2$ . What does the confidence interval mean in terms of the problem?

*Solution:*

- (a)  $H_0 : p_1 = p_2$  and  $H_a : p_1 \neq p_2$ .

$$\begin{aligned}\chi_W^2 &= \frac{(\hat{p}_1 - \hat{p}_2)^2}{(\hat{p}_1 + \hat{p}_2 - (\hat{p}_1 - \hat{p}_2)^2)/n} \\ &= \frac{(60/200 - 45/200)^2}{(60/200 + 45/200 - (60/200 - 45/200)^2)/200} \\ &= \boxed{2.167}\end{aligned}\tag{6.186}$$

Clearly,  $2.167 < 3.81 = \chi_{0.05}^2(1)$ . So, we fail to reject  $H_0$  – there is not enough evidence to reject the null hypothesis that the proportions of people who vote Yes and No are the same.

- (b) The 95% CI for  $p_1 - p_2$  is

$$\left( \frac{60}{200} - \frac{45}{200} \right) \pm z_{0.05/2} \left( \frac{\frac{60}{200} + \frac{45}{200} - \left( \frac{60}{200} - \frac{45}{200} \right)^2}{200} \right)^{1/2} \tag{6.187}$$

$$= (-0.025, 0.175). \tag{6.188}$$

Since the CI contains 0, we fail to reject  $H_0$ .

□

**6.5.6**

**6.5.6.** Let  $X_1, X_2, \dots, X_n$  and  $Y_1, Y_2, \dots, Y_m$  be independent random samples from the two normal distributions  $N(0, \theta_1)$  and  $N(0, \theta_2)$ .

- (a) Find the likelihood ratio  $\Lambda$  for testing the composite hypothesis  $H_0 : \theta_1 = \theta_2$  against the composite alternative  $H_1 : \theta_1 \neq \theta_2$ .
- (b) This  $\Lambda$  is a function of what  $F$ -statistic that would actually be used in this test?

*Solution:*

- (a)  $H_0 : \theta_1 = \theta_2$  and  $H_a : \theta_1 \neq \theta_2$ .  $X_i \sim N(0, \theta_1)$  and  $Y_i \sim N(0, \theta_2)$ . So,

$$\mathcal{L}(\theta_1, \theta_2) = (2\pi\theta_1)^{-n/2} (2\pi\theta_2)^{-m/2} \exp \left[ -\frac{1}{2\theta_1} \sum_{i=1}^n x_i^2 - \frac{1}{2\theta_2} \sum_{i=1}^m y_i^2 \right]. \quad (6.189)$$

Under  $H_0 : \theta_1 = \theta_2$ :

$$\begin{aligned} \partial_\theta \ln \mathcal{L}(\theta_1 = \theta_2 = \theta) &= 0 \implies -\frac{n+m}{2} - \frac{1}{2\theta} \left[ \sum_{i=1}^n x_i^2 + \sum_{i=1}^m y_i^2 \right] = 0 \\ &\implies \hat{\theta} = \frac{1}{n+m} \left[ \sum_{i=1}^n x_i^2 + \sum_{i=1}^m y_i^2 \right]. \end{aligned} \quad (6.190)$$

With this,

$$\begin{aligned} \mathcal{L}(\hat{\Theta}_0) &= \left( \frac{2\pi}{n+m} \left[ \sum_{i=1}^n x_i^2 + \sum_{i=1}^m y_i^2 \right] \right)^{-\frac{n+m}{2}} \exp \left[ -\frac{\sum_{i=1}^n x_i^2 + \sum_{i=1}^m y_i^2}{\frac{2}{n+m} (\sum_{i=1}^n x_i^2 + \sum_{i=1}^m y_i^2)} \right] \\ &= \left( \frac{2\pi}{n+m} \left[ \sum_{i=1}^n x_i^2 + \sum_{i=1}^m y_i^2 \right] \right)^{-\frac{n+m}{2}} \exp \left[ \frac{-n-m}{2} \right]. \end{aligned} \quad (6.191)$$

In the joint space,

$$\begin{aligned} \hat{\theta}_1 &= \frac{1}{n} \sum_{i=1}^n x_i^2 \\ \hat{\theta}_2 &= \frac{1}{m} \sum_{i=1}^m y_i^2. \end{aligned} \quad (6.192)$$

With these,

$$\mathcal{L}(\hat{\Theta}) = \left( \frac{2\pi}{n} \sum_{i=1}^n x_i^2 \right)^{-n/2} \left( \frac{2\pi}{m} \sum_{i=1}^m y_i^2 \right)^{-m/2} \exp \left[ \frac{-n-m}{2} \right]. \quad (6.193)$$

And so we have

$$\begin{aligned}
 \Lambda &= \frac{\mathcal{L}_0}{\mathcal{L}} = \frac{\left(\frac{1}{n+m} [\sum^n x_i^2 + \sum^m y_i^2]\right)^{-\frac{n+m}{2}}}{\left(\frac{1}{n} \sum^n x_i^2\right)^{-n/2} \left(\frac{1}{m} \sum^m y_i^2\right)^{-m/2}} \\
 &= \frac{(n+m)^{\frac{n+m}{2}}}{n^{n/2} m^{m/2}} \frac{(\sum^n x_i^2)^{n/2} (\sum^m y_i^2)^{m/2}}{[\sum^n x_i^2 + \sum^m y_i^2]^{\frac{n+m}{2}}} \\
 &= \frac{(n+m)^{\frac{n+m}{2}}}{n^{n/2} m^{m/2}} \left( \underbrace{\sum^n x_i^2}_{T} \right)^{n/2} \left( \underbrace{\sum^m y_i^2}_{1-T} \right)^{m/2}.
 \end{aligned} \tag{6.194}$$

(b)  $\Lambda \leq k$  is equivalent to  $W \leq c_1$  or  $W \geq c_2$  with

$$W = \frac{\frac{1}{n} \sum^n x_i^2}{\frac{1}{m} \sum^m y_i^2} \sim F(n, m)$$

(6.195)

such that  $1/T = 1 + 1/W$ . This is because  $\sum^n x_i^2/\theta \sim \chi^2(n)$ ,  $\sum^m y_i^2/\theta \sim \chi^2(m)$  under  $H_0$ .

□

**6.5.7**

**6.5.7.** Let  $X$  and  $Y$  be two independent random variables with respective pdfs

$$f(x; \theta_i) = \begin{cases} \left(\frac{1}{\theta_i}\right) e^{-x/\theta_i} & 0 < x < \infty, 0 < \theta_i < \infty \\ 0 & \text{elsewhere,} \end{cases}$$

for  $i = 1, 2$ . To test  $H_0 : \theta_1 = \theta_2$  against  $H_1 : \theta_1 \neq \theta_2$ , two independent samples of sizes  $n_1$  and  $n_2$ , respectively, were taken from these distributions. Find the likelihood ratio  $\Lambda$  and show that  $\Lambda$  can be written as a function of a statistic having an  $F$ -distribution, under  $H_0$ .

*Solution:* We did this in “class,” so I will skip a few steps. Let  $X_i \sim \exp(\theta_1)$  and  $Y_i \sim \exp(\theta_2)$ .  $H_0 : \theta_1 = \theta_2$  and  $H_a : \theta_1 \neq \theta_2$ . The null space is  $\{\{\theta_1, \theta_2\} : \theta_1 = \theta_2 > 0\}$ . The alternative space is  $\{\{\theta_1, \theta_2\} : \theta_1 \neq \theta_2, \theta_1, \theta_2 \neq 0\}$ . The joint space is just  $\{\{\theta_1, \theta_2\}, \theta_1 > 0, \theta_2 > 0\}$ . It’s easy to see that

$$\mathcal{L}(\theta_1, \theta_2 | X, Y) = \frac{1}{\theta_1^n} e^{\sum x_i / \theta_1} \frac{1}{\theta_2^m} e^{\sum y_i / \theta_2}. \quad (6.196)$$

From here one finds that under  $H_0 : \theta_1 = \theta_2 = \theta$ :

$$l(\Theta_0) = -(n+m) \ln \theta - \frac{1}{\theta} \left( \sum x_i + \sum y_i \right). \quad (6.197)$$

And so

$$\partial_\theta l(\theta) = 0 \implies \hat{\theta}_0 = \frac{1}{n+m} \left( \sum x_i + \sum y_i \right). \quad (6.198)$$

With this, we can plug back in to calculate the numerator:

$$\mathcal{L}(\hat{\Theta}_0) = \dots = \left( \frac{1}{n+m} \left[ \sum x_i + \sum y_j \right] \right)^{-n-m} \exp(-n-m). \quad (6.199)$$

In the joint space,

$$l(\Theta) = -n \ln \theta - \frac{1}{\theta} \sum x_i - m \ln \mu - \frac{1}{\mu} \sum y_j. \quad (6.200)$$

And so

$$\partial_{\theta_1} l(\Theta) = 0 \implies \hat{\theta}_1 = \bar{x} \quad (6.201)$$

$$\partial_{\theta_2} l(\Theta) = 0 \implies \hat{\theta}_2 = \bar{y}. \quad (6.202)$$

With these,

$$\mathcal{L}(\hat{\Theta}) = \dots \quad (6.203)$$

$$= \frac{1}{\left(\frac{1}{n} \sum x_i\right)^n} \exp \left[ -\frac{1}{\frac{1}{n} \sum x_i} \sum x_i \right] \frac{1}{\left(\frac{1}{m} \sum y_i\right)^m} \exp \left[ -\frac{1}{\frac{1}{m} \sum y_i} \sum y_i \right] \quad (6.204)$$

$$= \frac{1}{\left(\frac{1}{n} \sum x_i\right)^n} e^{-n} \frac{1}{\left(\frac{1}{m} \sum y_i\right)^m} e^{-m}. \quad (6.205)$$

Putting everything together, we find

$$\Lambda = \frac{\mathcal{L}_0}{\mathcal{L}} = \dots = \frac{\frac{1}{(n+m)^{-n-m}} (\sum x_i + \sum y_j)^{-n-m}}{\frac{1}{n^n} \frac{1}{m^m} (\sum x_i)^{-n} (\sum y_i)^{-m}}. \quad (6.206)$$

We reject if  $\Lambda < c$ , iff

$$\frac{(n+m)^{n+m}}{n^n m^m} \frac{(\sum x_i)^n (\sum y_i)^m}{(\sum x_i + \sum y_j)^{n+m}} < c \quad (6.207)$$

iff (letting  $c$  absorb the constant)

$$\frac{(\sum x_i)^n (\sum y_i)^m}{(\sum x_i + \sum y_j)^{n+m}} < c' \quad (6.208)$$

What does the distribution of  $\Lambda$  look like? Notice that we reject if

$$\Lambda = \frac{(n+m)^{n+m}}{n^n m^m} \frac{(\sum x_i)^n (\sum y_i)^m}{(\sum x_i + \sum y_j)^{n+m}} \quad (6.209)$$

$$= \frac{(n+m)^{n+m}}{n^n m^m} \left( \frac{\sum x_i}{\sum x_i + \sum y_j} \right)^n \left( \frac{\sum y_i}{\sum x_i + \sum y_j} \right)^m \quad (6.210)$$

$$= \frac{(n+m)^{n+m}}{n^n m^m} T^n (1-T)^m < c \quad (6.211)$$

which is akin to saying that we reject if  $T < a$  or  $T > b$ . What is the distribution of  $T$  under  $H_0$ ? Note that under  $H_0$ ,  $\sum x_i \sim \Gamma(n, \theta)$  and  $\sum y_i \sim \Gamma(m, \theta)$ . By transformation method, we can show that  $\boxed{T \sim \beta(n, m)}$ .

Notice that

$$\frac{1}{T} = 1 + \frac{\sum y_j}{\sum x_i} = 1 + \frac{m \sum y_j / m}{n \sum x_i / n} \quad (6.212)$$

Since  $\sum X_i \sim \Gamma(n, \theta)$ ,  $\sum Y_i \sim \Gamma(m, \theta)$  under  $H_0$  we have that  $2\theta X_i \sim \chi^2(2n)$  and  $2\theta Y_i \sim \chi^2(2m)$ . This means we can call

$$\boxed{W = \frac{\sum y_j / m}{\sum x_i / n} = \frac{2\theta \sum y_i / m}{2\theta \sum x_i / n} \sim F(2m, 2n)}. \quad (6.213)$$

□

## 6.7 Problem set 7

**6.8 Problem set 8**