


# The Hydrogen Atom and Harmonic Oscillator(s)

Huan Bui

MIT

ZGS, Mar 24, 2023


# Harmonic oscillator universe?



Harmonic oscillator in physics:

- Hooke's law
- QHO
- Einstein solid
- Atom-radiation interaction
- 2nd quantization of EM fields
- QFT
- ...

# Harmonic oscillator universe?



Harmonic oscillator in physics:

- Hooke's law
- QHO
- Einstein solid
- Atom-radiation interaction
- 2nd quantization of EM fields
- QFT
- ...
- Gravity? Inverse-square law?

# Coulomb-Kepler problem revisited

A particle in a central potential  $V(r) = -k/r$ :

$$H = \frac{\vec{p}^2}{2m} - \frac{k}{r}.$$

# Coulomb-Kepler problem revisited

A particle in a central potential  $V(r) = -k/r$ :

$$H = \frac{\vec{p}^2}{2m} - \frac{k}{r}.$$

Constants of motion:  $H$ ,

# Coulomb-Kepler problem revisited

A particle in a central potential  $V(r) = -k/r$ :

$$H = \frac{\vec{p}^2}{2m} - \frac{k}{r}.$$

Constants of motion:  $H, \vec{L} = \vec{r} \times \vec{p},$

# Coulomb-Kepler problem revisited

A particle in a central potential  $V(r) = -k/r$ :

$$H = \frac{\vec{p}^2}{2m} - \frac{k}{r}.$$

Constants of motion:  $H$ ,  $\vec{L} = \vec{r} \times \vec{p}$ , and  $\vec{A}$ ,

# Coulomb-Kepler problem revisited

A particle in a central potential  $V(r) = -k/r$ :

$$H = \frac{\vec{p}^2}{2m} - \frac{k}{r}.$$


Constants of motion:  $H$ ,  $\vec{L} = \vec{r} \times \vec{p}$ , and  $\vec{A}$ , the Laplace-Runge-Lenz vector:

$$\vec{A} = \vec{p} \times \vec{L} - mk \frac{\vec{r}}{r}.$$




# Coulomb-Kepler problem revisited

Brief review:



# Coulomb-Kepler problem revisited


Brief review:



$\vec{A}$  is in the plane of the orbit (so  $\vec{A} \cdot \vec{L} = 0$ ), with  $A^2 = m^2 k^2 + 2mEL^2$ .

# Coulomb-Kepler problem revisited

Brief review:



$\vec{A}$  is in the plane of the orbit (so  $\vec{A} \cdot \vec{L} = 0$ ), with  $A^2 = m^2 k^2 + 2mEL^2$ .

$\vec{A}$  determines the shape and orientation of the orbit.

# Coulomb-Kepler problem revisited

$\vec{A}$  determines the shape and orientation of the orbit:

# Coulomb-Kepler problem revisited

$\vec{A}$  determines the shape and orientation of the orbit:

- Shape: from

$$\vec{A} \cdot \vec{r} = \vec{r} \cdot (\vec{p} \times \vec{L}) - mk = (\vec{r} \times \vec{p}) \cdot \vec{L} - mk = L^2 - mk = Ar \cos \theta$$

# Coulomb-Kepler problem revisited

$\vec{A}$  determines the shape and orientation of the orbit:

- Shape: from

$$\vec{A} \cdot \vec{r} = \vec{r} \cdot (\vec{p} \times \vec{L}) - mk = (\vec{r} \times \vec{p}) \cdot \vec{L} - mk = L^2 - mk = Ar \cos \theta$$

we get the orbit equation

$$\frac{1}{r} = \frac{mk}{L^2} (1 + \epsilon \cos \theta), \quad \text{eccentricity } \epsilon = \frac{A}{|mk|} = \sqrt{1 + \frac{2EL^2}{mk^2}} \geq 0$$

# Coulomb-Kepler problem revisited

$\vec{A}$  determines the shape and orientation of the orbit:

- Shape: from

$$\vec{A} \cdot \vec{r} = \vec{r} \cdot (\vec{p} \times \vec{L}) - mk = (\vec{r} \times \vec{p}) \cdot \vec{L} - mk = L^2 - mk = Ar \cos \theta$$

we get the orbit equation

$$\frac{1}{r} = \frac{mk}{L^2} (1 + \epsilon \cos \theta), \quad \text{eccentricity } \epsilon = \frac{A}{|mk|} = \sqrt{1 + \frac{2EL^2}{mk^2}} \geq 0$$

- Orientation:  $\vec{A}$  points from source to periapsis

## Aside: Brief history of the LRL vector



## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

- Neither Laplace, Runge, nor Lenz discovered it. Laplace (1799)

## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

- Neither Laplace, Runge, nor Lenz discovered it. Laplace (1799)
- Jakob Hermann discovered  $|\vec{A}|$  (1710), recognized its relation to  $\epsilon$

## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

- Neither Laplace, Runge, nor Lenz discovered it. Laplace (1799)
- Jakob Hermann discovered  $|\vec{A}|$  (1710), recognized its relation to  $\epsilon$
- Johann Bernoulli generalized to  $\vec{A}$  (1710)

## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

- Neither Laplace, Runge, nor Lenz discovered it. Laplace (1799)
- Jakob Hermann discovered  $|\vec{A}|$  (1710), recognized its relation to  $\epsilon$
- Johann Bernoulli generalized to  $\vec{A}$  (1710)
- Hamilton "rediscovered"  $\vec{A}$  as  $\vec{A}/mk$  (~1850)

## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

- Neither Laplace, Runge, nor Lenz discovered it. Laplace (1799)
- Jakob Hermann discovered  $|\vec{A}|$  (1710), recognized its relation to  $\epsilon$
- Johann Bernoulli generalized to  $\vec{A}$  (1710)
- Hamilton "rediscovered"  $\vec{A}$  as  $\vec{A}/mk$  (~1850)

Pauli and the LRL vector in early QM (1926):

## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

- Neither Laplace, Runge, nor Lenz discovered it. Laplace (1799)
- Jakob Hermann discovered  $|\vec{A}|$  (1710), recognized its relation to  $\epsilon$
- Johann Bernoulli generalized to  $\vec{A}$  (1710)
- Hamilton "rediscovered"  $\vec{A}$  as  $\vec{A}/mk$  (~1850)

Pauli and the LRL vector in early QM (1926):

- Used  $\vec{A}$  to derive the spectrum of hydrogen (pre-SE!)

## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

- Neither Laplace, Runge, nor Lenz discovered it. Laplace (1799)
- Jakob Hermann discovered  $|\vec{A}|$  (1710), recognized its relation to  $\epsilon$
- Johann Bernoulli generalized to  $\vec{A}$  (1710)
- Hamilton "rediscovered"  $\vec{A}$  as  $\vec{A}/mk$  (~1850)

Pauli and the LRL vector in early QM (1926):

- Used  $\vec{A}$  to derive the spectrum of hydrogen (pre-SE!)
- Derived energy shifts in the presence of  $\vec{E}$  and  $\vec{B}$



## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

- Neither Laplace, Runge, nor Lenz discovered it. Laplace (1799)
- Jakob Hermann discovered  $|\vec{A}|$  (1710), recognized its relation to  $\epsilon$
- Johann Bernoulli generalized to  $\vec{A}$  (1710)
- Hamilton "rediscovered"  $\vec{A}$  as  $\vec{A}/mk$  (~1850)

Pauli and the LRL vector in early QM (1926):

- Used  $\vec{A}$  to derive the spectrum of hydrogen (pre-SE!)
- Derived energy shifts in the presence of  $\vec{E}$  and  $\vec{B}$

## Aside: Brief history of the LRL vector

Unclear origin, gets rediscovered repeatedly:

- Neither Laplace, Runge, nor Lenz discovered it. Laplace (1799)
- Jakob Hermann discovered  $|\vec{A}|$  (1710), recognized its relation to  $\epsilon$
- Johann Bernoulli generalized to  $\vec{A}$  (1710)
- Hamilton "rediscovered"  $\vec{A}$  as  $\vec{A}/mk$  (~1850)

Pauli and the LRL vector in early QM (1926):

- Used  $\vec{A}$  to derive the spectrum of hydrogen (pre-SE!)
- Derived energy shifts in the presence of  $\vec{E}$  and  $\vec{B}$

Further readings (so fun):

- History: [1], [2], [3], [4], [5]
- "Discoveries" and application: [6], [7], [8], [9], [10], [11]

# The Hydrogen Atom

The energy levels and wavefunctions for the bound states of hydrogen are gotten by solving the Schrödinger equation:

$$\left\{ -\frac{\hbar^2}{2m} \nabla^2 - \frac{e^2}{r} \right\} \psi = E\psi, \quad E < 0.$$

# The Hydrogen Atom

The energy levels and wavefunctions for the bound states of hydrogen are gotten by solving the Schrödinger equation:

$$\left\{ -\frac{\hbar^2}{2m} \nabla^2 - \frac{e^2}{r} \right\} \psi = E\psi, \quad E < 0.$$

With

$$\lambda = \frac{8}{a} \quad \alpha^4 = -\frac{8E}{e^2 a}, \quad a = \frac{\hbar^2}{me^2}, \quad (1)$$

the SE becomes

$$\left\{ 4\nabla^2 + \frac{\lambda}{r} - \alpha^4 \right\} \psi = 0. \quad (2)$$

# Where are the harmonic oscillators?

Following [12], introduce coordinates  $\zeta_A, \zeta_B \in \mathbb{C}$  and demand

$$x + iy = 2\zeta_A \overline{\zeta_B} \qquad z = \zeta_A \overline{\zeta_A} - \zeta_B \overline{\zeta_B}.$$

# Where are the harmonic oscillators?

Following [12], introduce coordinates  $\zeta_A, \zeta_B \in \mathbb{C}$  and demand

$$x + iy = 2\zeta_A \overline{\zeta_B} \qquad z = \zeta_A \overline{\zeta_A} - \zeta_B \overline{\zeta_B}.$$

With this,

$$r = \sqrt{x^2 + y^2 + z^2} = \zeta_A \overline{\zeta_A} + \zeta_B \overline{\zeta_B}$$

Note:

- Each pair  $(\zeta_A, \zeta_B)$  gives a unique point  $(x, y, z)$
- Converse is true up to arbitrary but equal arguments of  $\zeta_A, \zeta_B$

# Where are the harmonic oscillators?

Let  $\sigma = 2 \arg(\zeta_A) = 2 \arg(\zeta_B)$ . Can write  $\zeta_A, \zeta_B$  in spherical coordinates:

$$\zeta_A = r^{1/2} e^{i(\sigma+\varphi)/2} \cos \frac{\theta}{2} \quad \zeta_B = r^{1/2} e^{i(\sigma-\varphi)/2} \sin \frac{\theta}{2} \quad (3)$$

$\implies (x, y, z)$  determines  $(\zeta_A, \zeta_B)$  up to  $e^{i\sigma}$ .

# Where are the harmonic oscillators?

Let  $\sigma = 2 \arg(\zeta_A) = 2 \arg(\zeta_B)$ . Can write  $\zeta_A, \zeta_B$  in spherical coordinates:

$$\zeta_A = r^{1/2} e^{i(\sigma+\varphi)/2} \cos \frac{\theta}{2} \quad \zeta_B = r^{1/2} e^{i(\sigma-\varphi)/2} \sin \frac{\theta}{2} \quad (3)$$

$\implies (x, y, z)$  determines  $(\zeta_A, \zeta_B)$  up to  $e^{i\sigma}$ .

Can show that

$$r \nabla^2 \psi = (\partial_A \partial_{\bar{A}} + \partial_B \partial_{\bar{B}}) \psi.$$

$\implies$  Can now write SE in terms of  $\zeta_A, \zeta_B, \overline{\zeta_A}, \overline{\zeta_B}$ .



# Where are the harmonic oscillators?

SE in terms of  $\zeta_A, \zeta_B, \overline{\zeta_A}, \overline{\zeta_B}$ :

$$\left\{4\nabla^2 + \frac{\lambda}{r} - \alpha^4\right\}\psi = \left\{4(\partial_A\partial_{\bar{A}} + \partial_B\partial_{\bar{B}}) + \lambda - \alpha^4(\zeta_A\overline{\zeta_A} + \zeta_B\overline{\zeta_B})\right\}\psi = 0 \quad (4)$$

# Where are the harmonic oscillators?

SE in terms of  $\zeta_A, \zeta_B, \overline{\zeta_A}, \overline{\zeta_B}$ :

$$\left\{4\nabla^2 + \frac{\lambda}{r} - \alpha^4\right\}\psi = \left\{4(\partial_A\partial_{\bar{A}} + \partial_B\partial_{\bar{B}}) + \lambda - \alpha^4(\zeta_A\overline{\zeta_A} + \zeta_B\overline{\zeta_B})\right\}\psi = 0 \quad (4)$$

Since  $\psi(x, y, z)$  independent of  $\sigma$ ,

$$\frac{\partial\psi}{\partial\sigma} = 0 \quad \Longleftrightarrow \quad (\overline{\zeta_A}\partial_{\bar{A}} - \zeta_A\partial_A)\psi = -(\overline{\zeta_B}\partial_{\bar{B}} - \zeta_B\partial_B)\psi. \quad (5)$$

# Where are the harmonic oscillators?

SE in terms of  $\zeta_A, \zeta_B, \overline{\zeta_A}, \overline{\zeta_B}$ :

$$\left\{4\nabla^2 + \frac{\lambda}{r} - \alpha^4\right\}\psi = \left\{4(\partial_A\partial_{\bar{A}} + \partial_B\partial_{\bar{B}}) + \lambda - \alpha^4(\zeta_A\overline{\zeta_A} + \zeta_B\overline{\zeta_B})\right\}\psi = 0 \quad (4)$$

Since  $\psi(x, y, z)$  independent of  $\sigma$ ,

$$\frac{\partial\psi}{\partial\sigma} = 0 \quad \Longleftrightarrow \quad (\overline{\zeta_A}\partial_{\bar{A}} - \zeta_A\partial_A)\psi = -(\overline{\zeta_B}\partial_{\bar{B}} - \zeta_B\partial_B)\psi. \quad (5)$$

Together, (4) and (5) are equivalent to SE (2).

# Where are the harmonic oscillators?

Let  $\zeta_A = q_1 + iq_2$  and  $\zeta_B = q_3 + iq_4$ , then (4) is the equation for a 4D HO

$$[\partial_1^2 + \partial_2^2 + \partial_3^2 + \partial_4^2 + \lambda - \alpha^4(q_1^2 + q_2^2 + q_3^2 + q_4^2)]\psi = 0 \quad (6)$$

# Where are the harmonic oscillators?

Let  $\zeta_A = q_1 + iq_2$  and  $\zeta_B = q_3 + iq_4$ , then (4) is the equation for a 4D HO

$$[\partial_1^2 + \partial_2^2 + \partial_3^2 + \partial_4^2 + \lambda - \alpha^4(q_1^2 + q_2^2 + q_3^2 + q_4^2)]\psi = 0 \quad (6)$$

with frequency  $\omega$  and energy  $\epsilon$  given by (1):

$$\alpha^2 \equiv \sqrt{-\frac{8E}{e^2 a}} = \frac{m\omega}{\hbar} \quad \lambda \equiv \frac{8}{a} = \frac{2m\epsilon}{\hbar^2}.$$

# Where are the harmonic oscillators?

Let  $\zeta_A = q_1 + iq_2$  and  $\zeta_B = q_3 + iq_4$ , then (4) is the equation for a 4D HO

$$[\partial_1^2 + \partial_2^2 + \partial_3^2 + \partial_4^2 + \lambda - \alpha^4(q_1^2 + q_2^2 + q_3^2 + q_4^2)]\psi = 0 \quad (6)$$

with frequency  $\omega$  and energy  $\epsilon$  given by (1):

$$\alpha^2 \equiv \sqrt{-\frac{8E}{e^2 a}} = \frac{m\omega}{\hbar} \quad \lambda \equiv \frac{8}{a} = \frac{2m\epsilon}{\hbar^2}.$$

Condition (5) becomes

$$(q_1\partial_2 - q_2\partial_1)\psi = -(q_3\partial_4 - q_4\partial_3)\psi. \quad (7)$$

(6)+(7): **Two 2D HO's with equal and opposite angular momenta**

# From harmonic oscillators to hydrogen

Separating variables  $\psi = \psi(q_1, q_2)\psi(q_3, q_4)$ ,

$$[\partial_1^2 + \partial_2^2 + \lambda_A - \alpha^4(q_1^2 + q_2^2)]\psi_A = 0,$$

with  $\lambda_A = 2m\epsilon_A/\hbar^2$ .

# From harmonic oscillators to hydrogen

Separating variables  $\psi = \psi(q_1, q_2)\psi(q_3, q_4)$ ,

$$[\partial_1^2 + \partial_2^2 + \lambda_A - \alpha^4(q_1^2 + q_2^2)]\psi_A = 0,$$

with  $\lambda_A = 2m\epsilon_A/\hbar^2$ . Solution for  $A$ :

$$\psi_{An_A m_A} = C_{n_A m_A} \left( \frac{\zeta_A}{\bar{\zeta}_A} \right)^{m_A/2} (\alpha^2 \zeta_A \bar{\zeta}_A)^{|m_A|/2} e^{-\frac{\alpha^2 \zeta_A \bar{\zeta}_A}{2}} L_{n_A+|m_A|}^{|m_A|} (\alpha^2 \zeta_A \bar{\zeta}_A)$$

$$n_A = 0, 1, 2, \dots \quad m_A = 0, \pm 1, \pm 2, \dots$$

**Energy:**  $\epsilon_{An_A m_A} = \hbar\omega(2n_A + |m_A| + 1) = \frac{\hbar^2 \lambda_{An_A m_A}}{2m}$

**Angular momentum:**  $L_{An_A m_A} = m_A \hbar$

Similar solution for  $B$ .  $\lambda_A + \lambda_B = \lambda$  and  $m_A = -m_B = m$  due to (7).



# From harmonic oscillators to hydrogen

Full solution

$$\psi_{n_A n_B m} = \psi_{A n_A m}(\zeta_A, \overline{\zeta_A}) \psi_{B n_B - m}(\zeta_B, \overline{\zeta_B}).$$

# From harmonic oscillators to hydrogen

Full solution

$$\psi_{n_A n_B m} = \psi_{A n_A m}(\zeta_A, \overline{\zeta_A}) \psi_{B n_B - m}(\zeta_B, \overline{\zeta_B}).$$

Can relate this back to the hydrogen atom.

# From harmonic oscillators to hydrogen

Full solution

$$\psi_{n_A n_B m} = \psi_{A n_A m}(\zeta_A, \overline{\zeta_A}) \psi_{B n_B - m}(\zeta_B, \overline{\zeta_B}).$$

Can relate this back to the hydrogen atom. From

$$\lambda = \lambda_A + \lambda_B = 4\alpha^2(n_A + n_B + |m| + 1) = \frac{8}{a}$$

can get energy in terms of  $n_A, n_B, m$ :

$$E = \frac{-\alpha^4 e^2 a}{8} = -\frac{\alpha^4 e^2}{\lambda} = \frac{-e^2}{2a(n_A + n_B + |m| + 1)^2} \equiv \frac{-e^2}{2aN^2}.$$

# From harmonic oscillators to hydrogen

How about the wavefunctions?

# From harmonic oscillators to hydrogen

How about the wavefunctions? Going to parabolic coordinates  $(\xi, \eta, \varphi)$ :

$$\begin{aligned}x &= \sqrt{\xi\eta} \cos \varphi & y &= \sqrt{\xi\eta} \sin \varphi & z &= (\xi - \eta)/2 \\ \iff \xi &= 2r \cos^2(\theta/2) = 2|\zeta_A|^2 & \eta &= 2r \sin^2(\theta/2) = 2|\zeta_B|^2.\end{aligned}$$

# From harmonic oscillators to hydrogen

How about the wavefunctions? Going to parabolic coordinates  $(\xi, \eta, \varphi)$ :

$$\begin{aligned}x &= \sqrt{\xi\eta} \cos \varphi & y &= \sqrt{\xi\eta} \sin \varphi & z &= (\xi - \eta)/2 \\ \iff \xi &= 2r \cos^2(\theta/2) = 2|\zeta_A|^2 & \eta &= 2r \sin^2(\theta/2) = 2|\zeta_B|^2.\end{aligned}$$

we get

$$\psi_{n_A n_B m} = K_{n_A n_B m} e^{im\varphi} (\xi\eta)^{|m|/2} e^{-\frac{\alpha^2(\xi^2 + \eta^2)}{4}} L_{n_A+|m|}^{|m|} \left( \frac{\alpha^2 \xi}{2} \right) L_{n_B+|m|}^{|m|} \left( \frac{\alpha^2 \eta}{2} \right).$$

# From harmonic oscillators to hydrogen

How about the wavefunctions? Going to parabolic coordinates  $(\xi, \eta, \varphi)$ :

$$\begin{aligned}x &= \sqrt{\xi\eta} \cos \varphi & y &= \sqrt{\xi\eta} \sin \varphi & z &= (\xi - \eta)/2 \\ \iff \xi &= 2r \cos^2(\theta/2) = 2|\zeta_A|^2 & \eta &= 2r \sin^2(\theta/2) = 2|\zeta_B|^2.\end{aligned}$$

we get

$$\psi_{n_A n_B m} = K_{n_A n_B m} e^{im\varphi} (\xi\eta)^{|m|/2} e^{-\frac{\alpha^2(\xi^2 + \eta^2)}{4}} L_{n_A+|m|}^{|m|} \left( \frac{\alpha^2 \xi}{2} \right) L_{n_B+|m|}^{|m|} \left( \frac{\alpha^2 \eta}{2} \right).$$

These are simultaneous eigenfunctions of  $\mathbf{H}$ ,

# From harmonic oscillators to hydrogen

How about the wavefunctions? Going to parabolic coordinates  $(\xi, \eta, \varphi)$ :

$$\begin{aligned}x &= \sqrt{\xi\eta} \cos \varphi & y &= \sqrt{\xi\eta} \sin \varphi & z &= (\xi - \eta)/2 \\ \iff \xi &= 2r \cos^2(\theta/2) = 2|\zeta_A|^2 & \eta &= 2r \sin^2(\theta/2) = 2|\zeta_B|^2.\end{aligned}$$

we get

$$\psi_{n_A n_B m} = K_{n_A n_B m} e^{im\varphi} (\xi\eta)^{|m|/2} e^{-\frac{\alpha^2(\xi^2 + \eta^2)}{4}} L_{n_A+|m|}^{|m|} \left( \frac{\alpha^2 \xi}{2} \right) L_{n_B+|m|}^{|m|} \left( \frac{\alpha^2 \eta}{2} \right).$$

These are simultaneous eigenfunctions of  $\mathbf{H}$ ,  $\mathbf{L}_z$ ,



# From harmonic oscillators to hydrogen

How about the wavefunctions? Going to parabolic coordinates  $(\xi, \eta, \varphi)$ :

$$\begin{aligned}x &= \sqrt{\xi\eta} \cos \varphi & y &= \sqrt{\xi\eta} \sin \varphi & z &= (\xi - \eta)/2 \\ \iff \xi &= 2r \cos^2(\theta/2) = 2|\zeta_A|^2 & \eta &= 2r \sin^2(\theta/2) = 2|\zeta_B|^2.\end{aligned}$$

we get

$$\psi_{n_A n_B m} = K_{n_A n_B m} e^{im\varphi} (\xi\eta)^{|m|/2} e^{-\frac{\alpha^2(\xi^2 + \eta^2)}{4}} L_{n_A+|m|}^{|m|} \left( \frac{\alpha^2 \xi}{2} \right) L_{n_B+|m|}^{|m|} \left( \frac{\alpha^2 \eta}{2} \right).$$

These are simultaneous eigenfunctions of  $\mathbf{H}$ ,  $\mathbf{L}_z$ , and  $\mathbf{M}_z$  where

$$\mathbf{M} = \frac{1}{2m} (\mathbf{p} \times \mathbf{L} - \mathbf{L} \times \mathbf{p}) - \frac{e^2}{r} \mathbf{r}.$$

is the **Laplace-Runge-Lenz operator**, symmetrized by Pauli, 1926.

# From harmonic oscillators to hydrogen

From how  $(\zeta_A, \zeta_B)$  is defined:

$$\mathbf{M}_z = \frac{e^2 a}{r} \left[ |\zeta_B|^2 \partial_A \partial_{\bar{A}} - |\zeta_A|^2 \partial_B \partial_{\bar{B}} - \frac{1}{a} (|\zeta_A|^2 + |\zeta_B|^2) \right].$$

# From harmonic oscillators to hydrogen

From how  $(\zeta_A, \zeta_B)$  is defined:

$$\mathbf{M}_z = \frac{e^2 a}{r} \left[ |\zeta_B|^2 \partial_A \partial_{\bar{A}} - |\zeta_A|^2 \partial_B \partial_{\bar{B}} - \frac{1}{a} (|\zeta_A|^2 + |\zeta_B|^2) \right].$$

CSCO is  $\{\mathbf{H}, \mathbf{L}_z, \mathbf{M}_z\}$  instead of  $\{\mathbf{H}, \mathbf{L}^2, \mathbf{L}_z\}$ . Eigenvalue equations:

$$\mathbf{H} \psi_{n_A n_B m} = \frac{-e^2}{2a N^2} \psi_{n_A n_B m}$$

$$\mathbf{L}_z \psi_{n_A n_B m} = m \hbar \psi_{n_A n_B m}$$

$$\mathbf{M}_z \psi_{n_A n_B m} = \frac{e^2 (n_B - n_A)}{N} \psi_{n_A n_B m}.$$

# From harmonic oscillators to hydrogen

From how  $(\zeta_A, \zeta_B)$  is defined:

$$\mathbf{M}_z = \frac{e^2 a}{r} \left[ |\zeta_B|^2 \partial_A \partial_{\bar{A}} - |\zeta_A|^2 \partial_B \partial_{\bar{B}} - \frac{1}{a} (|\zeta_A|^2 + |\zeta_B|^2) \right].$$

CSCO is  $\{\mathbf{H}, \mathbf{L}_z, \mathbf{M}_z\}$  instead of  $\{\mathbf{H}, \mathbf{L}^2, \mathbf{L}_z\}$ . Eigenvalue equations:

$$\mathbf{H} \psi_{n_A n_B m} = \frac{-e^2}{2a N^2} \psi_{n_A n_B m}$$

$$\mathbf{L}_z \psi_{n_A n_B m} = m \hbar \psi_{n_A n_B m}$$

$$\mathbf{M}_z \psi_{n_A n_B m} = \frac{e^2 (n_B - n_A)}{N} \psi_{n_A n_B m}.$$

$$[\mathbf{M}_z, \mathbf{L}^2] \neq 0, \quad [\mathbf{M}_z, \mathbf{M}^2] \neq 0, \quad [\mathbf{M}^2, \mathbf{H}] = [\mathbf{M}^2, \mathbf{L}^2] = [\mathbf{M}^2, \mathbf{L}_z] = 0.$$

## Aside: Hydrogen wavefunctions in parabolic coordinates


What do eigenfunctions of  $\mathbf{M}_z$  look like?

$$\psi_{n_A n_B m} = K_{n_A n_B m} e^{im\varphi} (\xi\eta)^{|m|/2} e^{-\frac{\alpha^2(\xi^2+\eta^2)}{4}} L_{n_A+|m|}^{|m|}\left(\frac{\alpha^2\xi}{2}\right) L_{n_B+|m|}^{|m|}\left(\frac{\alpha^2\eta}{2}\right).$$


# Aside: Hydrogen wavefunctions in parabolic coordinates

What do eigenfunctions of  $\mathbf{M}_z$  look like?


$$\psi_{n_A n_B m} = K_{n_A n_B m} e^{im\varphi} (\xi\eta)^{|m|/2} e^{-\frac{\alpha^2(\xi^2+\eta^2)}{4}} L_{n_A+|m|}^{|m|}\left(\frac{\alpha^2\xi}{2}\right) L_{n_B+|m|}^{|m|}\left(\frac{\alpha^2\eta}{2}\right).$$




(0,0,0)



(2,0,0)



(2,1,2)



(4,1,3)

Figure:  $|\psi_{n_A n_B m}(x, 0, z)|^2$  for different values of  $(n_A, n_B, m)$  [13]

# Shouldn't the correspondence be classical?

Following [14], use the Kustaanheimo-Stiefel transformation  $S : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ :

$$x_1 = 2(s_1 s_3 - s_2 s_4)$$

$$x_2 = 2(s_1 s_4 + s_2 s_3)$$

$$x_3 = s_1^2 + s_2^2 - s_3^2 - s_4^2$$

# Shouldn't the correspondence be classical?

Following [14], use the Kustaanheimo-Stiefel transformation  $S : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ :

$$x_1 = 2(s_1 s_3 - s_2 s_4)$$

$$x_2 = 2(s_1 s_4 + s_2 s_3)$$

$$x_3 = s_1^2 + s_2^2 - s_3^2 - s_4^2$$

Only three of  $\{s_1, s_2, s_3, s_4\}$  are independent. What is the constraint?



# Shouldn't the correspondence be classical?

Following [14], use the Kustaanheimo-Stiefel transformation  $S : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ :

$$x_1 = 2(s_1 s_3 - s_2 s_4)$$

$$x_2 = 2(s_1 s_4 + s_2 s_3)$$

$$x_3 = s_1^2 + s_2^2 - s_3^2 - s_4^2$$

Only three of  $\{s_1, s_2, s_3, s_4\}$  are independent. What is the constraint?

$$\begin{cases} x_1 = r \sin \theta \cos \phi \\ x_2 = r \sin \theta \sin \phi \\ x_3 = r \cos \theta \end{cases} \quad \begin{cases} s_1 = s \cos \alpha \cos \beta \\ s_2 = s \cos \alpha \sin \beta \\ s_3 = s \sin \alpha \cos \gamma \\ s_4 = s \sin \alpha \sin \gamma \end{cases}$$

# Shouldn't the correspondence be classical?

Following [14], use the Kustaanheimo-Stiefel transformation  $S : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ :

$$x_1 = 2(s_1 s_3 - s_2 s_4)$$

$$x_2 = 2(s_1 s_4 + s_2 s_3)$$

$$x_3 = s_1^2 + s_2^2 - s_3^2 - s_4^2$$

Only three of  $\{s_1, s_2, s_3, s_4\}$  are independent. What is the constraint?

$$\begin{cases} x_1 = r \sin \theta \cos \phi \\ x_2 = r \sin \theta \sin \phi \\ x_3 = r \cos \theta \end{cases} \quad \begin{cases} s_1 = s \cos \alpha \cos \beta \\ s_2 = s \cos \alpha \sin \beta \\ s_3 = s \sin \alpha \cos \gamma \\ s_4 = s \sin \alpha \sin \gamma \end{cases}$$

Constraint on coordinates:  $r = s^2$ ,  $\theta = 2\alpha$ ,  $\phi = \beta + \gamma$

# Shouldn't the correspondence be classical?

Following [14], use the Kustaanheimo-Stiefel transformation  $S : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ :

$$x_1 = 2(s_1 s_3 - s_2 s_4)$$

$$x_2 = 2(s_1 s_4 + s_2 s_3)$$

$$x_3 = s_1^2 + s_2^2 - s_3^2 - s_4^2$$

Only three of  $\{s_1, s_2, s_3, s_4\}$  are independent. What is the constraint?

$$\begin{cases} x_1 = r \sin \theta \cos \phi \\ x_2 = r \sin \theta \sin \phi \\ x_3 = r \cos \theta \end{cases} \quad \begin{cases} s_1 = s \cos \alpha \cos \beta \\ s_2 = s \cos \alpha \sin \beta \\ s_3 = s \sin \alpha \cos \gamma \\ s_4 = s \sin \alpha \sin \gamma \end{cases}$$

Constraint on coordinates:  $r = s^2$ ,  $\theta = 2\alpha$ ,  $\phi = \beta + \gamma$

Constraint on velocities:  $s_2 \dot{s}_1 - s_1 \dot{s}_2 - s_4 \dot{s}_3 + s_3 \dot{s}_4 = 0$

# Shouldn't the correspondence be classical?

The equivalence: With  $4|E| = m\omega^2/2$  and  $\epsilon = 4k$ ,

$$\frac{1}{2}mv^2 - \frac{k}{r} = -|E| \quad \longrightarrow \quad \frac{1}{2}m\dot{s}^2 + \frac{1}{2}m\omega^2 s^2 = \epsilon \quad (4D \text{ H.O.})$$

# Shouldn't the correspondence be classical?

The equivalence: With  $4|E| = m\omega^2/2$  and  $\epsilon = 4k$ ,

$$\frac{1}{2}mv^2 - \frac{k}{r} = -|E| \quad \longrightarrow \quad \frac{1}{2}m\dot{s}^2 + \frac{1}{2}m\omega^2 s^2 = \epsilon \quad (4D \text{ H.O.})$$

In polar coordinates:

# Shouldn't the correspondence be classical?

The equivalence: With  $4|E| = m\omega^2/2$  and  $\epsilon = 4k$ ,

$$\frac{1}{2}mv^2 - \frac{k}{r} = -|E| \quad \longrightarrow \quad \frac{1}{2}m\dot{s}^2 + \frac{1}{2}m\omega^2 s^2 = \epsilon \quad (4D \text{ H.O.})$$

In polar coordinates:  $u = s \cos \alpha$ ,  $v = s \sin \alpha$

$$\frac{1}{2}m(\dot{u}^2 + u^2\dot{\beta}^2 + \dot{v}^2 + v^2\dot{\gamma}^2) + \frac{1}{2}m\omega^2(u^2 + v^2) = \epsilon.$$

# Shouldn't the correspondence be classical?

The equivalence: With  $4|E| = m\omega^2/2$  and  $\epsilon = 4k$ ,

$$\frac{1}{2}mv^2 - \frac{k}{r} = -|E| \quad \longrightarrow \quad \frac{1}{2}m\dot{s}^2 + \frac{1}{2}m\omega^2 s^2 = \epsilon \quad (4D \text{ H.O.})$$

In polar coordinates:  $u = s \cos \alpha$ ,  $v = s \sin \alpha$

$$\frac{1}{2}m(\dot{u}^2 + u^2\dot{\beta}^2 + \dot{v}^2 + v^2\dot{\gamma}^2) + \frac{1}{2}m\omega^2(u^2 + v^2) = \epsilon.$$

The constraint on velocities  $\iff \vec{L} \cdot \vec{M} = 0$

# Shouldn't the correspondence be classical?

The equivalence: With  $4|E| = m\omega^2/2$  and  $\epsilon = 4k$ ,

$$\frac{1}{2}mv^2 - \frac{k}{r} = -|E| \quad \longrightarrow \quad \frac{1}{2}m\dot{s}^2 + \frac{1}{2}m\omega^2 s^2 = \epsilon \quad (4D \text{ H.O.})$$

In polar coordinates:  $u = s \cos \alpha$ ,  $v = s \sin \alpha$

$$\frac{1}{2}m(\dot{u}^2 + u^2\dot{\beta}^2 + \dot{v}^2 + v^2\dot{\gamma}^2) + \frac{1}{2}m\omega^2(u^2 + v^2) = \epsilon.$$

The constraint on velocities  $\iff \vec{L} \cdot \vec{M} = 0$  and implies

$$-u^2\dot{\beta} + v^2\dot{\gamma} = 0 \implies mu^2\dot{\beta} = mv^2\dot{\gamma} \implies L_{\beta} = L_{\gamma}$$



# Shouldn't the correspondence be classical?

The equivalence: With  $4|E| = m\omega^2/2$  and  $\epsilon = 4k$ ,

$$\frac{1}{2}mv^2 - \frac{k}{r} = -|E| \quad \longrightarrow \quad \frac{1}{2}m\dot{s}^2 + \frac{1}{2}m\omega^2 s^2 = \epsilon \quad (4D \text{ H.O.})$$

In polar coordinates:  $u = s \cos \alpha$ ,  $v = s \sin \alpha$

$$\frac{1}{2}m(\dot{u}^2 + u^2\dot{\beta}^2 + \dot{v}^2 + v^2\dot{\gamma}^2) + \frac{1}{2}m\omega^2(u^2 + v^2) = \epsilon.$$


The constraint on velocities  $\iff \vec{L} \cdot \vec{M} = 0$  and implies

$$-u^2\dot{\beta} + v^2\dot{\gamma} = 0 \implies mu^2\dot{\beta} = mv^2\dot{\gamma} \implies L_{\beta} = L_{\gamma}$$

$\implies$  Two coupled 2D H.O.'s with equal angular momenta

# Something deeper?

Symmetry implies degeneracy.  $[\mathbf{M}, \mathbf{H}] = 0$  explains the  $n^2$  degeneracy in  $H$ .



More info:  $SO(4)$  symmetry of  $H$ , etc. See Chapter 14 of [15].

# References I

- [1] Galliano Valent.  
The hydrogen atom in electric and magnetic fields: Pauli's 1926 article.  
*American Journal of Physics*, 71(2):171–175, 2003.
- [2] AA Stahlhofen.  
Pauli and the runge–lenz vector.  
*American Journal of Physics*, 72(1):10–10, 2004.
- [3] Herbert Goldstein.  
Prehistory of the "runge–lenz" vector.  
*American Journal of Physics*, 43(8):737–738, 1975.
- [4] Herbert Goldstein.  
More on the prehistory of the laplace or runge–lenz vector.  
*Am. J. Phys*, 44(11):1123–1124, 1976.
- [5] Herbert Goldstein, Charles Poole, and John Safko.  
Classical mechanics, 2002.
- [6] Carl Runge.  
*Vektoranalysis*.  
S. Hirzel, 1919.
- [7] Pierre Simon Laplace.  
*Traité de mécanique céleste*, 1.  
Typ. Crapelet, 1823.
- [8] Wilhelm Lenz.  
Über den bewegungsverlauf und die quantenzustände der gestörten keplerbewegung.  
*Zeitschrift für Physik*, 24(1):197–207, 1924.

# References II

- [9] William Rowan Hamilton.  
On the application of the method of quaternions to some dynamical questions.  
In *Proc. Roy. Irish Acad*, volume 3, pages 441–448, 1847.
- [10] Jakob Hermann.  
Unknown title.  
*Giornale de Letterati D'Italia*, 2:447–467, 1710.
- [11] W Pauli Jr.  
Über das wasserstoffspektrum vom standpunkt der neuen quantenmechanik.  
*Zeitschrift für Physik A Hadrons and nuclei*, 36(5):336–363, 1926.
- [12] FHJ Cornish.  
The hydrogen atom and the four-dimensional harmonic oscillator.  
*Journal of Physics A: Mathematical and General*, 17(2):323, 1984.
- [13] S. M. Blinder.  
The hydrogen atom in parabolic coordinates, 2011.
- [14] Augustine C Chen.  
Coulomb–kepler problem and the harmonic oscillator.  
*American Journal of Physics*, 55(3):250–252, 1987.
- [15] Robert Gilmore.  
*Lie groups, physics, and geometry: an introduction for physicists, engineers and chemists*.  
Cambridge University Press, 2008.