Hydrodynamic Properties of the Unitary Fermi Gas

Eric Wolf, Huan Q Bui, Parth B Patel, Zhenjie Yan, Carsten Robens, Richard Fletcher, Martin Zwierlein

MIT-Harvard Center for Ultracold Atoms, Research Laboratory of Electronics,
Massachusetts Institute of Technology, Cambridge, MA 02139

Unitary Fermi Gas in a Box Potential

Unitary Fermi Gas

- Strongly-interacting fermion systems - difficult to analyze a priori
- Relevant to systems ranging from neutron stars to high-T_c superconductors
- Unitary Fermi gas is scale-invariant
- Realize unitarity with $|1\rangle - |3\rangle$ Feshbach resonance in ^{6}Li
- Evaporatively cool spin mixture to below T_F

Box Potential [1]

- Hollow blue-detuned beams realize (quasi) flat potential
- Reduces influence of trap averaging & targets smaller range of densities
- Residual harmonic trap in axial dimension allows momentum-space imaging

Temperature Response – Second Sound [3]

Second Sound

- Temperature oscillations: out-of-phase oscillations of normal and superfluid phase
- Excite oscillations by driving high-frequency sound waves to locally heat the gas
- Measure temperature locally using RF spectroscopy
- Second sound is only present below the superfluid transition temperature
- Damping rate of second sound gives thermal conductivity

Density Response – First Sound [2]

First Sound

- Density oscillations: in-phase oscillation of normal and superfluid phase
- Excite by shaking box walls
- Image density waves in-situ – extract wavevector k for a given ω to get c
- Speed of sound in scale-invariant system given by system energy

$$mc^2 = \frac{10}{9}\frac{E}{N}$$

- Agrees with experiment results

Resonant Modes and Dissipation

- Sound speed constrained by scale invariance, but dissipation is not
- Hydrodynamics predicts damping rate $\Gamma \propto k^2$, with proportionality constant D_s, the sound diffusivity
- D_s depends on shear viscosity and thermal conductivity
- Measure damping vs. k using width of resonant box modes – extract D_s

Hydrodynamics – Transport Properties

Hydrodynamic Quantities

- Hydrodynamic quantities above T_c: shear viscosity η and thermal conductivity κ
- Bulk viscosity vanishes for scale-invariant systems; below T_c, superfluidity density is conserved
- First sound dissipation given by κ and η; second sound dissipation by κ and c_P
- Together with equation of state [4], first and second sound give η and ρ above T_c

Results

- Viscosity in reasonable agreement with prior results and theory
- Thermal conductivity differs strongly from theory predictions near T_c
- Prandtl number $\frac{c_P\eta}{\kappa}$, ratio of momentum to thermal diffusivity, remains close to typical $2/3$ value for gases

Red diamond [5], green line [6], pink line [7], cyan line [8], orange line [9]

References

[1] B. Mukherjee et al., *PRL* 2017
[2] P.B. Patel et al., *Science* 2020
[3] Z. Yan et al., in preparation
[4] M.J.H. Ku et al., *Science* 2012
[5] L. Baird et al., *PRL* 2019
[6] B. Frank et al., *Phys Rev Research* 2020
[7] J. A. Joseph et al., *PRL* 2015
[8] M. Bluhm et al., *PRL* 2017
[9] T. Enss et al., *Annals of Physics* 2011

Funding

AFOSR-MURI, National Science Foundation, Office of Naval Research, Vannevar Bush Faculty Fellowship