

Hydrodynamic Properties of the Unitary Fermi Gas



Eric Wolf, Huan Q Bui, Parth B Patel, Zhenjie Yan, Carsten Robens, Richard Fletcher, Martin Zwierlein

MIT-Harvard Center for Ultracold Atoms, Research Laboratory of Electronics, Massachusetts Institute of Technology, Cambridge, MA 02139


Unitary Fermi Gas in a Box Potential


Unitary Fermi Gas

- Strongly-interacting fermion systems - difficult to analyze a priori
- Relevant to systems ranging from neutron stars to high-Tc superconductors
- Unitary Fermi gas is scale-invariant
- Realize unitarity with |1 > -|3 >Feshbach resonance in ⁶Li.
- Evaporatively cool spin mixture to below T_F

Box Potential [1]

- Hollow blue-detuned beams realize (quasi) flat potential
- Reduces influence of trap averaging & targets smaller range of densities
- Residual harmonic trap in axial dimension allows momentum-space imaging





700

B (G)

900

-20000


500


600


Temperature Response –Second Sound [3]


Second Sound

- Temperature oscillations: out-of-phase oscillations of normal and superfluid phase
- Excite oscillations by driving high-frequency sound waves to locally heat the gas
- Measure temperature locally using RF spectroscopy
- Second sound is only present below the superfluid transition temperature
- Damping rate of second sound gives thermal conductivity











Density Response – First Sound [2]


First Sound

- Density oscillations: in-phase oscillation of normal and superfluid phase
- Excite by shaking box walls
- Image density waves in situ extract wavevector k for a given ω to get c
- Speed of sound in scale-invariant system given by system energy: $mc^2 = \frac{10 E}{}$
- Agrees with experimental results.

1st Sound NF








Resonant Modes and Dissipation


- Sound speed constrained by scale invariance, but dissipation is not
- Hydrodynamics predicts damping rate $\Gamma \propto k^2$, with proportionality constant $D_{\rm S}$, the sound diffusivity D_{S} depends on shear
- viscosity and thermal conductivity
- Measure damping vs. k using width of resonant box modes – extract D_s



Hydrodynamics – Transport Properties


Hydrodynamic Quantities

- Hydrodynamic quantities above T_c : shear viscosity η and thermal conductivity κ
- Bulk viscosity vanishes for scale invariant system; below T_c , superfluid density is conserved
- First sound dissipation given by κ and η ; second sound dissipation by κ and c_P
- Together with equation of state [4], first and second sound give η and ρ above T_c



Results

- Viscosity in reasonable agreement with prior results and theory
- Thermal conductivity differs strongly from theory predictions near T_c
- Prandtl number $\frac{c_P\eta}{\kappa}$, ratio of momentum to thermal diffusivity, remains close to typical 2/3 value for gases



Red diamond [5], green line [6], pink line [7], cyan line [8], orange line [9]

References

- [1] B. Mukherjee et al., PRL 2017
- [2] P.B. Patel et al., Science 2020 [3] Z. Yan et al., in preparation

[4] M.J.H. Ku et al., *Science* 2012

- [5] L. Baird et al., *PRL* 2019
- [6] B. Frank et al., Phys Rev Research 2020
- [7] J. A. Joseph et al., *PRL* 2015
- [8] M. Bluhm et al., PRL 2017
- [9] T. Enss et al., Annals of Physics 2011

Funding