

Name: **Huan Q. Bui**
Course: **8.422 - AMO II**
Problem set: **#3**
Due: Friday, Mar 3, 2022.

1. Classical Coherence of Light.

Consider a classical light field. The classical expressions for first-order and second-order coherence are:

$$g^{(1)}(\tau) = \frac{\langle E^*(t)E(t+\tau) \rangle}{\langle E^*(t)E(t) \rangle}$$

$$g^{(2)}(\tau) = \frac{\langle I(t)I(t+\tau) \rangle}{\langle I(t) \rangle^2} = \frac{\langle E^*(t)E^*(t+\tau)E(t+\tau)E(t) \rangle}{\langle E^*(t)E(t) \rangle^2}$$

where $\langle \cdot \rangle$ denotes a statistical averaging over many measurements, which can be interpreted as a time average. Moreover, $\langle f^*(t)g(t) \rangle$ defines a scalar product.

- a) Here we show that $|g^{(1)}(\tau)| \leq 1$. In view of Cauchy's inequality, and the fact that $\langle E^*(t+\tau)E(t+\tau) \rangle = \langle E^*(t)E(t) \rangle$

$$|g^{(1)}(\tau)|^2 = \frac{|\langle E^*(t)E(t+\tau) \rangle|^2}{|\langle E^*(t)E(t) \rangle|^2} \leq \frac{\langle E^*(t)E(t) \rangle \langle E^*(t+\tau)E(t+\tau) \rangle}{|\langle E^*(t)E(t) \rangle|^2} = \frac{\langle E^*(t)E(t) \rangle \langle E^*(t)E(t) \rangle}{|\langle E^*(t)E(t) \rangle|^2} = 1.$$

So, $|g^{(1)}(\tau)| \leq 1$ as desired.

- b) Here we show that for zero time-delay, the second-order coherence obeys $g^{(2)}(0) \geq 1$. To this end, we notice that

$$0 \leq \langle [I(t) - \langle I(t) \rangle]^2 \rangle = \langle I(t)^2 - 2I(t)\langle I(t) \rangle + \langle I(t) \rangle^2 \rangle = \langle I(t)^2 \rangle - 2\langle I(t) \rangle^2 + \langle I(t) \rangle^2 = \langle I(t)^2 \rangle - \langle I(t) \rangle^2.$$

Thus, for $\tau = 0$, we have

$$g^{(2)}(0) = \frac{\langle I(t)^2 \rangle}{\langle I(t) \rangle^2} \geq \frac{\langle I(t) \rangle^2}{\langle I(t) \rangle^2} = 1,$$

as desired. This implies that light in a number state, which has $g^{(2)}(0) < 1$, has no classical analog.

- c) Here we show that $g^{(2)}(\tau) \leq g^{(2)}(0)$. Viewing $\langle I(t)I(t+\tau) \rangle$ as the inner product $\langle I(t), I(t+\tau) \rangle$ (which is possible since intensities are real numbers), we see immediately that $\langle I(t)I(t+\tau) \rangle \leq \langle I(t)I(t) \rangle$, which gives

$$g^{(2)}(\tau) = \frac{\langle I(t)I(t+\tau) \rangle}{\langle I(t) \rangle^2} \leq \frac{\langle I(t)I(t) \rangle}{\langle I(t) \rangle^2} = g^{(2)}(0).$$

This implies that anti-bunched light, which has $g^{(2)}(\tau) > g^{(2)}(0)$, has no classical analog.

- d) Consider chaotic classical light generated by an ensemble of ν atoms. The total electric field can be expressed as $E(t) = \sum_{i=1}^{\nu} E_i(t)$, where the phases of E_i are random. Here we show that when ν is large,

$$g^{(2)}(\tau) = 1 + |g^{(1)}(\tau)|^2.$$

Since statistical averages in which the random electric field phases do not cancel are zero, we have

$$\langle E^*(t)E(t+\tau) \rangle =$$

2. Quantum Coherence of Light.

Consider light in a single mode of the radiation field. The quantum mechanical expressions for first-order and second-order coherence are

$$g^{(1)}(\mathbf{r}_1, t_1, \mathbf{r}_2, t_2) = \frac{\langle \hat{E}^-(\mathbf{r}_1, t_1) \hat{E}^+(\mathbf{r}_2, t_2) \rangle}{\langle \hat{E}^-(\mathbf{r}_1, t_1) \hat{E}^+(\mathbf{r}_1, t_1) \rangle^{1/2} \langle \hat{E}^-(\mathbf{r}_2, t_2) \hat{E}^+(\mathbf{r}_2, t_2) \rangle^{1/2}}$$

$$g^{(2)}(\mathbf{r}_1, t_1, \mathbf{r}_2, t_2; \mathbf{r}_2, t_2, \mathbf{r}_1, t_1) = \frac{\langle \hat{E}^-(\mathbf{r}_1, t_1) \hat{E}^-(\mathbf{r}_2, t_2) \hat{E}^+(\mathbf{r}_2, t_2) \hat{E}^+(\mathbf{r}_1, t_1) \rangle}{\langle \hat{E}^-(\mathbf{r}_1, t_1) \hat{E}^+(\mathbf{r}_1, t_1) \rangle \langle \hat{E}^-(\mathbf{r}_2, t_2) \hat{E}^+(\mathbf{r}_2, t_2) \rangle}$$

where

$$\hat{E}^+(\mathbf{r}, t) = i \sqrt{\frac{\hbar \omega}{2 \epsilon_0 V}} a e^{-i(\omega t - \mathbf{k} \cdot \mathbf{r})}$$

$$\hat{E}^-(\mathbf{r}, t) = -i \sqrt{\frac{\hbar \omega}{2 \epsilon_0 V}} a^\dagger e^{+i(\omega t - \mathbf{k} \cdot \mathbf{r})}$$

a) Here we show that

$$g^{(2)}(0) = \frac{\langle a^\dagger a^\dagger a a \rangle}{\langle a^\dagger a \rangle^2} = \frac{\langle n^2 \rangle - \langle n \rangle}{\langle n \rangle^2}$$

b) Next we show that for number states of light $|n\rangle$ where $n > 2$,

$$|g^{(1)}| = 1 \quad g^{(2)} = 1 - \frac{1}{n}$$

independent of space-time separation.

For $n = 0$, we calculate $g^{(1)}$ and $g^{(2)}$:

$$g_{n=0}^{(1)} =$$

$$g_{n=0}^{(2)} =$$

For $n = 1$, we calculate $g^{(1)}$ and $g^{(2)}$

$$g_{n=1}^{(1)} =$$

$$g_{n=1}^{(2)} =$$

c) Consider a coherent state $|\alpha\rangle$, we show that $|g^{(1)}| = |g^{(2)}| = 1$

$$|g_{|\alpha\rangle}^{(1)}| =$$

$$|g_{|\alpha\rangle}^{(2)}| =$$

d) Finally, consider chaotic light with density matrix

$$\hat{\rho} = \left(1 - e^{-\hbar\omega/k_B T}\right) \sum_n e^{-n\hbar\omega/k_B T} |n\rangle\langle n|.$$

Here we show that $|g^{(1)}| = 1$ and $|g^{(2)}| = 2$.

Compare these results to what we found in Problem 1... What about multi-mode chaotic light?

e) Consider the states

$$|\psi_{\pm}\rangle = \frac{|\alpha\rangle \pm |-\alpha\rangle}{\sqrt{2}\sqrt{1 \pm e^{-2|\alpha|^2}}}.$$

Let us compute $g^{(2)}(\tau)$ for these states as a function of α .

Do either of these two states show non-classical second-order coherence? Why (or why not)? (Make sure you agree with the normalization given)

3. The Quantum Beamsplitter.

Let the beamsplitter operator B , acting with angle \angle on modes a and b , by defined by

$$B = \exp \left[\theta \left(a^{\dagger} b - a b^{\dagger} \right) \right].$$

a) Here we show that B conserves photon number and leaves coherent states as coherent states.

b) Let $|\alpha\rangle$ be a coherent state. Here we compute $B |0\rangle_b |\alpha\rangle_a$.

From here, we see that the output is a tensor product of coherent states for all θ . This makes sense: the beamsplitter has well-defined transmission and reflection coefficients. What are these in terms of θ ?

c) There is close connection between the Lie group $SU(2)$ and the algebra of two coupled harmonic oscillators, which is useful for understanding B . Define

$$s_z = a^{\dagger} a - b^{\dagger} b \quad s_+ = a^{\dagger} a \quad s_- = b^{\dagger} b,$$

and let $s_{\pm} = (s_x \pm i s_y)/\sqrt{2}$.

What is $B(\theta)$ in spin space?

What is $a^{\dagger} a + b^{\dagger} b$ in spin space?

We now show that s_x, s_y, s_z have the same commutation relations as Pauli matrices.

How does this explain why $a^{\dagger} a + b^{\dagger} b$ is invariant under B . It is the Casimir operator of the algebra.

d) Here we look at how a beamsplitter transforms an input of a photon-number eigenstate. Let

$$B(\theta) = \exp \left[\theta \left(-a^{\dagger} b + a b^{\dagger} \right) \right],$$

and $B = B(\pi/4)$ be a 50/50 beamsplitter, such that

$$B a B^{\dagger} = \frac{a + b}{\sqrt{2}} \quad B b B^{\dagger} = \frac{-a + b}{\sqrt{2}}.$$

Let us compute $B |0\rangle_b |n\rangle_a$.

Note that the result is NOT $|n/2\rangle |n/2\rangle$ since $|n\rangle$ is a number state and not a coherent state.

What photon number states have the largest amplitude?

How sharp is the distribution for $n = 10$, and $n = 100$, or as a function of n , if a general solution exists? To do this, we look at the binomial expansion for $(a^{\dagger} + b^{\dagger})^n$.

4. The Hanbury-Brown and Twiss experiment and $g^{(2)}(\tau)$.

Here we look at how the HBT experiment measures $g^{(2)}(\tau)$. To this end, let $a, a^\dagger, b, b^\dagger$ be the raising and lowering operators for the two modes of light input to the beamsplitter, and let the unitary transformation performed by the beamsplitter be defined by

$$a_1 = UaU^\dagger = \frac{a+b}{\sqrt{2}} \quad b_1 = UbU^\dagger = \frac{a-b}{\sqrt{2}}.$$

For light input in state $|\psi_a\rangle$, suppose that the output of the coincidence circuit is a voltage

$$V_{\psi_a, 0_b} = V_0 \langle \psi_a, 0_b | a_1^\dagger a_1 b_1^\dagger b_1 | \psi_a, 0_b \rangle,$$

which is nothing but the average of the product of the two detected photon signals. We will show that $V_{\psi_a, 0_b}$ gives measure of $g^{(2)}(\tau)$ up to an additive offset and normalization.