

1 Ramsey Fringes Overview:

Following a double pulse, the population of the excited state is:

$$P_2 = 4 \sin^2 \theta \sin^2 \frac{\Omega' \tau}{2} \left\{ \cos \frac{\Omega' \tau}{2} \cos \frac{\Delta_0 T}{2} - \cos \theta \sin \frac{\Omega' \tau}{2} \sin \frac{\Delta_0 T}{2} \right\}$$

under the assumption that initially,

$$\begin{pmatrix} C_1(0) \\ C_2(0) \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

The final state vector is:

$$\begin{pmatrix} C_1(2\tau + T) \\ C_2(2\tau + T) \end{pmatrix} = \rho_2 D \rho_1 \begin{pmatrix} C_1(0) \\ C_2(0) \end{pmatrix}$$

where $|C_1|^2 + |C_2|^2 = 1$ for all value of time, and ρ_1 and ρ_2 are propagators associated with Pulse 1 and Pulse 2 (both with width τ), respectively. D is a propagator associated with the field-free evolution of duration T .

Specifically, in the interaction representation:

$$\rho_1 = e^{-i\bar{\Delta}\tau} \begin{pmatrix} e^{-i\frac{\Delta_0}{2}\tau} \left(\cos \frac{\Omega'\tau}{2} + i \cos \theta \sin \frac{\Omega'\tau}{2} \right) & i e^{-i(\frac{\Delta_0}{2}\tau - \phi_0)} \sin \theta \sin \frac{\Omega'\tau}{2} \\ i e^{i(\frac{\Delta_0}{2}\tau + \phi_0)} \sin \theta \sin \frac{\Omega'\tau}{2} & e^{i\frac{\Delta_0}{2}\tau} \left(\cos \frac{\Omega'\tau}{2} - i \cos \theta \sin \frac{\Omega'\tau}{2} \right) \end{pmatrix}$$

$$\rho_2 = e^{-i\bar{\Delta}\tau} \begin{pmatrix} e^{-i\frac{\Delta_0}{2}\tau} \left(\cos \frac{\Omega'\tau}{2} + i \cos \theta \sin \frac{\Omega'\tau}{2} \right) & i e^{-i(\frac{\Delta_0}{2}\tau + \phi_0)} \sin \theta \sin \frac{\Omega'\tau}{2} e^{-i\Delta_0(\tau+T)} \\ i e^{i(\frac{\Delta_0}{2}\tau + \phi_0)} \sin \theta \sin \frac{\Omega'\tau}{2} e^{i\Delta_0(\tau+T)} & e^{i\frac{\Delta_0}{2}\tau} \left(\cos \frac{\Omega'\tau}{2} - i \cos \theta \sin \frac{\Omega'\tau}{2} \right) \end{pmatrix}$$

and

$$D = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Note that D , in the interaction representation, is the identity matrix. This is different from Ramsey's original approach in which the state vector does evolve and change during the delay time T . The angle θ is defined as:

$$\sin \theta = \frac{\Omega_0^*}{\Omega'}$$

and

$$\cos \theta = \frac{\Delta_0 + \Delta_d}{\Omega'}$$

where Ω_0^* is the complex conjugate of the Rabi rate, and Ω' can be defined as the "effective Rabi rate."

$$\Omega' = \sqrt{|\Omega_0^*| + (\Delta_0 + \Delta_d)^2}$$

2 Detailed Derivation for P_f

In the interaction representation,

$$i \begin{pmatrix} \dot{a}_i(t) \\ \dot{a}_f(t) \end{pmatrix} = \begin{pmatrix} \Delta_i & -\frac{\Omega_0^*}{2} e^{-i\Delta_0 t} \\ -\frac{\Omega_0}{2} e^{i\Delta_0 t} & \Delta_f \end{pmatrix} \begin{pmatrix} a_i(t) \\ a_f(t) \end{pmatrix} \quad (1)$$

where Δ_i is the ac Stark shift in the $|i\rangle$ state and Δ_f is the ac Stark shift in the $|f\rangle$ state. We first solve for a_i :

$$i\ddot{a}_i = \Delta_i \dot{a}_i - \frac{\Omega_0^*}{2} e^{-i\Delta_0 t} \dot{a}_f + \frac{i\Delta_0}{2} \Omega_0^* e^{-i\Delta_0 t} a_f, \quad (2)$$

where

$$a_f = \left(\frac{\Omega_0^*}{2} e^{-i\Delta_0 t} \right)^{-1} (\Delta_i a_i - i\dot{a}_i) = \frac{2}{\Omega_0^*} e^{i\Delta_0 t} (\Delta_i a_i - i\dot{a}_i). \quad (3)$$

From Eq. (1), we also have

$$\begin{aligned} \dot{a}_f &= i^{-1} \left(-\frac{\Omega_0}{2} e^{i\Delta_0 t} a_i + \Delta_f a_f \right) \\ &= i^{-1} \left[-\frac{\Omega_0}{2} e^{i\Delta_0 t} a_i + \Delta_f (\Delta_i a_i - i\dot{a}_i) \left(\frac{2}{\Omega_0^*} e^{i\Delta_0 t} \right) \right] \\ &= \frac{2i}{\Omega_0^*} e^{i\Delta_0 t} \left(\frac{|\Omega_0|^2}{4} a_i - \Delta_f \Delta_i a_i + i\Delta_f \dot{a}_i \right). \end{aligned} \quad (4)$$

Therefore,

$$\begin{aligned} \ddot{a}_i &= -i\Delta_i \dot{a}_i + \frac{i\Omega_0^*}{2} e^{-i\Delta_0 t} \left[\frac{2i}{\Omega_0^*} e^{i\Delta_0 t} \left(\frac{|\Omega_0|^2}{4} a_i - \Delta_f \Delta_i a_i + i\Delta_f \dot{a}_i \right) \right] \\ &\quad + \frac{\Delta_0}{2} \Omega_0^* e^{-i\Delta_0 t} \frac{2}{\Omega_0^*} e^{i\Delta_0 t} (\Delta_i a_i - i\dot{a}_i) \\ &= -i\Delta_i \dot{a}_i - \frac{|\Omega_0|^2}{4} a_i + \Delta_f \Delta_i a_i - i\Delta_f \dot{a}_i + \Delta_0 \Delta_i a_i - i\Delta_0 \dot{a}_i \\ &= -i(\Delta_0 + \Delta_i + \Delta_f) \dot{a}_i - \left[\frac{|\Omega_0|^2}{4} - \Delta_i(\Delta_f + \Delta_0) \right] a_i \\ &= -i(\Delta_0 + \Delta_i + \Delta_f) \dot{a}_i - \frac{1}{4} (|\Omega_0|^2 - 4\Delta_i \Delta_f - 4\Delta_i \Delta_0) a_i. \end{aligned} \quad (5)$$

We obtain the first second-order homogeneous differential equation:

$$\ddot{a}_i + i(\Delta_0 + \Delta_i + \Delta_f) \dot{a}_i + \left[\frac{|\Omega_0|^2}{4} - \Delta_i(\Delta_f + \Delta_0) \right] a_i = 0. \quad (6)$$

Let a guess solution be $a_i(t) = a_0 e^{i\omega t}$. The characteristic equation is:

$$\begin{aligned} -\omega^2 + i(\Delta_0 + \Delta_i + \Delta_f) \omega + \frac{1}{4} (|\Omega_0|^2 - 4\Delta_0 \Delta_i - 4\Delta_i \Delta_f) &= 0 \\ -\omega^2 - (\Delta_0 + \Delta_i + \Delta_f) \omega + \frac{1}{4} (|\Omega_0|^2 - 4\Delta_0 \Delta_i - 4\Delta_i \Delta_f) &= 0. \end{aligned} \quad (7)$$

Solving the quadratic equation (7) and obtain w :

$$\begin{aligned}\omega &= -\frac{\Delta_0 + \Delta_f + \Delta_i}{2} \pm \frac{1}{2}\sqrt{(\Delta_0 + \Delta_f + \Delta_i)^2 + |\Omega_0|^2 - 4\Delta_0\Delta_i - 4\Delta_i\Delta_f} \\ &= -\frac{\Delta_0 + \Delta_f + \Delta_i}{2} \\ &\quad \pm \frac{1}{2}\sqrt{(\Delta_0 + \Delta_f + \Delta_i)^2 + \Delta_0^2 + |\Omega_0|^2 + (\Delta_i - \Delta_f)^2 - 2\Delta_0(\Delta_i - \Delta_f)}\end{aligned}\quad (8)$$

Let $\bar{\Delta} = (\Delta_i + \Delta_f)/2$ and $\Delta_d = \Delta_f - \Delta_i$, this gives

$$\omega = -\frac{\Delta_0}{2} - \bar{\Delta} \pm \frac{1}{2}\sqrt{|\Omega_0|^2 + (\Delta_0 + \Delta_d)^2}\quad (9)$$

Next, let the “effective Rabi rate” be Ω' , defined as

$$\Omega' = \sqrt{|\Omega_0|^2 + (\Delta_0 + \Delta_d)^2}.\quad (10)$$

The general solution to eq. (7) is:

$$\begin{aligned}a_i &= a_+ e^{i\omega_+ t} + a_- e^{i\omega_- t} \\ &= e^{-i\bar{\Delta}t} e^{-i\frac{\Delta_0}{2}t} \left(a_+ e^{i\frac{\Omega'}{2}t} + a_- e^{-i\frac{\Omega'}{2}t} \right)\end{aligned}\quad (11)$$

So,

$$a_i = e^{-i\bar{\Delta}t} e^{-i\frac{\Delta_0}{2}t} \left(A \cos \frac{\Omega't}{2} + B \sin \frac{\Omega't}{2} \right)\quad (12)$$

Next, we solve for a_f . From eq. (3):

$$\begin{aligned}a_f &= \frac{2}{\Omega_0^*} e^{i\Delta_0 t} (\Delta_i a_i - i\dot{a}_i) \\ &= \frac{2}{\Omega_0^*} e^{i\Delta_0 t} \left[\Delta_i e^{-i\bar{\Delta}t} e^{-i\frac{\Delta_0}{2}t} \left(A \cos \frac{\Omega't}{2} + B \sin \frac{\Omega't}{2} \right) - i\dot{a}_i \right] \\ &= \frac{2}{\Omega_0^*} e^{i\frac{\Delta_0}{2}t} \left[\Delta_i e^{-i\bar{\Delta}t} \left(A \cos \frac{\Omega't}{2} + B \sin \frac{\Omega't}{2} \right) - i\dot{a}_i \right]\end{aligned}\quad (13)$$

where

$$\begin{aligned}-i\dot{a}_i &= (-i)^2 \left(\bar{\Delta} + \frac{\Delta_0}{2} \right) e^{-i\bar{\Delta}t} e^{-i\frac{\Delta_0}{2}t} \left[\left(A \cos \frac{\Omega't}{2} + B \sin \frac{\Omega't}{2} \right) \right. \\ &\quad \left. + -i\frac{\Omega'}{2} \left(-A \sin \frac{\Omega't}{2} + B \cos \frac{\Omega't}{2} \right) \right] \\ &= e^{-i\bar{\Delta}t} e^{-i\frac{\Delta_0}{2}t} \left[-\left(\bar{\Delta} + \frac{\Delta_0}{2} \right) \left(A \cos \frac{\Omega't}{2} + B \sin \frac{\Omega't}{2} \right) \right. \\ &\quad \left. + i\frac{\Omega'}{2} \left(A \sin \frac{\Omega't}{2} - B \cos \frac{\Omega't}{2} \right) \right].\end{aligned}\quad (14)$$

Assume that at $t = 0$, $A = a_i(0)$ and

$$B = i \frac{\Omega_0^*}{\Omega'} a_f(0) + i \frac{\Delta_d + \Delta_0}{\Omega'} a_i(0). \quad (15)$$

So, from Eq. (12):

$$a_i(t) = e^{-i(\bar{\Delta} + \frac{\Delta_0}{2})t} \left\{ a_i(0) \left[\cos \frac{\Omega' t}{2} + i \frac{\Delta_0 + \Delta_d}{\Omega'} \sin \frac{\Omega' t}{2} \right] + a_f(0) \frac{i\Omega_0^*}{\Omega'} \sin \frac{\Omega' t}{2} \right\} \quad (16)$$

From Eq. (13) and (14), we obtain an expression for $a_f(t)$:

$$\begin{aligned} a_f(t) &= \frac{2}{\Omega_0^*} e^{-i\bar{\Delta}t} e^{i\frac{\Delta_0}{2}t} \left\{ \Delta_i \left(a_i(0) \cos \frac{\Omega' t}{2} + \left(\frac{i\Omega_0^*}{\Omega'} a_f(0) + i \frac{\Delta_d + \Delta_0}{\Omega'} a_i(0) \right) \sin \frac{\Omega' t}{2} \right) \right. \\ &\quad \left. - \left(\bar{\Delta} + \frac{\Delta_0}{2} \left(a_i \cos \frac{\Omega' t}{2} + \left(\frac{i\Omega_0^*}{\Omega'} a_f(0) + \frac{i}{2} \frac{\Delta_d + \Delta_0}{\Omega'} a_i(0) \right) \sin \frac{\Omega' t}{2} \right) \right) \right. \\ &\quad \left. i \frac{\Omega'}{2} \left(a_i \sin \frac{\Omega' t}{2} - \left(\frac{i\Omega_0^*}{\Omega'} a_f(0) + \frac{i}{2} \frac{\Delta_d + \Delta_0}{\Omega'} a_i(0) \right) \cos \frac{\Omega' t}{2} \right) \right\} \\ &= \frac{2}{\Omega_0^*} e^{-i\bar{\Delta}t} e^{i\frac{\Delta_0}{2}t} \left\{ \left(\Delta_i - \bar{\Delta} - \frac{\Delta_0}{2} \right) \left(A \cos \frac{\Omega' t}{2} + B \sin \frac{\Omega' t}{2} \right) \right. \\ &\quad \left. \frac{i\Omega'}{2} \left(A \sin \frac{\Omega' t}{2} - B \cos \frac{\Omega' t}{2} \right) \right\}. \end{aligned} \quad (17)$$

Now, notice that

$$\Delta_i - \bar{\Delta} = \Delta_i - \frac{\Delta_i + \Delta_f}{2} = -\frac{\Delta_d}{2}. \quad (18)$$

So,

$$\begin{aligned} a_f(t) &= e^{-i\bar{\Delta}t} e^{i\frac{\Delta_0}{2}t} \left\{ A \left(-\frac{\Delta_d + \Delta_0}{\Omega_0^*} \cos \frac{\Omega' t}{2} + \frac{i\Omega'}{\Omega_0^*} \sin \frac{\Omega' t}{2} \right) \right. \\ &\quad \left. - B \left(\frac{\Delta_d + \Delta_0}{\Omega_0^*} \sin \frac{\Omega' t}{2} + \frac{i\Omega'}{\Omega_0^*} \cos \frac{\Omega' t}{2} \right) \right\} \\ &= e^{-i\bar{\Delta}t} e^{i\frac{\Delta_0}{2}t} \left\{ a_i(0) \left(-\frac{\Delta_d + \Delta_0}{\Omega_0^*} \cos \frac{\Omega' t}{2} + \frac{i\Omega'}{\Omega_0^*} \sin \frac{\Omega' t}{2} \right) \right. \\ &\quad \left. - \left(i \frac{\Omega_0^*}{\Omega'} a_f(0) + i \frac{\Delta_d + \Delta_0}{\Omega'} a_i(0) \right) \left(\frac{\Delta_d + \Delta_0}{\Omega_0^*} \sin \frac{\Omega' t}{2} + \frac{i\Omega'}{\Omega_0^*} \cos \frac{\Omega' t}{2} \right) \right\} \\ &= e^{-i\bar{\Delta}t} e^{i\frac{\Delta_0}{2}t} \left\{ a_i \sin \frac{\Omega' t}{2} \left(\frac{i\Omega'}{\Omega_0^* \Omega'} - \frac{i(\Delta_d + \Delta_0)^2}{\Omega_0^* \Omega'} \right) \right. \\ &\quad \left. a_f(0) \left(\cos \frac{\Omega' t}{2} - i \frac{\Delta_d + \Delta_0}{\Omega'} \sin \frac{\Omega' t}{2} \right) \right\}. \end{aligned} \quad (19)$$

Next, note that

$$\Omega'^2 = (\Delta_0 + \Delta_d)^2 + |\Omega_0|^2 = (\Delta_0 + \Delta_d)^2 + \Omega_0 \Omega_0^*. \quad (20)$$

So

$$a_f(t) = e^{-i\bar{\Delta}t} e^{i\frac{\Delta_0}{2}t} \left\{ a_i(0) \frac{i\Omega_0}{\Omega'} \sin \frac{\Omega't}{2} + a_f(0) \left(\cos \frac{\Omega't}{2} - i \frac{\Delta_d + \Delta_0}{\Omega'} \sin \frac{\Omega't}{2} \right) \right\}. \quad (21)$$

Finally, let us put everything together in matrix form:

$$\begin{pmatrix} a_i(t) \\ a_f(t) \end{pmatrix} = \mathcal{M} \begin{pmatrix} a_i(0) \\ a_f(0) \end{pmatrix}, \quad (22)$$

where \mathcal{M} is the matrix

$$e^{-i\bar{\Delta}t} \begin{pmatrix} e^{-i\frac{\Delta_0}{2}t} \left(\cos \frac{\Omega't}{2} + i \frac{\Delta_d + \Delta_0}{\Omega'} \sin \frac{\Omega't}{2} \right) & e^{-i\frac{\Delta_0}{2}t} \frac{i\Omega_0^*}{\Omega'} \sin \frac{\Omega't}{2} \\ e^{i\frac{\Delta_0}{2}t} \frac{i\Omega_0}{\Omega'} \sin \frac{\Omega't}{2} & e^{i\frac{\Delta_0}{2}t} \left(\cos \frac{\Omega't}{2} - i \frac{\Delta_d + \Delta_0}{\Omega'} \sin \frac{\Omega't}{2} \right) \end{pmatrix} \quad (23)$$