

Name: **Huan Q. Bui**  
 Course: **8.421 - AMO I**  
 Problem set: **#8**  
 Due: Friday, April 8, 2022.

**1. Optical Traps and Scattering.** What are the proper power and wavelength needed to trap an ultracold atomic gas? Consider an alkali atom with resonance frequency  $\omega_0$  on the principal  $nS \rightarrow nP$  transition. A sample of atoms in the ground state  $nS$  are exposed to monochromatic radiation of intensity  $I$  and frequency  $\omega_L < \omega_0$ . Using the fact that essentially all of the oscillator strength out of the ground state comes from the  $nS \rightarrow nP$  transition, we have

$$\alpha(\omega_L) \approx \frac{2e^2}{\hbar} |\langle nP | z | nS \rangle|^2 \frac{\omega_0}{\omega_0^2 - \omega_L^2} \implies |\langle nP | z | nS \rangle|^2 = \frac{\hbar \alpha(\omega_L)}{2e^2} \frac{\omega_0^2 - \omega_L^2}{\omega_0}.$$

(a) AC Stark shift:

(i) From lecture, the AC Stark shift  $U_i$  from time-dependent perturbation theory is given by

$$U_i = -\frac{1}{4} \alpha(\omega_L) \mathcal{E}^2 = -\frac{2I\alpha(\omega_L)}{4c\epsilon_0} = -\frac{I\alpha(\omega_L)}{2c\epsilon_0}.$$

(ii) Now, we want to use the rotating wave approximation to obtain the AC Stark shift. This can be done by first writing down the true (symmetrized) Hamiltonian:

$$\mathcal{H} = \frac{\hbar}{2} \begin{pmatrix} -\omega_0 & \omega_R e^{i\omega_L t} \\ \omega_R^* e^{-i\omega_L t} & \omega_0 \end{pmatrix}.$$

By going into the rotating frame, plus making the rotating wave approximation, we find that

$$\mathcal{H}_{\text{rot}}^{\text{RWA}} = \frac{\hbar}{2} \begin{pmatrix} -\delta & \omega_R \\ \omega_R & \delta \end{pmatrix}$$

where  $\delta = \omega_0 - \omega_L$ . The energy shifts can be obtained from the eigenvalues:

$$\Delta E = \pm \frac{\hbar}{2} \sqrt{\omega_R^2 + \delta^2} = \pm \frac{\hbar}{2} \sqrt{\omega_R^2 + (\omega_0 - \omega_L)^2}.$$

In the limit where the Rabi frequency is much less than the detuning, we simply have that

$$|U_{ii}| = \frac{\hbar}{2} |\omega_0 - \omega_L|.$$

In particular, the energy of the lower state gets shifted down while the energy of the higher state gets shifted up (since we're red-detuning).

(iii) From the previous two parts, we find that

$$\frac{U_i}{U_{ii}} = \frac{I\alpha(\omega_L)}{2c\epsilon_0} \frac{2}{\hbar(\omega_0 - \omega_L)} = \frac{I}{c\epsilon_0} \frac{2e^2}{\hbar^2} |\langle nP | z | nS \rangle|^2 \frac{\omega_0}{(\omega_0 - \omega_L)^2(\omega_0 + \omega_L)}.$$

When  $\omega_L \approx 0$ , we have

$$\frac{U_i}{U_{ii}} \approx \frac{I}{c\epsilon_0} \frac{2e^2}{\hbar^2} |\langle nP | z | nS \rangle|^2 \frac{1}{\omega_0^2} \left(1 + \frac{\omega_L}{\omega_0} + \dots\right).$$

When  $\omega_L \approx \omega_0$ , we may write  $\omega_L + \omega_0 = 2\omega_0$ , so that

$$\frac{U_i}{U_{ii}} \approx \frac{I}{c\epsilon_0} \frac{2e^2}{\hbar^2} |\langle nP | z | nS \rangle|^2 \frac{1}{2(\omega_0 - \omega_L)^2}$$

We see that if the intensity has spatial structure, with the appropriate detuning, the AC Stark shift can have energy minima where the atoms can be trapped.

(b) Using time-dependent perturbation theory, we have that

*blah*

In the RWA picture, we know that

$$P_{e,ii}(t) = \frac{\omega_R^2}{\omega_R^2 + \delta^2} \sin^2 \left( \frac{\sqrt{\omega_R^2 + (\omega_0 - \omega_L)^2} t}{2} \right).$$

Under time-averaging this is

$$P_{e,ii} = \frac{\omega_R^2}{2(\omega_R^2 + \delta^2)}$$

(c) (i)

(ii)

(iii)

(d) (i)

(ii)

## 2. Magic Wavelength Optical Trap.

(a) (i)

(ii)

(iii)

(b) (i)

(ii)

## 3. Species-Dependent and Spin-Dependent AC Stark shift

(a) (i)

(ii)

(b) (i)

(ii)