MATRIX ANALYSIS

A Quick Guide

Huan Bui

Colby College Physics & Statistics Class of 2021

February 11, 2019

Preface

Greetings,

Matrix Analysis: A Quick Guide to is compiled based on my MA353: Matrix Analysis notes with professor Leo Livshits. The sections are based on a number of resources: Linear Algebra Done Right by Axler, A Second Course in Linear Algebra by Horn and Garcia, Matrices and Linear Transformations by Cullen, Matrices: Methods and Applications by Barnett, Problems and Theorems in Linear Algebra by Prasolov, Matrix Operations by Richard Bronson, and professor Leo Livshits' own textbook (in the making).

The development of this text will come in layers. The first layer, one that I am working on during the course of S'19 MA353, will be an overview of the key topics listed in the table of contents. As the semester progresses, I will be constantly updating the existing notes, as well as adding prof. Livshits' problems and my solutions to the problems. The second layer will come after the course is over, when concepts will have hopefully "come together."

I will decide how much narrative I should put into the text as text is developed over the semester. I'm thinking that I will only add detailed explanations wherever I find fit or necessary for my own studies. I will most likely keep the text as condensed as I can.

Technical note: for the conjugate transpose operation, I will use " \dagger " and " \ast " interchangeably. I'm more biased towards \dagger as a physics student, but to appeal to my mathematics professor, I should also learn the mathematicians' notations. Enjoy!

Contents

	Preface	1			
1	List of Special Matrices & Their Properties				
2	List of Operations				
3 List of Algorithms					
4	Complex Numbers 4.1 A different point of view	8			
5	Vector Spaces & Linear Functions 5.1 Review of Linear Spaces and Subspaces 5.2 Review of Linear Maps 5.3 Review of Kernels and Images 5.4 Atrices 5.5 Rank-Nullity Theorem 5.6 Isomorphisms 5.7 Coordinatization & matricial representation of linear functions	12 12 13 13 13 13			
6	Products of vector spaces & Sums of subspaces 6.1 Direct Sums	14 14 14 14 15 15			
7	Idempotents & Resolutions of Identity	16			
8	Block-representations of operators 8.1 Direct sums of operators	17 17			
9	Invariant subspaces 9.1 Reducing subspaces	18 18			
10	Polynomials applied to operators 10.1 Minimal polynomials of block- Δ^r operators 10.2 Minimal polynomials at a vector	19 19			
	Eigentheory 11.1 Spectral Mapping Theorem	20			

12	Triangularization	21			
	12.1 Compression to invariant subspaces	21			
	12.2 Simultaneously Δ -ity of commuting families	21			
13	Diagonalization	22			
	13.1 Spectral resolutions	22			
	13.2 Compressions to reducing subspaces	22			
	13.3 Simultaneous diagonalizability for commuting families	22			
14	14 Primary decomposition over $\mathbb C$ and generalized eigenspaces				
15	Cyclic decomposition and Jordan form	24			
	15.1 Square roots of operators	24			
	15.2 Similarity of a matrix and its transpose	24			
	15.3 Similarity of a matrix and its conjugate	24			
	15.4 Jordan forms of \mathcal{AB} and \mathcal{BA}	24			
	15.5 Power-convergent operators	24			
	15.6 Power-bounded operators	24			
	15.7 Row-stochastic matrices	24			
16	Determinant & Trace	25			
	16.1 Classical adjoints	25			
	16.2 Cayley-Hamilton theorem	25			
17	Inner products and norms	26			
	17.1 Riesz representation theorem	27			
	17.2 Adjoints	27			
	17.3 Grammians	27			
	17.4 Orthogonal complements and orthogonal decompositions	27			
	17.5 Ortho-projections	27			
	17.6 Closest point solutions	27			
	17.7 Gram-Schmidt and orthonormal bases	27			
10	Isomethics and unitary an austans	28			
10	Isometries and unitary operators	40			
19	Ortho-triangularization	29			
20	Consideration and the constant of the constant	90			
20	Spectral resolutions	30			
21 Ortho-diagonalization; self-adjoint and normal operators; spec-					
	tral theorems	31			
99	22 Positive (gami) definite energy				
44	Positive (semi-)definite operators 22.1 Classification of inner products	32 32			
	22.1 Classification of finner products	$\frac{32}{32}$			
	22.2 I OSITIVE SQUARE TOOKS	JΔ			
23	Polar decomposition	33			

24	Sing	gle value decomposition	3 4
	24.1	Spectral/operator norm	34
	24.2	Singular values and approximation	34
	24.3	Singular values and eigenvalues	34

1 List of Special Matrices & Their Properties

1. **Hermitian/Self-adjoint**: $H = H^{\dagger}$. A Hermitian matrix is matrix that is equal to its own conjugate transpose:

$$H$$
 is Hermitian $\iff H_{ij} = \bar{H}_{ji}$

Properties 1.1.

- (a) H is Hermitian $\iff \langle w, Hv \rangle = \langle Hw, v \rangle$, where \langle , \rangle denotes the inner product.
- (b) H is Hermitian $\iff \langle v, Hv \rangle \in \mathbb{R}$.
- (c) H is Hermitian ⇐⇒ it is unitarily diagonalizable with real eigenvalues.

Unitary: $U^*U = UU^* = I = U^{\dagger}U = UU^{\dagger}$. The real analogue of a unitary matrix is an orthogonal matrix. The following list contain the properties of U:

(a) U preserves the inner product:

$$\langle Ux, Uy \rangle = \langle x, y \rangle.$$

- (b) U is normal: it commutes with $U^* = U^{\dagger}$.
- (c) U is diagonalizable:

$$U = VDV^*$$
,

where D is diagonal and unitary, and V is unitary.

- (d) $|\det(U)| = 1$ (hence the real analogue to U is an orthogonal matrix)
- (e) Its eigenspaces are orthogonal.
- (f) U can be written as

$$U = e^{iH}$$
.

where H is a Hermitian matrix.

- (g) Any square matrix with unit Euclidean norm is the average of two unitary matrices.
- 2. Cofactor: ?

2 List of Operations

- 1. Conjugate transpose is what its name suggests.
- 2. Classical adjoint/Adjugate/adjunct of a square matrix is the transpose of its cofactor matrix.

3 List of Algorithms

4 Complex Numbers

4.1 A different point of view

We often think of complex numbers as

$$a+ib$$

where $a, b \in \mathbb{R}$ and $i = \sqrt{-1}$. While there is nothing "bad" about this way of thinking - in fact thinking of complex numbers as a+ib allows us to very quickly and intuitively do arithmetics operations on them - a "matrix representation" of complex numbers can give us some insights on "what we actually do" when we perform complex arithmetics.

Let us think of

$$\begin{pmatrix} a & -b \\ b & a \end{pmatrix}$$

as a different representation of the same object - the same complex number "a+ib." Note that it does not make sense to say the matrix representation **equals** the complex number itself. But we shall see that a lot of the properties of complex numbers are carried into this matrix representation under interesting matricial properties.

First, let us break the the matrix down:

$$a+ib=a\times 1+i\times b\sim \begin{pmatrix} a & -b\\ b & a \end{pmatrix}=a\begin{pmatrix} 1 & 0\\ 0 & 1 \end{pmatrix}+b\begin{pmatrix} 0 & -1\\ 1 & 0 \end{pmatrix}=aI+b\mathcal{I}.$$

Right away, we can make some "mental connections" between the representations:

$$I \sim 1$$

$$\mathcal{I} \sim i.$$

Now, we know that complex number multiplications commute:

$$(a+ib)(c+id) = (c+id)(a+ib).$$

Matrix multiplications are not commutative. So, we might wonder whether commutativity holds under the this new representation of complex numbers. Well, the answer is yes. We can readily verify that

$$(aI + b\mathcal{I})(cI + b\mathcal{I}) = (cI + b\mathcal{I})(aI + b\mathcal{I}).$$

How about additions? Let's check:

$$(a+ib)+(c+id)=(a+c)+i(b+d) \sim \begin{pmatrix} a+c & -(b+d) \\ (b+d) & a+c \end{pmatrix} = \begin{pmatrix} a & -b \\ b & a \end{pmatrix} + \begin{pmatrix} c & -d \\ d & c \end{pmatrix}.$$

Ah! Additions work. So, the new representation of complex numbers seems to be working flawlessly. However, we have yet to gain any interesting insights into the connections between the representations. To do that, we have to look into changing the form of the matrix. First, let's see what conjugation does:

$$(a+ib)^* = a-ib \sim \begin{pmatrix} a & b \\ -b & a \end{pmatrix} = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}^{\top}$$

Ah, so conjugation to a complex number in the traditional representation is the same as transposition in the matrix representations. What about the amplitude square? Let us call

$$M = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}.$$

We have

$$(a+ib)(a-ib) \sim \begin{pmatrix} a & -b \\ b & a \end{pmatrix} \begin{pmatrix} a & b \\ -b & a \end{pmatrix} = MM^{\top} = (a^2+b^2)I = \det(M)I$$

Interesting. But observe that if $det(M) \neq 0$

$$\frac{1}{\det(M)}MM^{\top} = I.$$

This tells us that

$$M^{\top} = M^{-1},$$

where M^{-1} is the inverse of M, and, not surprisingly, it corresponds to the reciprocal to the complex number a+ib. We can readily show that

$$M^{-1} \sim (a+ib)^{-1} = \frac{1}{a^2 + b^2}(a-ib).$$

Remember that we can also think of a complex number as a column vector:

$$c + id \sim \begin{pmatrix} c \\ d \end{pmatrix}$$
.

Let us look back at complex number multiplication under matrix representation:

$$(a+ib)(c+id) = (ac-bd) + i(bc+ad) \sim \begin{pmatrix} a & -b \\ b & a \end{pmatrix} \begin{pmatrix} c \\ d \end{pmatrix} = \begin{pmatrix} ac-bd \\ bc+ad \end{pmatrix}.$$

Multiplication actually works in this "mixed" way of representing complex numbers as well. Now, observe that what we just did was performing a linear transformation on a vector in \mathbb{R}^2 . It is always interesting to look at the geometrical interpretation of this transformation. To do this, let us call N the "normalized" version of M:

$$N = \frac{1}{\sqrt{a^2 + b^2}} \begin{pmatrix} a & -b \\ b & a \end{pmatrix}.$$

We immediately recognize that N is an orthogonal matrix. This means N is an orthogonal transformation (length preserving). Now, it is reasonable to define

$$\cos \theta = \frac{a}{\sqrt{a^2 + b^2}}$$
$$\sin \theta = \frac{b}{\sqrt{a^2 + b^2}}.$$

We can write N as

$$N = (\cos \theta - \sin \theta \sin \theta \cos \theta)$$
,

which is a physicists' favorite matrix: the rotation by θ . So, let us write M in terms of N:

$$M = \begin{pmatrix} a & -b \\ b & a \end{pmatrix} = \sqrt{a^2 + b^2} N = \sqrt{a^2 + b^2} \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

We can interpret M as a rotation by θ , followed by a scaling by $\sqrt{(a^2+b^2)}$. But what $\sqrt{a^2+b^2}$ exactly is just the "length" or the "amplitude" of the complex number a+ib, if we think of it as an arrow in a plane.

4.2 Relevant properties and definitions

- 1. The modulus of z=a+ib is the "amplitude" of z, denoted by $|z|=\sqrt{a^2+b^2}=z\bar{z}.$
- 2. The modulus is *multiplicative*, i.e.

$$|wz| = |w||z|.$$

3. Triangle inequality:

$$|z+w| < |z| + |w|.$$

We can readily show this geometrically, or algebraically.

4. The argument of z = a + ib is θ , where

$$\theta = \begin{cases} \tan^{-1}\left(\frac{b}{a}\right), \text{ if } a > 0\\ \frac{\pi}{2} + k2\pi, k \in \mathbb{R} \text{ if } a = 0, b > 0\\ -\frac{\pi}{2} + k2\pi, k \in \mathbb{R} \text{ if } a = 0, b < 0\\ \text{Undefined if } a = b = 0. \end{cases}$$

5. The *conjugate* of a+ib is a-ib. Conjugation is *additive* and *multiplicative*, i.e.

$$z + \bar{w} = \bar{z} + \bar{w}$$
$$\bar{w}z = \bar{w}\bar{z}.$$

Note that we can also show the multiplicative property with the matrix representation as well:

$$\bar{wz} \sim (WZ)^{\top} = Z^{\top}W^{\top} \sim \bar{z}\bar{w} = \bar{w}\bar{z}.$$

6. Euler's identity, generalized to de Moivre's formula:

$$z^n = r^n e^{in\theta}.$$

5 Vector Spaces & Linear Functions

5.1 Review of Linear Spaces and Subspaces

Properties 5.1. of linear spaces:

- 1. Commutativity and associativity of addition
- 2. Existence of an additively neutral element (null element). Zero multiples of elements give the null element: $0 \cdot V = \mathbf{0}$
- 3. Every element has an (unique) additively antipodal element
- 4. Scalar multiplication distributes over addition
- 5. Multiplicative identity:
- 6. $ab \cdot V = a \cdot (bV)$
- 7. (a + b)V = aV + bV

W is a subspace of V if

- 1. $S \subseteq V$
- 2. S is non-empty
- 3. S is closed under addition and scalar multiplication

Properties 5.2. that are interesting/important/maybe-not-so-obvious:

- 1. If S is a subspace of V and $S \neq V$ then S is a proper subspace of V.
- 2. If X in a subspace of Y and Y is a subspace of Z, then X is a subspace of W.
- 3. Non-trivial linear (non-singleton) spaces are infinite.

5.2 Review of Linear Maps

Consider linear spaces V and W and elements $v \in V$ and $w \in W$ and scalars $\alpha, \beta \in \mathbb{R}$, a function $F: V \to W$ is a linear map if

$$F[\alpha v + \beta w] = \alpha F[v] + \beta F[w].$$

Properties 5.3.

- 1. $F[\mathbf{0}_V] = \mathbf{0}_W$
- 2. $G[w] = (\alpha \cdot F)[w] = \alpha \cdot F[w]$
- 3. Given $F: V \to W$ and $G: V \to W$, H[v] = F[v] + G[v] = (F+G)[v] is call the sum of the functions F and G.

- 4. Linear combinations of linear maps are linear.
- 5. Compositions of linear maps are linear.
- 6. Compositions distributes over linear combinations of linear maps.
- 7. Inverses of linear functions (if they exist) are linear.
- 8. Inverse of a bijective linear function is a bijective linear function

5.3 Review of Kernels and Images

Definition 5.1. Let $F: V \to W$ be given. The kernel of F is defined as

$$\ker(F) = \{ v \in V | F[v] = \mathbf{0}_W \}.$$

Properties 5.4. Let $F: V \to W$ a linear map be given. Also, consider a linear map G such that $F \circ G$ is defined

- 1. F is null $\iff \ker(F) = V \iff \operatorname{Im}(F) = \mathbf{0}_W$
- 2. ker(F) is a subspace of V
- 3. Im(F) is a subspace of W
- 4. F is injective $\iff \ker(F) = \mathbf{0}_V$
- 5. $\ker(F) \subseteq \ker(F \circ G)$.
- 6. F injective $\implies \ker(F) = \ker(F \circ G)$

Definition 5.2. Let $F: V \to W$ be given. The image of F is defined as

$$\operatorname{Im}(F) = \{ w \in W | \exists v \in V, F[v] = w \}.$$

- 5.4 Atrices
- 5.5 Rank-Nullity Theorem
- 5.6 Isomorphisms
- 5.7 Coordinatization & matricial representation of linear functions

6 Products of vector spaces & Sums of subspaces

6.1 Direct Sums

Definition 6.1. Let U_j , $j=1,2,\ldots m$ are subspaces of V. $\sum_1^m U_j$ is a direct sum if each $u \in \sum U_j$ can be written in only one way as $u = \sum_1^m u_j$. The direct sum $\sum_i^m U_i$ is denoted as $U_1 \oplus \cdots \oplus U_m$.

Properties 6.1.

- 1. Condition for direct sum: If all U_j are subspaces of V, then $\sum_{1}^{m} U_j$ is a direct sum \iff the only way to write 0 as $\sum_{1}^{m} u_j$, where $u_j \in U_j$ is to take $u_j = 0$ for all j.
- 2. If U, W are subspaces of V and $U \cap W = \{0\}$ then U + W is a direct sum.

6.2 Products and Quotients of Vector Spaces

6.3 Products of Vector Spaces

Definition 6.2. Product of vectors spaces

$$V_1 \times \cdots \times V_m = \{(v_1, \dots, v_m) : v_j \in V_j, j = 1, 2, \dots, m\}.$$

Definition 6.3. Addition on $V_1 \times \cdots \times V_m$:

$$(u_1, \ldots, u_m) + (v_1, \ldots, v_m) = (u_1 + v_1, \ldots, v_m + u_m).$$

Definition 6.4. Scalar multiplication on $V_1 \times \cdots \times V_m$:

$$\lambda(v_1,\ldots,v_m)=(\lambda v_1,\ldots,\lambda v_m).$$

Properties 6.2.

- 1. Product of vectors spaces is a vector space.
 - V_j are vectors spaces over $\mathbb{F} \implies V_1 \times \cdots \times V_m$ is a vector space over \mathbb{F} .
- 2. Dimension of a product is the sum of dimensions:

$$\dim(V_1 \times \cdots \times V_m) = \sum_{1}^{m} \dim(V_j)$$

6.4 Products & Direct Sums

Properties 6.3.

1. Let U_1, \ldots, U_m be subspaces of V. Define a linear map $\Gamma: U_1 \times \cdots \times U_m \to U_1 + \cdots + U_m$ by:

$$\Gamma(u_1,\ldots,u_m)=\sum_1^m u_j.$$

 $U_1 + \cdots + U_m$ is a direct sum $\iff \Gamma$ is injective.

2. Let U_j be finite-dimensional and are subspaces of V.

$$U_1 \oplus \cdots \oplus U_m \iff \dim(U_1 + \cdots + U_m) = \sum_{1}^{m} \dim(U_j)$$

- 6.5 Quotients of Vector Spaces
- 6.6 Nullspaces
- 6.7 Ranges of Operator Powers

7 Idempotents & Resolutions of Identity

- 8 Block-representations of operators
- 8.1 Direct sums of operators

- 9 Invariant subspaces
- 9.1 Reducing subspaces

- 10 Polynomials applied to operators
- 10.1 Minimal polynomials of block- Δ^r operators
- 10.2 Minimal polynomials at a vector

- 11 Eigentheory
- 11.1 Spectral Mapping Theorem

- 12 Triangularization
- 12.1 Compression to invariant subspaces
- 12.2 Simultaneously Δ -ity of commuting families

- 13 Diagonalization
- 13.1 Spectral resolutions
- 13.2 Compressions to reducing subspaces
- 13.3 Simultaneous diagonalizability for commuting families

14 Primary decomposition over $\mathbb C$ and generalized eigenspaces

- 15 Cyclic decomposition and Jordan form
- 15.1 Square roots of operators
- 15.2 Similarity of a matrix and its transpose
- 15.3 Similarity of a matrix and its conjugate
- 15.4 Jordan forms of AB and BA
- 15.5 Power-convergent operators
- 15.6 Power-bounded operators
- 15.7 Row-stochastic matrices

- 16 Determinant & Trace
- 16.1 Classical adjoints
- 16.2 Cayley-Hamilton theorem

17 Inner products and norms

- 17.1 Riesz representation theorem
- 17.2 Adjoints
- 17.3 Grammians
- 17.4 Orthogonal complements and orthogonal decompositions
- 17.5 Ortho-projections
- 17.6 Closest point solutions
- 17.7 Gram-Schmidt and orthonormal bases

18 Isometries and unitary operators

19 Ortho-triangularization

20 Spectral resolutions

21 Ortho-diagonalization; self-adjoint and normal operators; spectral theorems

- 22 Positive (semi-)definite operators
- 22.1 Classification of inner products
- 22.2 Positive square roots

23 Polar decomposition

- 24 Single value decomposition
- 24.1 Spectral/operator norm
- 24.2 Singular values and approximation
- 24.3 Singular values and eigenvalues