

BCS order parameter, critical  
temperature and bandgap

# Selfconsistency equation for $\Delta$

$$b_k = \langle \hat{c}_{-k\downarrow} \hat{c}_{k\uparrow} \rangle = u_k v_k \langle 1 - \hat{a}_{k0}^+ \hat{a}_{k0} - \hat{a}_{k1}^+ \hat{a}_{k1} \rangle$$

Quasiparticles are fermions:  $\langle \hat{a}_{ki}^+ \hat{a}_{ki} \rangle = f(E_k)$  Fermi-Dirac distribution

Order parameter  $\Delta_k = \sum_{\vec{k}'} V_{k,k'} b_k$

# Selfconsistency equation for $\Delta$

$$b_k = \langle \hat{c}_{-k\downarrow} \hat{c}_{k\uparrow} \rangle = u_k v_k \langle 1 - \hat{a}_{k0}^+ \hat{a}_{k0} - \hat{a}_{k1}^+ \hat{a}_{k1} \rangle$$

Quasiparticles  
are fermions:

$$\langle \hat{a}_{ki}^+ \hat{a}_{ki} \rangle = f(E_k)$$

Fermi-Dirac  
distribution

Order  
parameter

$$\Delta_k = \sum_{\vec{k}'} V_{k,k'} b_{k'}$$

$$\Delta_k = - \sum_{k'} V_{k,k'} \frac{\Delta_{k'}}{2 E_{k'}} [1 - 2f(E_{k'})]$$

# Selfconsistency equation for $\Delta$

$$b_k = \langle \hat{c}_{-k\downarrow} \hat{c}_{k\uparrow} \rangle = u_k v_k \langle 1 - \hat{a}_{k0}^+ \hat{a}_{k0} - \hat{a}_{k1}^+ \hat{a}_{k1} \rangle$$

Quasiparticles are fermions:  $\langle \hat{a}_{ki}^+ \hat{a}_{ki} \rangle = f(E_k)$  Fermi-Dirac distribution

Order parameter  $\Delta_k = \sum_{\vec{k}'} V_{k,k'} b_k$

$$\Delta_k = - \sum_{k'} V_{k,k'} \frac{\Delta_{k'}}{2 E_{k'}} [1 - 2f(E_{k'})]$$

Ground state satisfies:  $a_{k,0} |g_s\rangle = 0, \quad a_{k,1} |g_s\rangle = 0$

This gives:  $|g_s\rangle = \prod_k \left( u_k + v_k c_{k\uparrow}^+ c_{-k\downarrow}^+ \right) |0\rangle$

# BCS critical temperature

$$\Delta_k = - \sum_{k'} V_{k,k'} \frac{\Delta_{k'}}{2 E_{k'}} \tanh \left( \frac{E_{k'}}{2 k_B T} \right)$$

Apprx: k-independent interaction

(s-wave scattering)  $V_{k,k'} = -U \quad (-\omega_D < \epsilon_k, \epsilon_{k'} < \omega_D)$

Gap eqn  
for  $\Delta(T)$

# BCS critical temperature

$$\Delta_k = - \sum_{k'} V_{k,k'} \frac{\Delta_{k'}}{2 E_{k'}} \tanh \left( \frac{E_{k'}}{2 k_B T} \right)$$

Apprx: k-independent interaction

(s-wave scattering)  $V_{k,k'} = -U \quad (-\omega_D < \epsilon_k, \epsilon_{k'} < \omega_D)$

Gap eqn  
for  $\Delta(T)$

$$\frac{1}{U} = \sum_k \frac{\tanh(E_k / 2 k_B T)}{2 E_k}$$

$$\begin{aligned} \sum_k \dots &= \int_{-\omega_D}^{\omega_D} \dots v(\epsilon) d\epsilon \\ &\approx v(\epsilon_F) \int_{-\omega_D}^{\omega_D} \dots d\epsilon \end{aligned}$$

# BCS critical temperature

$$\Delta_k = - \sum_{k'} V_{k,k'} \frac{\Delta_{k'}}{2 E_{k'}} \tanh \left( \frac{E_{k'}}{2 k_B T} \right)$$

Apprx: k-independent interaction

(s-wave scattering)  $V_{k,k'} = -U$  ( $-\omega_D < \epsilon_k, \epsilon_{k'} < \omega_D$ )

Gap eqn  
for  $\Delta(T)$

$$\frac{1}{U} = \sum_k \frac{\tanh(E_k / 2 k_B T)}{2 E_k}$$

$$\begin{aligned} \sum_k \dots &= \int_{-\omega_D}^{\omega_D} \dots v(\epsilon) d\epsilon \\ &\approx v(\epsilon_F) \int_{-\omega_D}^{\omega_D} \dots d\epsilon \end{aligned}$$

Critical temperature: set  $\Delta=0$

$$\frac{1}{U v(\epsilon_F)} = \int_{-\omega_D}^{\omega_D} \frac{d\epsilon}{2\epsilon} \tanh \left( \frac{\epsilon}{2 k_B T_c} \right) = \int_0^{\omega_D / 2 k_B T_c} \frac{\tanh x}{x} dx \approx \ln \left( \frac{2\gamma}{\pi} \frac{\omega_D}{2 k_B T_c} \right)$$

$\gamma = 1.78 \dots$

Euler constant

# BCS critical temperature

$$\Delta_k = - \sum_{k'} V_{k,k'} \frac{\Delta_{k'}}{2 E_{k'}} \tanh \left( \frac{E_{k'}}{2 k_B T} \right)$$

Apprx: k-independent interaction

(s-wave scattering)  $V_{k,k'} = -U$  ( $-\omega_D < \epsilon_k, \epsilon_{k'} < \omega_D$ )

Gap eqn  
for  $\Delta(T)$

$$\frac{1}{U} = \sum_k \frac{\tanh(E_k / 2 k_B T)}{2 E_k}$$

$$\sum_k \dots = \int_{-\omega_D}^{\omega_D} \dots v(\epsilon) d\epsilon$$

$$\approx v(\epsilon_F) \int_{-\omega_D}^{\omega_D} \dots d\epsilon$$

Critical temperature: set  $\Delta=0$

$$\frac{1}{U v(\epsilon_F)} = \int_{-\omega_D}^{\omega_D} \frac{d\epsilon}{2\epsilon} \tanh \left( \frac{\epsilon}{2 k_B T_c} \right) = \int_0^{\omega_D / 2 k_B T} \frac{\tanh x}{x} dx \approx \ln \left( \frac{2\gamma}{\pi} \frac{\omega_D}{2 k_B T_c} \right)$$

$k_B T_c \ll \hbar \omega_D$

$\gamma = 1.78 \dots$

$$k_B T_c = 1.13 \hbar \omega_D \exp \left( - \frac{1}{U v(\epsilon_F)} \right)$$

Euler constant



# Temperature dependence, exper tests

Determine the value  $\Delta$  at  $T=0$

$$\frac{1}{U \nu(\epsilon_F)} = \int_{-\omega_D}^{\omega_D} \frac{d\epsilon}{2\sqrt{\epsilon^2 + \Delta^2}} \approx \ln\left(\frac{2\omega_D}{\Delta}\right)$$

gives

$$\Delta = 2 \hbar \omega_D \exp\left(-\frac{1}{U \nu(\epsilon_F)}\right)$$

(agrees w our  
prev result)

# Temperature dependence, exper tests

Determine the value  $\Delta$  at  $T=0$

$$\frac{1}{U \nu(\epsilon_F)} = \int_{-\omega_D}^{\omega_D} \frac{d\epsilon}{2\sqrt{\epsilon^2 + \Delta^2}} \approx \ln\left(\frac{2\omega_D}{\Delta}\right)$$

gives

$$\Delta = 2 \hbar \omega_D \exp\left(-\frac{1}{U \nu(\epsilon_F)}\right)$$

(agrees w our  
prev result)

A universal relation (testable):

$$2 \Delta_{T=0} \approx 3.52 k_B T_c$$

# Temperature dependence, exper tests

Determine the value  $\Delta$  at  $T=0$

$$\frac{1}{U v(\epsilon_F)} = \int_{-\omega_D}^{\omega_D} \frac{d\epsilon}{2\sqrt{\epsilon^2 + \Delta^2}} \approx \ln\left(\frac{2\omega_D}{\Delta}\right)$$

gives

$$\Delta = 2 \hbar \omega_D \exp\left(-\frac{1}{U v(\epsilon_F)}\right)$$

(agrees w our  
prev result)

A universal relation (testable):

$$2 \Delta_{T=0} \approx 3.52 k_B T_c$$

**Temperature dependence  $\Delta(T)$ :**

\* Second-order phase transition,

$\Delta(T)$  vanishes at  $T=T_c$

\* measured in microwave  
absorption experiments

