

Midterm #2

Name Huan Q. Bui

Course 8.333 Stat Mech 1

Prof. Kardar

Date: Oct 29, 2021

(1)

⑦ Poisson brackets

(a) $I = \int dP A \{B, C\}$

$$= \int dP A \left\{ \frac{\partial B}{\partial \vec{q}} \frac{\partial C}{\partial \vec{p}} - \frac{\partial B}{\partial \vec{p}} \frac{\partial C}{\partial \vec{q}} \right\}$$

Integration by parts \rightarrow $= \int dP \left\{ -\frac{\partial C}{\partial \vec{p}} \cdot \frac{\partial A}{\partial \vec{q}} \cdot B + B \frac{\partial A}{\partial \vec{p}} \frac{\partial C}{\partial \vec{q}} \right\}$

$$= \int dP B \left\{ \frac{\partial C}{\partial \vec{q}} \frac{\partial A}{\partial \vec{p}} - \frac{\partial C}{\partial \vec{p}} \frac{\partial A}{\partial \vec{q}} \right\}$$

$$= \int dP B \{C, A\} \quad \checkmark$$

□

(b) If $C(A) = F(A)$ then

$$\int dP A \{B, C\} = \int dP B \{C, A\} = \int dP B \{F(A), A\} = 0$$

\uparrow
part (a)

Since $\{F(A), A\} = \frac{\partial F(A)}{\partial \vec{q}} \cdot \frac{\partial A}{\partial \vec{p}} - \frac{\partial F(A)}{\partial \vec{p}} \cdot \frac{\partial A}{\partial \vec{q}}$

$$= F' \frac{\partial A}{\partial \vec{q}} \cdot \frac{\partial A}{\partial \vec{p}} - F' \frac{\partial A}{\partial \vec{p}} \cdot \frac{\partial A}{\partial \vec{q}} = 0$$

□

Note that we could also do part (a) by using a Poisson bracket identity -- (Leibniz rule)

$$I = \int d\Gamma A \{B, C\}$$

$$= \int d\Gamma (\{AB, C\} - \{A, C\}B)$$

Now $\int d\Gamma \{AB, C\}$

$$= \int d\Gamma \frac{\partial(AB)}{\partial \vec{q}} \frac{\partial C}{\partial \vec{p}} - \frac{\partial(AB)}{\partial \vec{p}} \frac{\partial C}{\partial \vec{q}}$$

(\therefore by parts) $= - \int d\Gamma AB \frac{\partial^2 C}{\partial \vec{p} \partial \vec{q}} + AB \frac{\partial C}{\partial \vec{p} \partial \vec{q}} = 0$

So

$$I = \int d\Gamma A \{B, C\} = - \int d\Gamma B \{A, C\}$$

$$= \int d\Gamma B \{C, A\}$$

✓

$$(c) \quad \partial_t \rho = \{ \mathcal{H}, \rho \}$$

$$S(t) = - \int d\Gamma \rho(\Gamma, t) \ln \rho(\Gamma, t)$$

Then

$$\frac{dS}{dt} = \frac{\partial S}{\partial t} = - \int d\Gamma (\partial_t \rho) \ln \rho(\Gamma, t) + \frac{\rho(\Gamma, t)}{\rho(\Gamma, t)} (\partial_t \rho(\Gamma, t))$$

$$= - \int d\Gamma (\partial_t \rho) (\ln \rho(\Gamma, t) + 1)$$

$$= - \int d\Gamma \{ \mathcal{H}, \rho \} (\ln \rho(\Gamma, t) + 1)$$

$$= - \int d\Gamma \{ \mathcal{H}, \rho \} \ln \rho(\Gamma, t)$$

$$\int d\Gamma \{ \mathcal{H}, \rho \} = 0$$

by integration
by parts



Assuming

$$u = \vec{p}^2 / 2m + u$$

~~$$= - \int d\Gamma \rho \{ \mathcal{H}, \rho \} \ln \rho$$~~

$$= - \int d\Gamma (\ln \rho) \left\{ \frac{\partial u}{\partial \vec{q}} \cdot \frac{\partial \rho}{\partial \vec{p}} - \frac{\vec{p}}{m} \frac{\partial \rho}{\partial \vec{q}} \right\}$$

$$(\text{integration by parts}) = \int d\Gamma \rho \frac{\partial u}{\partial \vec{q}} \cdot \frac{1}{\rho} \cdot \frac{\partial \rho}{\partial \vec{p}} - \int d\Gamma \rho \frac{\vec{p}}{m} \frac{1}{\rho} \frac{\partial \rho}{\partial \vec{q}}$$

$$(\text{by parts again}) = - \int d\Gamma \rho \frac{\partial^2 u}{\partial \vec{p}^2 \partial \vec{q}} + \int d\Gamma \rho \frac{\partial^2 u}{\partial \vec{q}^2} \frac{\vec{p}}{m}$$

$$= 0 + 0$$

$$= 0$$

$$\Rightarrow \boxed{\frac{dS(t)}{dt} = 0}$$

✓

(3)

(1)

$$\langle A \rangle(t) = \int dP_p (P, t) A(P)$$

$$\frac{d\langle A \rangle}{dt} = \int dP_p \quad \text{Since } \frac{\partial_t A(P)}{A(P)} = 0.$$

$$= \int dP_p A - \frac{\partial_p}{\partial t} (P) \quad \partial_t p = \{H, p\}$$

$$= \sum_{\alpha=1}^{3N} \int dP_p A \times \left(\frac{\partial p}{\partial q_\alpha} \frac{\partial^2 H}{\partial q_\alpha} - \frac{\partial p}{\partial q_\alpha} \frac{\partial^2 H}{\partial p_\alpha} \right)$$

Integration by parts...

$$= - \sum_{\alpha=1}^{3N} \int dP_p \left\{ \overbrace{\left[\frac{\partial A}{\partial p_\alpha} \frac{\partial^2 H}{\partial q_\alpha} - \frac{\partial A}{\partial q_\alpha} \frac{\partial^2 H}{\partial p_\alpha} \right]}^{\{H, A\}} + A \left(\frac{\partial^2 H}{\partial p_\alpha \partial q_\alpha} - \frac{\partial^2 H}{\partial q_\alpha \partial p_\alpha} \right) \right\} \rightarrow 0$$

$$= - \int dP_p \{H, A\} = \langle \{A, H\} \rangle$$

✓

(4)

(2)

(a) Boltzmann Eqns

~~$$L_a f_a = C_{aa} + C_{cb} + C_{ac}$$~~

$$\begin{cases} L_a f_a = C_{aa} + C_{cb} + C_{ac} \\ L_b f_b = C_{bb} + C_{ba} + C_{bc} \\ L_c f_c = C_{cc} + C_{ca} + C_{cb} \end{cases}$$

where $L_a = \partial_t + \frac{\vec{p}_a}{m_a} \cdot \vec{\nabla}$

(b) No collision $\Rightarrow C_{\alpha\beta} = \delta_{\alpha\beta} C_{\alpha\alpha}$

\rightarrow zeroth order solutions are simply the local equilibrium solutions:

$$f_a^0(\vec{p}, \vec{q}, t) = \frac{n_a(\vec{q}, t)}{[2\pi k_B T(\vec{q}, t) m]^{3/2}} \exp \left\{ - \frac{(\vec{p} - m \vec{u}(\vec{q}, t))^2}{2 m k_B T(\vec{q}, t)} \right\}$$

for $\alpha = a, b, c$ \rightarrow More generally, $f_\alpha^{(0)} \propto A_\alpha \exp(-\beta H_\alpha)$

(c) Suppose only (a), (b) interact and (c) does nothing, then f_c stays the same, i.e. $f_c' = 0$ since there's no collision with (c).

\rightarrow get solution with interaction between a & b ...

$$\begin{cases} L_a f_a = C_{aa} + C_{ab} \\ L_b f_b = C_{bb} + C_{ba} \end{cases} \quad L_c f_c = 0$$

In this case, f_a, f_b will have corrections from (5)
higher order terms --- For example, the first order correction is

$$f'_\alpha(\vec{p}, \vec{z}, t) = f_\alpha^0 \left(1 - \tau_\alpha \mathcal{L}_\alpha [\ln f_\alpha^0] \right) \rightarrow \text{first order correction}$$

where $\alpha = a, b$.

we can work this out through tedious calculations...

- ① For (a) and (c), they will have the same form if c they only interact with (b)

$$\mathcal{L}_a f_a = C_{aa} + C_{ab} \rightarrow \text{same form as part (c)}$$

$$\mathcal{L}_c f_c = C_{cc} + C_{bc} \rightarrow \text{same form as part (c)}$$

$$\mathcal{L}_b f_b = C_{bb} + C_{ba} + C_{bc}$$

- ② What are the slow hydrodynamic modes?

In the slow hydrodynamic modes, collision terms can be considered zero, and we get that the densities follow the Vlasov equations...

(6)

In this case, we have that $f_a^{(s)}$ satisfies

$$\left\{ \partial_t + \sum_{n=1}^5 \left(\frac{\vec{p}_n}{m_a} \cdot \frac{\partial}{\partial \vec{q}_n} - \frac{\partial U_{eff,a}}{\partial \vec{q}_n} \cdot \frac{\partial}{\partial \vec{p}_n} \right) \right\} f_a^{(s)} = 0$$

where $U_{eff,a} = U_a(\vec{q}) + N_a \int dV' V(\vec{q} - \vec{q}') \rho_1^{(a)}(\vec{x}', t)$

here $V_a = V_{ab} + V_{ba} + V_{ac}$

$V_b = V_{ba} + V_{bc} + V_{ba} \dots$

$V_c = V_{cb} + V_{ca} + V_{cc}$

(f) Consider N_a, N_b, N_c in a box of volume V .

Then we have

$$H_{eff}^{(a)} = \sum_{i=1}^N \left\{ \frac{\vec{p}_i^2}{2m} + U_{eff}^{(a)}(\vec{q}_i) \right\}$$

where

$$U_{eff}^{(a)} = 0 + N_a \int dV' V(\vec{q} - \vec{q}') \frac{1}{V} g_a^{(a)}(\vec{p}) = \frac{N_a}{V} \int d\vec{q}' V(\vec{q} - \vec{q}') g_a^{(a)}(\vec{p})$$

where $\rho_1^{(a)} = \frac{g_a^{(a)}(\vec{p})}{V}$

Substituting this into the Vlasov eqn gives

$$\left(\partial_t + \frac{\vec{p}}{m} \cdot \frac{\partial}{\partial \vec{q}} \right) \rho_1^{(a)} = 0 \rightarrow \text{stationary solution}$$

And so the equilibrium form for f_a is

$$f_1^{(a)}(\vec{p}, \vec{q}) = \frac{N_a}{V} g^{(a)}(\vec{p}) \quad \text{where we have used } f_1^{(a)} = N_a \rho_1^{(a)} \quad (\text{for any } g^{(a)}(\vec{p})) \rightarrow \text{same momentum dist.}$$

(7)

In general,
~~However~~ We can recover $f^{(d)}$ by independence

$$f_s^{(d)} = \frac{N_A!}{(N_A - s)!} \prod_{n=1}^s f_7^{(d)}(\vec{x}_n, t)$$

So

$$\boxed{f_s^{(d)} = N_A! \prod_{n=1}^s f^{(d)}(\vec{p}, \vec{x}) / V}$$

(setting $s = N$)

no more \leftrightarrow dependence
 since we're at equilibrium.