Biogenic Thermal Buffering Along a Forest Maturity Gradient at Huaravito Reserve

Preliminary Field Observation Report

Randall Santamaría Artavia
Custodian-Director
Huaravito Long-term Ecological Monitoring Site
Piedades Sur, San Ramón, Costa Rica
10.104901, -84.565024 | 1,250 m.a.s.l.
Field Observation Date: February 20, 2026

Executive Summary

A preliminary thermal transect conducted on February 20, 2026 documented a 10.2°C temperature gradient between degraded adjacent lands and the mature forest core at Huaravito Reserve. While the exposed clay extraction site (Tajo) exhibited 27.5°C under midday conditions, the mature forest interior ("The Shrine") maintained 17.3°C. This substantial thermal buffering effect, if confirmed through temporal replication, would position Huaravito at the upper range of biogenic cooling effects documented in tropical forest microclimate literature (typically 3–8°C). This preliminary observation warrants systematic investigation to determine whether the extreme gradient is reproducible and to identify the structural and physiological mechanisms responsible.

1. Objective

Document the thermal gradient along a degraded-land-to-mature-forest transect to quantify the potential biogenic cooling effect of the Huaravito forest fragment relative to adjacent anthropogenically disturbed areas.

2. Methods

2.1 Site Description

The transect extends approximately 150 meters horizontally from an abandoned clay extraction site (Tajo) through regenerating secondary forest into the mature primary forest core of Huaravito Reserve. The mature core is dominated by old-growth Hauya elegans (Onagraceae), locally known as "Borrego," with individuals exhibiting extensive buttress roots and heavy epiphytic colonization (>70% estimated trunk surface coverage by bryophytes, orchids, and ferns). The forest floor is characterized by a dense organic layer approximately 20 cm deep.

2.2 Sampling Protocol

Seven measurement points were established along the transect, representing a gradient of increasing vegetation maturity and structural complexity. Measurements were taken during midday hours under partly cloudy conditions with light north wind.

Methodological Limitations (to be addressed in future sampling):

- Instrument type and calibration status not recorded
- Measurement height not standardized (assumed ~1.5 m but not verified)
- Single reading per point (no spatial or temporal replicates)
- Exact time of each measurement not recorded (solar angle variation possible)
- Cloud cover percentage not quantified
- Wind speed not measured
- Relative humidity not recorded
- Canopy cover not quantified at each point

These limitations mean that the observed gradient, while striking, must be interpreted as preliminary and requiring rigorous replication before drawing definitive conclusions.

3. Results

3.1 Thermal Gradient Data

Sampling Point	Temp (°C)	ΔT from Tajo	Vegetation Structure
Abandoned Tajo (center)	**27.5**	0.0	Bare clay, full sun exposure, no vegetation
Tajo perimeter	26.4	-1.1	Grassland buffer, partial shade
East entrance	23.6	-3.9	Young secondary forest (~1.5 m shrubs)
West entrance	23.5	-4.0	Transition to mature canopy
Mature Hauya elegans tree	21.6	-5.9	Old-growth with buttress roots, dense epiphytes
Reserve deep core	20.0	-7.5	High biomass, closed canopy
The Shrine (forest interior)	**17.3**	**-10.2**	Maximum canopy closure, 20 cm organic layer

Table 1. Temperature measurements along degraded-land-to-mature-forest transect. Bold values indicate maximum cooling effect. ΔT = temperature difference relative to Tajo baseline.

3.2 Ecological Context

During the field session, iNaturalist observations for the Huaravito project increased from 82 to 238, documenting 116 distinct species. Notable indicator species observed within the mature Hauya elegans stands include Coffea arabica (naturalized coffee) and Syzygium jambos (rose apple), both of which indicate stable, moist microclimate conditions. The dense organic carpet (20 cm depth) consists of fine roots, decomposing leaf litter, and extensive white mycelial networks, likely representing ectomycorrhizal or saprotrophic fungal communities.

4. Preliminary Interpretation

4.1 Proposed Mechanisms

The observed 10.2°C gradient appears to be driven by progressive increases in the following structural and functional attributes along the transect:

- Canopy closure: reducing incident solar radiation reaching the ground
- Epiphytic biomass: increasing overall thermal inertia and moisture retention on vertical surfaces
- Organic soil layer depth: insulating mineral soil against radiative heat gain
- Evapotranspiration: latent heat flux from canopy and soil surface
- Wind velocity reduction: preventing advective heat transport into forest interior

4.2 Comparison with Existing Literature

Published studies on tropical forest cooling effects typically report temperature differentials of 3–8°C between forest interiors and adjacent open areas (Hardwick et al. 2015, Agricultural and Forest Meteorology; de Frenne et al. 2019, Global Change Biology). The 10.2°C gradient observed at Huaravito is at the upper end of this range. Possible explanations for the extreme value include:

1. Exceptionally high structural complexity of mature Hauya elegans stands with extensive epiphytic loading
2. Deep organic soil layer (20 cm) providing superior insulation
3. Topographic fog interception at northeast boundary (see Biogenic Cooling Hypothesis document for details)
4. Sampling artifact: single-day observation may have coincided with optimal conditions for thermal decoupling

Distinguishing between these explanations requires temporal replication and measurement of additional environmental covariates.

5. Potential Significance

If the 10.2°C cooling effect is reproducible across multiple sampling dates and weather conditions, this finding would have implications for:

- Climate refugia: Small forest fragments with mature structure may provide thermal microrefugia for heat-sensitive species under regional warming scenarios
- Restoration targets: The data suggest that passive regeneration leading to mature forest structure can yield substantial thermal regulation services, even in small, edge-exposed fragments
- Microclimate monitoring: Temperature gradients may serve as a cost-effective proxy for assessing forest structural recovery and ecosystem function
- Conservation prioritization: Sites exhibiting extreme thermal buffering may warrant protection as “cooling cores” in fragmented landscapes

6. Critical Limitations and Required Next Steps

6.1 Current Limitations

This report documents a single-day, single-measurement observation. The following limitations prevent definitive conclusions:

- No temporal replication: The gradient may not be consistent across days, seasons, or weather conditions
- No spatial replication: Multiple transects are needed to assess site-wide representativeness

- Unknown measurement error: Instrument precision, calibration status, and operator technique not documented
- Confounding variables not controlled: Time of day, cloud movement, wind shifts could all influence individual readings
- No mechanistic data: Canopy cover, humidity, wind speed, and soil temperature were not measured

6.2 Recommended Protocol for Replication Study

To validate and extend this preliminary observation, the following protocol is recommended:

5. Replicate transect on 5–10 dates spanning dry and wet seasons, always at same time of day (e.g., 12:00–13:00)
6. Standardize measurement height (1.5 m above ground) and instrument (calibrated digital thermometer or datalogger)
7. Take 3 replicate readings per point, separated by 2 minutes, and average
8. Record timestamp, cloud cover (%), estimated wind speed, and relative humidity (if sensor available) for each measurement
9. Quantify canopy cover at each point using hemispheric photography or mobile canopy app (e.g., Canopy by PLT)
10. Measure organic layer depth at each forest point (minimum 3 replicates)
11. Install permanent temperature dataloggers at Tajo and Shrine to capture diurnal and seasonal variation
12. Conduct statistical analysis: ANOVA to test for differences among points; linear regression of temperature vs. canopy cover and organic layer depth

7. Working Hypothesis for Future Investigation

Based on five years of qualitative observation and the thermal data presented here, I hypothesize that the extreme cooling effect at Huaravito results from the synergistic interaction of multiple biophysical processes:

- Orographic fog interception: Northeast boundary channels cool, moist air into the reserve (see separate Biogenic Cooling Hypothesis document)
- High epiphytic biomass: >70% trunk coverage increases thermal inertia and moisture retention
- Deep organic soil layer: 20 cm insulating mat prevents radiative soil heating
- Convective circulation: East-to-west air flow driven by temperature and density gradients (speculative; requires tracer studies)

This hypothesis predicts that sites with similar structural attributes (high epiphyte loading, deep organic layers, topographic fog interception) will exhibit comparable thermal buffering, regardless of fragment size. Comparative studies at other premontane forest sites in Costa Rica would allow testing of this prediction.

8. Contextual Note on “Thermal Memory”

In my original field notes, I referred to the 17.3°C at The Shrine as potentially representing the “thermal memory” of pre-colonial primary forests (~500 years ago). While this is a compelling concept, it is speculative in the absence of paleoclimatic data. What can be stated with confidence is that the mature forest core maintains conditions substantially decoupled from contemporary anthropogenic warming in the

surrounding matrix. Whether this represents a "pre-industrial baseline" or simply the maximal cooling achievable by this forest structure remains an open question requiring historical climate reconstruction (e.g., tree ring isotope analysis, pollen core analysis) to address definitively.

9. Conclusion

The 10.2°C thermal gradient documented on February 20, 2026 is a striking preliminary observation that, if confirmed through rigorous replication, would represent one of the most extreme examples of biogenic thermal buffering in a small tropical forest fragment. However, the single-day, single-measurement design of this initial survey precludes definitive conclusions. Systematic temporal and spatial replication, coupled with measurement of structural and microclimatic covariates, is required to:

13. Confirm that the gradient is reproducible
14. Identify the relative contributions of canopy closure, epiphytic biomass, organic layer depth, and evapotranspiration
15. Assess whether the effect varies seasonally or with weather conditions
16. Compare Huaravito's thermal buffering capacity to other premontane fragments

This preliminary report serves as a foundation for designing a more rigorous follow-up study. The potential implications for climate refugia, restoration science, and ecosystem service valuation justify the investment in systematic data collection.

References

de Frenne, P., et al. (2019). Global buffering of temperatures under forest canopies. Nature Ecology & Evolution, 3(5), 744–749.

Hardwick, S. R., et al. (2015). The relationship between leaf area index and microclimate in tropical forest and oil palm plantation: Forest disturbance drives changes in microclimate. Agricultural and Forest Meteorology, 201, 187–195.

Acknowledgment

This observation is the product of daily custodial presence and intimate site knowledge accumulated over five years. While the data collection protocol requires refinement for publication-grade science, the phenomenological pattern — extreme thermal decoupling between degraded and mature forest — is empirically robust and merits systematic investigation.

Randall Santamaría Artavia

Huaravito Long-term Ecological Monitoring Site

Piedades Sur, San Ramón, Costa Rica

February 2026

RESERVA HUARAVITO