Huaravito Reserve
Long-Term Ecological Monitoring Site

(LTEMS)

Monitoring Framework 2026–2031

Integrated Under the Sentinel Protocol

Randall Santamaría Artavia
Custodian-Director
Piedades Sur, San Ramón, Alajuela, Costa Rica

February 2026

"Where the Sentinel Protocol protects the forest, the LTEMS measures its function."

Executive Summary

The Huaravito Reserve Long-Term Ecological Monitoring Site (LTEMS) is an independent ecological monitoring initiative established within the Reserva Huaravito, Costa Rica. The site encompasses approximately 4,500 m² of mature secondary forest, locally known as Bosque Sagrado Toyopán, characterized by a high structural density of Hauya elegans (Guayabón), with an estimated 75–100 mature individuals concentrated within the western core sector. This initiative integrates microclimatic monitoring, forest structural assessment, and long-term biodiversity documentation into a unified data framework operating under the governance principles of the Sentinel Protocol.

Purpose and Origin

In February 2026, preliminary field measurements documented a 10.2°C thermal gradient between an exposed degraded area (former clay extraction zone) and the forest core (Shrine sector). This observation, detailed in the Biogenic Thermal Buffering – Preliminary Report, suggests the presence of a strong microclimatic buffering effect associated with forest structure. Given the magnitude of the observed difference—exceeding typical documented forest cooling ranges (3–8°C)—a structured, multi-year monitoring system is required to verify reproducibility, assess seasonal stability, determine structural drivers, and quantify long-term variance.

The LTEMS is established to provide that structured evidence base. It does not assert conclusions; it establishes the infrastructure necessary to evaluate them.

Core Objective (2026–2031)

To develop and maintain a five-year, replicable, georeferenced ecological monitoring system capable of evaluating microclimatic stability, forest structural dynamics, and species composition trends within the Huaravito Reserve.

1. Institutional Framework: Integration with the Sentinel Protocol

1.1 Governance Alignment

The Huaravito Reserve LTEMS operates under the governance principles defined in the Sentinel Protocol. The Sentinel Protocol establishes:

- Site protection commitment
- Data continuity responsibility
- Structural conservation mandate
- Long-term stewardship ethics

The LTEMS constitutes the scientific monitoring arm of this protocol. Where the Sentinel Protocol defines protection, the LTEMS defines measurement.

1.2 The Sentinel Principle: Detect, Document, Defend

The Monitoring Framework follows three operational pillars derived from the Sentinel Protocol:

A. Detect

Systematic measurement of environmental parameters that indicate ecosystem function:

- Air temperature gradients (microclimatic buffering capacity)
- Structural forest density (canopy complexity and biomass)
- Soil organic layer depth (carbon storage and moisture retention)
- Species richness trajectory (biodiversity stability)

B. Document

Georeferenced data storage through transparent, reproducible protocols:

- Fixed monitoring stations with standardized measurement heights
- Permanent forest plots with re-measurement cycles
- Platform-integrated biodiversity records (iNaturalist, Terraware, Restor)
- Annual summary reporting with full methodological transparency

C. Defend

Evidence-based conservation positioning through longitudinal dataset accumulation:

- Quantifiable metrics for microclimatic refuge status
- Transparent methodological reporting enabling external review
- Institutional readiness for academic or governmental collaboration
- Data-driven advocacy for site preservation under climate change scenarios

2. Site Description

2.1 Geographic and Ecological Context

The LTEMS is located within the Reserva Huaravito, Piedades Sur, San Ramón, Alajuela Province, Costa Rica (10.104811, -84.565125; 1,250 m.a.s.l.). The site encompasses approximately 4,500 m² of mature secondary forest known as Bosque Sagrado Toyopán, classified as premontane humid forest (Holdridge Life Zone system).

2.2 Forest Structure

The western core sector contains an estimated 75–100 mature individuals of Hauya elegans (Guayabón, Onagraceae), forming a structurally dominant canopy system. Key structural attributes include:

- Large-diameter trees (>50 cm DBH estimated) with extensive buttress roots
- >70% epiphytic colonization of trunk surfaces (bryophytes, orchids, ferns)
- Dense organic soil layer (~20 cm depth) with active mycorrhizal networks
- High canopy closure creating multi-layered vertical structure

2.3 Matrix Context

The site borders a historically degraded clay extraction zone (former tajo) to the east, creating a sharp structural and thermal contrast that enables comparative microclimatic monitoring. The surrounding landscape matrix is mixed agriculture and secondary growth, positioning the reserve as an isolated forest fragment along a stream corridor (Quebrada Honda – Quebradillas).

3. Scientific Rationale

3.1 The 10.2°C Observation

Preliminary field measurements conducted on February 20, 2026, documented a 10.2°C temperature differential between exposed degraded terrain (27.5°C) and the forest core at The Shrine (17.3°C) over approximately 150 meters horizontal distance under midday conditions. This magnitude exceeds typical forest buffering values reported in tropical microclimate literature, where differentials of 3–8°C are more commonly observed (Hardwick et al. 2015; de Frenne et al. 2019).

3.2 Research Questions

The LTEMS does not assume causality. It establishes the infrastructure required to test:

- Reproducibility: Is the 10.2°C gradient consistent across multiple sampling dates?
- Seasonal persistence: Does the buffering effect vary between dry and wet seasons?
- Structural dependency: Is the cooling effect correlated with canopy cover, epiphytic biomass, or organic layer depth?
- Interannual stability: Does the forest maintain thermal homeostasis across years despite regional climate variability?

3.3 Hypothesis Framework

The observed thermal gradient is hypothesized to result from synergistic biogenic cooling mechanisms (see Proposed Mechanisms of Biogenic Cooling document):

- Orographic fog interception at the northeast boundary
- Epiphytic-bryophytic thermal buffering on trunk surfaces
- Convective air circulation along the east-west gradient (speculative)
- Evaporative cooling from the dense organic soil layer

The LTEMS monitoring architecture is designed to provide the empirical data required to evaluate these mechanisms.

4. Monitoring Architecture (2026–2031)

The LTEMS integrates four primary monitoring components, each with standardized protocols and re-measurement cycles.

4.1 Microclimate Monitoring Network

Objective

Quantify spatial and temporal variation in air temperature across the degraded-to-mature forest gradient.

Infrastructure

- Three fixed temperature stations: (1) Open area (Tajo baseline), (2) Transition zone (forest edge), (3) Forest core (The Shrine)
- Standardized measurement height: 1.5 m above ground
- Data logging frequency: Continuous (dataloggers) or weekly (manual readings, if continuous logging unavailable)
- Primary metric: ΔT (temperature differential) between open area and forest core

Protocol

- Calibrate all sensors against a reference thermometer at site initiation and annually thereafter
- Record timestamp, cloud cover (%), estimated wind speed, and relative humidity (if sensor available) for each measurement
- Calculate monthly mean temperatures for each station
- Analyze dry season (Dec–Apr) vs. wet season (May–Nov) thermal regimes

Target

≥ 48 months of structured thermal data by 2031, enabling interannual variance analysis.

4.2 Forest Structure Plot

Objective

Establish a permanent vegetation plot to monitor tree density, size distribution, and species composition over time.

Infrastructure

- Permanent 30 × 30 m plot established within the western core sector (high H. elegans density)
- Plot corners marked with permanent stakes and GPS coordinates recorded
- All trees ≥10 cm DBH tagged with unique identifiers

Protocol

- Measure DBH (cm) at 1.3 m height using diameter tape

- Estimate total height (m) using clinometer or visual estimation (document method)
- Confirm species identification with photographic voucher uploaded to iNaturalist
- Record health status: herbivory score (0–4 scale), foliage color, structural damage
- Calculate density (trees/ha) for H. elegans and all species combined

Re-measurement Cycle

Every 2 years (2026, 2028, 2030). Intermediate annual walk-throughs to document mortality or major structural changes.

Target

Complete tree census of the core forest plot with ≥3 re-measurement cycles by 2031.

4.3 Soil and Organic Layer Assessment

Objective

Monitor organic matter accumulation and soil surface conditions as indicators of long-term carbon storage and moisture retention capacity.

Infrastructure

- 10 permanent sampling points distributed across the 30 × 30 m plot
- Points marked with subsurface markers (e.g., buried metal washers) and GPS coordinates

Protocol

- Measure organic layer depth (cm) using a graduated metal rod inserted vertically until resistance from mineral soil is encountered
- Take 3 replicate measurements per point and average
- Photograph soil surface condition (litter type, fungal presence, root visibility)
- Record qualitative notes: moisture status, compaction, disturbance signs

Re-measurement Cycle

Annual (each February, coinciding with dry season to minimize seasonal variability).

Target

≥5 years of organic layer depth data enabling trend analysis.

4.4 Biodiversity Documentation

Objective

Track species richness trajectory and maintain a comprehensive, publicly verifiable biodiversity inventory.

Infrastructure

- Continued use of iNaturalist as primary documentation platform (Project: Reserva Huaravito / Sitio Natural Sagrado)
- Cross-referencing with Terraware (organizationId=213), Restor, and NASA GLOBE Observer (Trees module) for multi-platform validation

Protocol

- Upload all field observations with GPS coordinates, timestamp, and photographic voucher
- Target taxonomic verification to species level where possible, genus level minimum
- Track annual totals: total observations, unique species count, H. elegans observations
- Calculate species accumulation curve to assess sampling completeness

Target

≥300 total observations and ≥150 unique species documented by 2031.

5. Governance and Ethical Commitments

5.1 Sentinel Protocol Compliance

Under the Sentinel Protocol, the LTEMS commits to:

- Non-extractive monitoring practices: No collection of plant material beyond increment cores (if dendrochronology is conducted) or voucher specimens for taxonomic verification
- Structural canopy preservation: No pruning, clearing, or intentional modification of forest structure
- Microclimate integrity: No intentional alteration of core microclimate (e.g., no experimental tree removal, no artificial ventilation)
- Transparent documentation: Full methodological disclosure in all reports, including changes to protocols and reasons for modifications

The monitoring system is adaptive but conservative. Changes to protocols require documented justification and do not compromise longitudinal data comparability.

5.2 Data Governance

The LTEMS operates as an open-data-aligned monitoring site:

- All datasets are structured to enable external review and academic collaboration
- Raw data (temperature logs, tree census spreadsheets, soil measurements) are maintained in archival formats (CSV, TXT) with metadata documentation
- Annual summary reports are prepared for public release and institutional sharing
- Data sharing agreements with academic or governmental collaborators will specify: (a) anonymization of site-specific results in publications, (b) generalized location data (regional level) in public datasets, (c) opportunity for the Custodian-Director to review manuscripts prior to submission

5.3 Institutional Readiness

While the LTEMS is currently an independent initiative, its data architecture is designed for potential integration with:

- University research programs (e.g., UCR, UNA, OTS, EARTH)
- National biodiversity monitoring networks (e.g., SINAC, INBio legacy databases)
- International restoration monitoring initiatives (e.g., SNAPP-MORE)
- Climate refugia mapping projects

6. Five-Year Targets and Deliverables (2026–2031)

By December 2031, the LTEMS aims to achieve the following measurable outcomes:

6.1 Microclimatic Dataset

- ≥ 48 months of continuous or semi-continuous temperature data from all three stations (open area, transition, core)
- Quantified mean annual ΔT value with 95% confidence interval
- Interannual variance analysis (σ) demonstrating thermal stability or identifying trends
- Seasonal decomposition: dry season vs. wet season thermal regimes

6.2 Forest Structure Dataset

- Complete structural census of the 30 × 30 m permanent plot with ≥3 re-measurement cycles
- DBH distribution histogram for all species and H. elegans separately
- Density (trees/ha) calculation with temporal comparison (2026 vs. 2028 vs. 2030)
- Mortality and recruitment rates documented

6.3 Soil Organic Layer Dataset

- ≥5 years of annual organic layer depth measurements
- Mean depth with standard deviation calculated across 10 sampling points
- Trend analysis: stable, increasing, or decreasing organic matter accumulation

6.4 Biodiversity Dataset

- ≥300 total iNaturalist observations
- ≥150 unique species documented
- Species accumulation curve with asymptote assessment
- H. elegans representation trajectory (% of total observations per year)

6.5 Technical Synthesis Report

- Publication of a formal Technical Synthesis Report (2031) consolidating five years of monitoring data
- Report to include: site description, methods, results, statistical analyses, discussion of microclimatic buffering hypothesis, limitations, and recommendations for future monitoring
- Report prepared in format suitable for submission to peer-reviewed journals (e.g., Brenesia, Revista de Biología Tropical) or institutional repositories

7. Strategic Position and Long-Term Vision

7.1 The LTEMS Role

The Sentinel Protocol protects the forest. The LTEMS measures its function. Together, they establish Huaravito as:

- A locally governed ecological observatory: Demonstrating that small-scale, custodian-led monitoring can generate scientifically rigorous datasets
- A microclimate buffering case study: Providing empirical evidence for exceptional thermal regulation in a small, edge-exposed forest fragment
- A potential long-term climate refugium candidate: Contributing data to regional assessments of sites likely to maintain stable microclimates under future warming scenarios

7.2 Beyond 2031

The five-year framework (2026–2031) is intended as a foundation, not a terminus. If microclimatic stability and forest structural integrity are maintained, the LTEMS is designed for indefinite continuation. Long-term monitoring (10+ years) would enable:

- Decadal trend analysis of thermal buffering capacity under regional climate change
- Generational forest dynamics: recruitment, growth, and mortality of H. elegans cohorts
- Species composition shifts in response to altered temperature or precipitation regimes
- Comparative analysis with other premontane forest sites in Costa Rica

7.3 Core Principle

The Huaravito LTEMS does not assert conclusions. It establishes the infrastructure necessary to evaluate them. It operates at the intersection of local stewardship, empirical ecological monitoring, and long-term climate buffering research.

References

de Frenne, P., et al. (2019). Global buffering of temperatures under forest canopies. Nature Ecology & Evolution, 3(5), 744–749.

Hardwick, S. R., et al. (2015). The relationship between leaf area index and microclimate in tropical forest and oil palm plantation. Agricultural and Forest Meteorology, 201, 187–195.

Santamaría Artavia, R. (2026). Biogenic Thermal Buffering Along a Forest Maturity Gradient at Huaravito Reserve: Preliminary Field Observation Report. Huaravito Reserve LTEMS.

Document prepared by:

Randall Santamaría Artavia

Custodian-Director

Huaravito Reserve Long-Term Ecological Monitoring Site

Piedades Sur, San Ramón, Costa Rica

February 2026