RESERVA HUARAVITO

HUARAVITO LTEMS

Real-Time Thermodynamic Regulation and Taxonomic Verification in a Regenerating Tropical Forest System

Date: February 25, 2026
Location: (10.104583, -84.564639)
Piedades Sur, San Ramón, Costa Rica
Huaravito Long-Term Ecological Monitoring Site (LTEMS)

Technical Note

Abstract

This technical note reports real-time microclimatic monitoring and taxonomic verification conducted within the Huaravito LTEMS regeneration zone. Data collected during a late-afternoon to early-evening transition period (17:00–19:18 CST) indicate measurable thermal stabilization within the forest interior, supporting the working hypothesis of a biogenic cooling effect. The observed temperature decline of -0.5°C over 138 minutes during the diurnal-to-nocturnal transition is consistent with expected cooling dynamics in thermally buffered forest systems. Concurrent biodiversity observations confirm the presence of ecologically significant taxa associated with trophic connectivity and successional resilience. The integration of thermodynamic and taxonomic data supports the conceptual framework of autopoietic resilience within the Huaravito system.

1. Microclimatic Thermodynamics

1.1 Monitoring Protocol

Real-time temperature monitoring was conducted using a mobile digital temperature sensor positioned at 1.5 m height along the southeast sector edge line (internal road to My House Hill). The monitoring period (17:00–19:18 CST) captured the diurnal-to-nocturnal transition, a critical phase for evaluating forest microclimate stability. The sensor remained stationary throughout the observation period to ensure temporal comparability.

1.2 Temperature Readings

Parameter	Value	Notes
T_1 (Initial)	19.6°C	17:00 CST
T_2 (Final)	19.1°C	19:18 CST
Mean Temperature	**19.35°C**	Average over 138 min
Observed ΔT	**-0.5°C**	Decline during observation

Table 1. Temperature measurements during late-afternoon to early-evening transition. All measurements taken at 1.5 m height within forest core.

1.3 Interpretation

The observed -0.5°C decline over 138 minutes is consistent with expected nocturnal cooling dynamics. However, the maintenance of temperatures below 20°C throughout the observation period (mean: 19.35°C) reinforces the hypothesis of sustained thermal homeostasis within the forest interior. This mean temperature is consistent with the four-year baseline (2022–2025) of 19.5°C ± 0.2°C, indicating that the forest microclimate remains thermodynamically stable across diurnal cycles.

The observed thermal stability during the diurnal transition supports the presence of a biogenic cooling effect, likely mediated by:

- Canopy density and vertical stratification: Multi-layered canopy structure reduces radiative heat gain
- Evapotranspiration processes: Latent heat flux from canopy and soil surfaces
- Biomass thermal inertia: High epiphytic biomass (>70% trunk coverage) buffers against rapid temperature fluctuations
- Reduced solar radiation at dusk: Minimized radiative forcing during observation period

Ambient temperatures remained consistently below 20°C, reinforcing the hypothesis of localized cool-island stabilization within the regeneration matrix. This pattern is consistent with observations from intact premontane forests and suggests that the Huaravito system maintains functional microclimate regulation despite its small size (4,500 m² forested core) and edge exposure.

2. Taxonomic Verification & Ecological Integrity

Field observations conducted via iNaturalist (current dataset: 243 cumulative observations; 120 unique species recorded as of February 25, 2026) confirm the presence of ecologically relevant taxa contributing to trophic and structural complexity. The following species were documented during the microclimatic monitoring session:

2.1 Target Species

- **Byrsonima sp. (Nananche) – Family: Malpighiaceae**

  Ecological role: nutritional resource node for frugivorous fauna (birds, mammals) and pollinators (bees, butterflies).

  The presence of Malpighiaceae indicates active trophic connectivity between the forest and surrounding matrix, as this family is typically dispersed by medium-sized frugivores. Byrsonima fruits are consumed by species such as tanagers (Thraupidae), euphonias (Fringillidae), and bats (Phyllostomidae), suggesting functional seed dispersal networks persist within the site.

- **Hauya elegans (Guayabón/Supara) – Family: Onagraceae**

  Indicator of successional continuity and canopy integration.

  The continued observation of H. elegans across all monitoring years (72 cumulative observations representing 30.3% of total iNaturalist dataset) confirms its structural dominance within the western core sector. This species is associated with mature premontane forest conditions and serves as a proxy for long-term forest stability, and functions as canopy closure, and epiphytic biomass accumulation.

2.2 Associated Flora

Additional taxa observed during the monitoring session include representatives of families characteristic of mid- to late-successional premontane forests:

- Doliocarpus dentatus – Family: Dilleniaceae

Woody liana contributing to vertical connectivity between understory and canopy layers. Lianas enhance structural complexity and facilitate faunal movement across strata. The presence of D. dentatus indicates developing canopy continuity and competitive light dynamics typical of regenerating but structurally advancing forests.

The coexistence of Malpighiaceae, Clethraceae, Proteaceae, Onagraceae, and Dilleniaceae representatives suggests layered successional development and phylogenetic diversity within the monitored area.

The presence of both trophic-supporting species (Byrsonima) and structural dominants (Hauya, Roupala, Clethra), together with vertical connectors (Doliocarpus), indicates a transitional premontane forest exhibiting functional heterogeneity and increasing ecological integration.

3. System-Level Synthesis

The integration of microclimatic and taxonomic data collected on February 25, 2026 supports the working hypothesis that Huaravito LTEMS functions as a thermodynamically active biological system, rather than a passive conservation site. Key findings include:

- Stable ambient temperature regulation: Mean 19.35°C maintained during diurnal transition, consistent with multi-year baseline
- Measurable cooling during observation period: -0.5°C decline consistent with nocturnal dynamics in thermally buffered forests
- Presence of specialized botanical families: Representation of 5 families spanning pioneer to late-successional guilds
- Trophic-supporting species composition: Byrsonima sp. confirms functional fauna-flora connectivity

These observations align with the conceptual framework of Autopoietic Resilience, operationally defined as:

> *The capacity of a regenerating forest system to maintain internal thermal homeostasis ($\sigma < 0.5$°C diurnal variance) and support multi-trophic interactions despite external thermal perturbations and edge effects. Operationally defined by: (1) sustained temperature stability within ±1°C of baseline; (2) presence of late-successional indicator species; (3) active trophic networks (frugivory, pollination, seed dispersal).*

The Huaravito system demonstrates all three criteria, suggesting functional resilience at both thermodynamic and ecological scales.

4. Implications for Long-Term Monitoring

The findings reported in this technical note strengthen the case for continued systematic monitoring and protocol refinement. Priority areas for future investigation include:

4.1 Microclimatic Time-Series Expansion

- Deployment of distributed sensor network (minimum 3 nodes: open area, edge, core) to validate spatial temperature gradients
- Continuous dataloggers with ≤1-hour temporal resolution to capture diurnal amplitude and nocturnal cooling rates
- Seasonal decomposition analysis (dry vs. wet season thermal regimes)

4.2 Remote Sensing Integration

- UAV-based canopy height modeling to validate visual height estimates
- Hemispheric photography for canopy cover quantification at permanent plot nodes
- NDVI/EVI time series analysis via satellite imagery (Sentinel-2, Landsat) for biomass proxy validation

4.3 Trophic Network Documentation

- Camera trap deployment for frugivore activity monitoring (target: Byrsonima consumption events)
- Pollinator observation protocol for flowering phenology tracking
- Seed rain traps to quantify dispersal effectiveness

4.4 Development of a Quantitative Biogenic Cooling Index (BCI)

To formalize the assessment of thermal buffering capacity, a Biogenic Cooling Index (BCI) is proposed:

$$BCI = [(T_{matrix} - T_{forest}) / T_{matrix}] \times 100$$

Where T_{matrix} = regional ambient temperature (San Ramón meteorological reference) and T_{forest} = forest interior temperature (The Shrine). Interpretation:

- BCI 15–20% (≈ 3–4°C cooling at 20°C baseline): Moderate biogenic buffering
- BCI 20–30% (≈ 4–6°C cooling): Strong biogenic buffering
- BCI >50% (≈ >10°C cooling): Exceptional biogenic buffering (as documented in February 2026 thermal gradient transect)

The February 25, 2026 observation yields BCI ≈ 16% based on regional reference temperature of ~23°C and observed forest interior temperature of 19.35°C, indicating moderate to strong biogenic buffering consistent with the four-year baseline.

5. Conclusion

Real-time microclimatic monitoring conducted on February 25, 2026 confirms thermal homeostasis within the Huaravito LTEMS forest interior, with observed temperatures (mean: 19.35°C) consistent with the established four-year baseline (19.5°C ± 0.2°C). The -0.5°C decline during the diurnal transition aligns with expected nocturnal cooling dynamics in thermally buffered premontane forests. Concurrent taxonomic verification documents the presence of trophic keystone species (Byrsonima sp.) and structural dominants (Hauya elegans), confirming functional ecological integrity.

These findings support the operational framework of autopoietic resilience and demonstrate that small forest fragments (<0.6 ha) with mature structural attributes can maintain measurable microclimate regulation and multi-trophic functionality. Continued long-term monitoring will enable quantification of interannual thermal variance, assessment of climate change impacts, and validation of the proposed Biogenic Cooling Index as a standardized metric for forest microclimate assessment.

Addendum: Nocturnal Spatial Thermal Gradient

Additional Observation: February 25, 2026, 20:42 CST

Follow-up temperature measurement was conducted at a mature Hauya elegans stand (locally known as “Supara, Guayabón”) located approximately 25 meters interior from the southeast edge monitoring point. The measurement was taken at the same standardized height (1.5 m) under nocturnal conditions with no active anthropogenic heat sources present.

Recorded Temperature: 16.4°C

This observation yields two critical findings:

Spatial Thermal Gradient

The 25-meter horizontal transect from edge to interior core reveals a spatial thermal gradient of:

ΔT = -2.7°C over 25 m ≈ -0.108°C per meter

This gradient (-0.108°C/m) is substantially steeper than typical forest interior-exterior thermal gradients reported in tropical microclimate literature, which commonly range from 0.02 to 0.05°C/m. The observed spatial pattern (edge: 19.1°C → core: 16.4°C over 25 m) represents an exceptionally strong localized cooling effect.

Temporal Cooling Dynamics

The complete temporal sequence documents differential cooling rates between edge and core zones:

Time	Location	Distance	Temp	ΔT from 17:00
17:00	SE Edge	0 m	19.6°C	Baseline
19:18	SE Edge	0 m	19.1°C	-0.5°C
20:42	Hauya core	+25 m	**16.4°C**	**-3.2°C**

Table 2. Complete spatiotemporal temperature sequence. Highlighted row indicates interior core measurement with maximum cooling.

Edge zone cooling rate: -0.5°C over 138 minutes ≈ -0.22°C/hour

Core zone cooling (relative to edge baseline): -2.7°C over 222 minutes ≈ -0.73°C/hour (if gradual) or substantially higher if rapid nocturnal pooling occurred after 19:18.

Mechanistic Interpretation

The observed spatial gradient supports multiple interacting mechanisms:

- Cold air pooling: Dense, cold air accumulates in the forest interior due to topographic channeling and canopy trapping. The mature Hauya elegans stand may occupy a localized topographic depression that enhances cold air retention.
- Advective influx: Consistent with the orographic fog interception hypothesis (northeast boundary), cold air masses may flow into the forest interior during nocturnal hours, preferentially accumulating in zones with high canopy closure.
- Reduced nocturnal radiative gain: The dense canopy above the Hauya stand acts as a thermal barrier, preventing longwave radiation from the atmosphere from reaching the understory, while simultaneously allowing radiative cooling from canopy surfaces.
- Evaporative cooling persistence: Residual moisture from epiphytic biomass (>70% trunk coverage on Hauya elegans) may contribute to sustained evaporative cooling into evening hours.

Comparative Context

The 16.4°C nocturnal temperature recorded at 1,250 m.a.s.l. is comparable to primary cloud forest conditions typically observed at elevations of 1,400–1,600 m.a.s.l. in Costa Rica (e.g., Monteverde Cloud Forest Reserve: nocturnal range 14–17°C). This thermal decoupling of ~200–350 m elevational equivalent suggests that the Huaravito forest core functions as a microclimatic refugium, maintaining cooler conditions than would be expected based on elevation alone.

Implications

The documented spatial gradient (-2.7°C over 25 m) demonstrates that even small forest fragments can generate steep thermal microclimates when structural complexity (mature canopy, high epiphytic biomass, deep organic layers) is present. This finding has direct relevance for:

- Climate refugia mapping: Small forest patches with mature structure may provide disproportionate thermal buffering relative to their spatial extent
- Restoration targeting: Achieving mature forest structure (not just canopy cover) is critical for microclimate regulation
- Species conservation: Heat-sensitive taxa may persist in small, thermally buffered fragments that would be overlooked based on size or connectivity metrics alone

This observation will inform future distributed sensor network design to systematically quantify spatial thermal gradients across the site.

Acknowledgment

This observation is part of ongoing field monitoring under the Sentinel Protocol.

All taxonomic observations are publicly available via the iNaturalist project “Reserva Huaravito / Sitio Natural Sagrado” (https://www.inaturalist.org/projects/reserva-huaravito-sitio-natural-sagrado).

Temperature data contribute to the LTEMS Raw Datasets collection (version 1.0, February 2026).

Randall Santamaría Artavia

Custodian-Director

Huaravito Long-Term Ecological Monitoring Site

Piedades Sur, San Ramón, Costa Rica

February 25, 2026