EXECUTIVE SUMMARY

Biogenic Microclimatic Buffering and Passive Forest Restoration at Huaravito Reserve LTEMS

Custodian-Director: Randall Santamaría Artavia
Huaravito Long-Term Ecological Monitoring Site (LTEMS)
Piedades Sur, San Ramón, Costa Rica
10.104901°N, 84.565024°W | 1,250 m a.s.l.
February 2026

The Huaravito Reserve LTEMS (0.57 ha, mature core 0.4 ha), embedded within the Reserva de Biosfera Cordillera Volcánica Central (UNESCO-MAB) and Corredor Biológico Montes del Aguacate, serves as a reference site for long-term ecological monitoring and passive restoration in premontane-montane tropical forest fragments. Five years of structured field observations (2021–2026) and four years of continuous microclimatic monitoring (2022–2025) reveal persistent thermal homeostasis (interannual variance $\sigma \approx 0.2$°C) and a sustained biogenic cooling effect of -3.7°C relative to the surrounding anthropogenically disturbed matrix. A preliminary midday thermal transect (February 20, 2026) documented a maximum gradient of 10.2°C (27.5°C at the abandoned clay extraction site [Tajo] vs. 17.3°C in the mature forest core [The Shrine]), placing Huaravito at the upper end of reported biogenic cooling magnitudes in tropical forest literature (typically 3–8°C).

Key Scientific Highlights

1. Exceptional Thermal Buffering Capacity

- Mean annual core temperature: 19.5°C ($\sigma \approx 0.2$°C interannual)
- Sustained cooling: -3.7°C vs. adjacent matrix; extreme-event buffering: +6.0°C during February 2026 polar intrusion
- Transect gradient (150 m, 7 points): progressive decline from exposed clay (27.5°C) through secondary regeneration to mature canopy (17.3°C), driven by increasing structural complexity and functional attributes

2. Proposed Biogenic Heat Exchanger Hypothesis

A mechanistic framework integrates four interdependent processes:

- (1) Orographic fog interception at the NE boundary, channeled by topographic depression from an abandoned mine; mid-canopy trees (e.g., Hauya elegans) and dense epiphytic load (mosses, orchids, ferns) facilitate condensation and evaporative cooling
- (2) Epiphytic-bryophytic thermal buffering: >70% trunk surface coverage increases thermal inertia, insulates cambial tissues, and reduces wind penetration via secondary shade layers (vines/lianas)
- (3) Convective displacement along an E-W gradient ('chimney effect'): density-driven circulation evacuates warmer air through canopy gaps while maintaining cool understory conditions
- (4) Soil-atmosphere interface ('living crust'): dense 20-cm organic horizon (fine roots, decomposing litter, fungal hyphae/mycelia) enhances evaporative cooling and nutrient cycling

All mechanisms are consistent with established forest microclimate theory but remain speculative pending distributed instrumentation (temperature/humidity loggers, anemometers, fog collectors, infrared thermography, tracer gas/smoke dispersion tests).

3. Sentinel Role of Hauya elegans (Onagraceae)

- Dominant in the mature core (72/238 iNaturalist records; 30.3% of total; 116 species documented)
- Population of >100 mature individuals — exceptional density in the Valle Central, where the species persists only as scattered, senescent individuals in riverine barrancos (e.g., Río Virilla)
- Structural traits (buttress roots, high epiphytic colonization) facilitate fog interception, thermal inertia, and microclimatic stability, positioning H. elegans as a key facilitator in passive restoration and a sentinel species for monitoring success in fragmented premontane systems

4. Conservation and Scientific Significance

- The site exemplifies passive regeneration (non-assisted recruitment, closed multi-layered canopy, high biomass) within a human-dominated matrix, contributing to national (SINAC guidelines) and global targets (UN Decade on Ecosystem Restoration 2021–2030, SDG 13/15, CBD Kunming-Montreal Targets 2/3)
- Integrated low-impact Field Lodge infrastructure supports LTEMS continuity, participatory biodiversity documentation, and regenerative design demonstration
- Data quality: high-resolution time series and transect records; limitations (single-sensor monitoring, lack of spatial/temporal replication) highlight the need for expanded instrumentation networks

This preliminary evidence underscores Huaravito LTEMS as a model for quantifying biogenic cooling as a proxy of restoration success in small, fragmented tropical forests under climate pressure. Systematic replication and multi-sensor deployment are recommended to test mechanistic predictions and scale findings regionally.



Online visualization:
https://claude.ai/public/artifacts/2fd66654-ca14-4920-bbf2-10b3ae9b46e3

For complete technical documentation, see:

- Monitoring Framework 2026–2031
- Biogenic Thermal Buffering – Preliminary Report
- Hauya elegans Population Analysis
- Data Annex (Raw Datasets v1.0)
- Basal Area Analysis

Contact: Randall Santamaría Artavia
Huaravito Reserve LTEMS | Piedades Sur, Costa Rica