

EDS241: Assignment 3

Shale Hunter

02/18/2022

1 Load Data

```
smoke = read_csv(here("HW3/SMOKING_EDS241.csv"))
```

2 Questions

- a) What is the unadjusted mean difference in birth weight of infants with smoking and non- smoking mothers? Under what hypothesis does this correspond to the average treatment effect of maternal smoking during pregnancy on infant birth weight? Provide some simple empirical evidence for or against this hypothesis.

The unadjusted mean difference in birth weight of infants with smoking and non- smoking mothers is -244.5393875grams. This corresponds to the ATE under the assumption that mothers are assigned randomly to the smoking/non-smoking group. This is probably not a strong assumption because it is likely that there are external factors that effect a mother's likelihood to smoke during pregnancy, such as (only thinking about other variables in our dataset) whether or not this is the mother's first child (mothers might know less about the harmful effects of smoking when they are on their first child) or mother's education level (women with more education may be less likely to smoke because of a greater awareness of the harmful effects of smoking). Both these effects are shown to be significant in the models below, showing that smoking is not random across these other conditions.

```
summary(estimatr::lm_robust(data = smoke, tobacco ~ first))
```

```
##
## Call:
## estimatr::lm_robust(formula = tobacco ~ first, data = smoke)
##
## Standard error type: HC2
##
## Coefficients:
##           Estimate Std. Error t value Pr(>|t|) CI Lower CI Upper    DF
## (Intercept)  0.2120   0.001752  120.98 0.000e+00  0.20858  0.21545 94171
## first       -0.0456   0.002561  -17.81 8.202e-71 -0.05062 -0.04058 94171
##
## Multiple R-squared:  0.003261 , Adjusted R-squared:  0.00325
## F-statistic: 317.1 on 1 and 94171 DF, p-value: < 2.2e-16
```

```
summary(estimatr::lm_robust(data = smoke, tobacco ~ meduc))
```

```
##
## Call:
## estimatr::lm_robust(formula = tobacco ~ meduc, data = smoke)
```

```
##
## Standard error type: HC2
##
## Coefficients:
##           Estimate Std. Error t value Pr(>|t|) CI Lower CI Upper DF
## (Intercept)  0.78933  0.0077083  102.40      0  0.77423  0.80444 94171
## meduc       -0.04594  0.0005421  -84.75      0 -0.04701 -0.04488 94171
##
## Multiple R-squared:  0.06057 , Adjusted R-squared:  0.06056
## F-statistic:  7183 on 1 and 94171 DF, p-value: < 2.2e-16
```

- b) Assume that maternal smoking is randomly assigned conditional on the observable covariates listed above. Estimate the effect of maternal smoking on birth weight using a linear regression. Report the estimated coefficient on tobacco and its standard error.

```
tbco_lm = summary(estimatr::lm_robust(data = smoke, birthwgt ~ .))
tbco_lm
```

```
##
## Call:
## estimatr::lm_robust(formula = birthwgt ~ ., data = smoke)
##
## Standard error type: HC2
##
## Coefficients:
##           Estimate Std. Error t value Pr(>|t|) CI Lower CI Upper DF
## (Intercept) 3362.258  12.0765 278.4133 0.000e+00 3338.588 3385.92805 94164
## anemia      -4.796  17.8739 -0.2683 7.884e-01 -39.829  30.23630 94164
## diabete     73.228  13.2355  5.5327 3.162e-08  47.286  99.16895 94164
## tobacco    -228.073   4.2768 -53.3282 0.000e+00 -236.456 -219.69063 94164
## alcohol    -77.350  14.0392 -5.5096 3.607e-08 -104.866 -49.83312 94164
## mblack     -240.030   5.3478 -44.8842 0.000e+00 -250.512 -229.54873 94164
## first      -96.944   3.4880 -27.7934 2.528e-169 -103.781 -90.10763 94164
## mage       -0.694   0.3682 -1.8849 5.944e-02  -1.416   0.02764 94164
## meduc       11.688   0.8618  13.5630 7.262e-42   9.999  13.37742 94164
##
## Multiple R-squared:  0.0717 , Adjusted R-squared:  0.07162
## F-statistic: 877.6 on 8 and 94164 DF, p-value: < 2.2e-16
```

The model above predicts a coefficient of -228.07 (grams) for tobacco, with a standard error of 4.27 grams.

- c) Use the exact matching estimator to estimate the effect of maternal smoking on birth weight. For simplicity, consider the following covariates in your matching estimator: create a 0-1 indicator for mother's age (=1 if mage>=34), and a 0-1 indicator for mother's education (1 if meduc>=16), mother's race (mblack), and alcohol consumption indicator (alcohol). These 4 covariates will create $2*2*2*2 = 16$ cells. Report the estimated average treatment effect of smoking on birthweight using the exact matching estimator and its linear regression analogue (Lecture 6, slides 12-14).

```
# dummy variables
age = as.numeric(as.logical(smoke$mage >= 34))
edu = as.numeric(as.logical(smoke$meduc >= 16))
blk = smoke$mblack
alc = smoke$alcohol

# grouped variable
grp = paste0(age, edu, blk, alc)
```

```
smoke = smoke %>% mutate(grp = grp)

analogue_lm = estimatr::lm_robust(data = smoke, birthwgt ~ tobacco + factor(grp))
EMcompare = summary(analogue_lm)
```

2.0.1 Exact Matching

```
treatment_table <- smoke %>%
  group_by(grp,tobacco)%>%
  # Calculate number of observations and Y mean by X by treatment cells:
  summarise(n_obs = n(),
            Y_mean = mean(birthwgt, na.rm = T))%>%
  #old way to pivot_longer: gather(variables, values, n_obs:Y_mean)
  pivot_longer(names_to = "variables", values_to = "values", n_obs:Y_mean) %>%
  # Combine the treatment and variables for re-resaping
  mutate(variables = paste0(variables, "_", tobacco))%>%
  # Reshape data by treatment and X cell
  pivot_wider(id_cols = grp, names_from = variables, values_from = values)%>%
  ungroup()%>% #Ungroup from X values
  mutate(Y_diff = Y_mean_1 - Y_mean_0, #calculate Y_diff
         w_ATE = (n_obs_0+n_obs_1)/(sum(n_obs_0)+sum(n_obs_1)),
         w_ATT = n_obs_1/sum(n_obs_1))%>% #calculate weights
  mutate_if(is.numeric, round, 2) #Round data

stargazer(treatment_table, type= "text", summary = FALSE, digits = 2)
```

```
##
## =====
##      grp  n_obs_0 Y_mean_0 n_obs_1 Y_mean_1 Y_diff  w_ATE w_ATT
## -----
## 1  0000  44274  3445.69  13443  3220.25  -225.44  0.61  0.74
## 2  0001   214  3450.28   448  3124.25  -326.03  0.01  0.02
## 3  0010  7007  3195.97  1980  3006.31  -189.66  0.1  0.11
## 4  0011   71  3120.07   226  2817.34  -302.73  0  0.01
## 5  0100 13425  3483.02   535  3273.94  -209.08  0.15  0.03
## 6  0101  130  3510.95    29  3413.21  -97.74  0  0
## 7  0110  625  3319.22    61  3159.05  -160.17  0.01  0
## 8  0111    4  2983.5    10  3097.7  114.2  0  0
## 9  1000  5115  3467.41   976  3171.42  -295.98  0.06  0.05
## 10 1001   56  3358.32    45  3097.73  -260.59  0  0
## 11 1010  396  3185.08   135  2994.67  -190.41  0.01  0.01
## 12 1011    7  2739.71    26  2846.38  106.67  0  0
## 13 1100  4492  3487.19   201  3249.45  -237.74  0.05  0.01
## 14 1101   57  3534.91    17  3037.47  -497.44  0  0
## 15 1110  147  3328.29    19  2852.16  -476.13  0  0
## 16 1111    1  3459     1  2835  -624  0  0
## -----
## # MULTIVARIATE MATCHING ESTIMATES OF ATE AND ATT
## ATE=sum((treatment_table$w_ATE)*(treatment_table$Y_diff))
## ATE
```

```
## [1] -224.2583
```

```
ATT=sum((treatment_table$w_ATT)*(treatment_table$Y_diff))
ATT
```

```
## [1] -222.589
```

The estimated average treatment effect of smoking on birthweight using the exact matching estimator is **-224.2583** grams, while the comparable linear regression gives an average treatment effect of **c(tobacco = -226.245032864622)** grams. These values are quite close.

- d) Estimate the propensity score for maternal smoking using a logit estimator and based on the following specification: mother's age, mother's age squared, mother's education, and indicators for mother's race, and alcohol consumption.

```
smoke = smoke %>% mutate(mage2 = (mage * mage))

ps_model = glm(data = smoke, formula = tobacco ~ mage + mage2 + meduc + mblack + alcohol, family = binomial)
propensity = predict(ps_model, type = "response")
```

- e) Use the propensity score weighted regression (WLS) to estimate the effect of maternal smoking on birth weight (Lecture 7, slide 12).

```
ps_wt = (smoke$tobacco / propensity) + ((1-smoke$tobacco) / (1-propensity))

wps_model = estimatr::lm_robust(data = smoke, birthwgt ~ tobacco + mage + (mage * mage) + meduc + mblack)
summary(wps_model)
```

```
##
## Call:
## estimatr::lm_robust(formula = birthwgt ~ tobacco + mage + (mage *
##     mage) + meduc + mblack + alcohol, data = smoke, weights = ps_wt)
##
## Weighted, Standard error type: HC2
##
## Coefficients:
##              Estimate Std. Error t value    Pr(>|t|)    CI Lower CI Upper    DF
## (Intercept)  3309.739    22.4120  147.677  0.000e+00  3265.8114  3353.666  94167
## tobacco      -220.909     5.0576  -43.679  0.000e+00  -230.8219 -210.996  94167
## mage           1.310     0.5178   2.531  1.137e-02    0.2957   2.325  94167
## meduc          8.316     1.5593   5.333  9.676e-08    5.2596  11.372  94167
## mblack       -225.628     8.2190  -27.452  2.964e-165 -241.7373 -209.519  94167
## alcohol       -69.401    16.7878  -4.134  3.567e-05 -102.3052  -36.498  94167
##
## Multiple R-squared:  0.07265 ,    Adjusted R-squared:  0.0726
## F-statistic: 887.5 on 5 and 94167 DF,  p-value: < 2.2e-16
```

The effect of maternal smoking on birth weight is **-220.9** grams.