
endothelial cell
adipocyte
Schwann cell
glial cell
chondrocyte
pericyte
immature B cell
naive B cell
memory B cell
plasma cell
transitional stage B cell
precursor B cell
pro-B cell
smooth muscle cell
osteoblast
basophil
stromal cell
lymphocyte of B lineage
fibroblast
common myeloid progenitor
common lymphoid progenitor
granulocyte monocyte progenitor cell
megakaryocyte progenitor cell
early lymphoid progenitor
hematopoietic stem cell
megakaryocyte-erythroid progenitor cell
megakaryocyte
hematopoietic multipotent progenitor cell
mesenchymal stem cell
bone marrow
erythroid lineage cell
Homo sapiens
immature conventional dendritic cell
conventional dendritic cell
dendritic cell, human
plasmacytoid dendritic cell, human
promyelocyte
regulatory T cell
CD8-positive, alpha-beta T cell
CD4-positive, alpha-beta memory T cell
effector memory CD8-positive, alpha-beta T cell
naive thymus-derived CD8-positive, alpha-beta T cell
effector memory CD4-positive, alpha-beta T cell, terminally differentiated
naive thymus-derived CD4-positive, alpha-beta T cell
CD8-positive, alpha-beta memory T cell
effector memory CD8-positive, alpha-beta T cell, terminally differentiated
T cell
eosinophil
mast cell
obsolete female germ line stem cell (sensu Vertebrata)
granulocyte
neutrophil
myeloid lineage restricted progenitor cell
myelocyte
CD14-positive monocyte
CD14-low, CD16-positive monocyte
monocyte
promonocyte
CD16-negative, CD56-bright natural killer cell, human
mature natural killer cell
mature NK T cell
ccf_part_of
rdfs:subClassOf
ccf_located_in
develops from
part of
has part