

Kernel Methods in Machine Learning - Course Notes

Hugo Cisneros

1 Kernels and RKHS

1.1 Positive Definite Kernels

Definition 1

A kernel K is a comparison function $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$.

With n data point $\{x_1, x_2, \dots, x_n\}$ a $n \times n$ matrix \mathbf{K} can be defined by $\mathbf{K}_{ij} = K(x_i, x_j)$.

A kernel K is **positive definite** (p.d.) if it is **symmetric** ($K(x, x') = K(x', x)$) and for all sets of a and x

$$\sum_i \sum_j a_i a_j K(x_i, x_j) \geq 0$$

This is equivalent to the kernel matrix being **positive semi-definite**.

Examples:

- Kernel on $\mathbb{R} \times \mathbb{R}$ defined by $K(x, x') = xx'$ is p.d. ($xx' = x'x$ and $\sum_i \sum_j a_i a_j K(x_i, x_j) = (\sum_i a_i x_i)^2 \geq 0$).
- Linear kernel ($K(x, x') = \langle x, x' \rangle_{\mathbb{R}^d}$) is p.d.
- More generally for any set \mathcal{X} , and function $\Phi : \mathcal{X} \rightarrow \mathbb{R}^d$, the kernel defined by $K(x, x') = \langle \Phi(x), \Phi(x') \rangle_{\mathbb{R}^d}$ is p.d.

Theorem 1: Aronszajn, 1950

K is a p.d. kernel on the set \mathcal{X} if and only if there exists a **Hilbert space** \mathcal{H} and a **mapping** $\Phi : \mathcal{X} \rightarrow \mathcal{H}$ such that, for any x, x' in \mathcal{X} :

$$K(x, x') = \langle \Phi(x), \Phi(x') \rangle_{\mathcal{H}}$$

(A Hilbert space is a vector space with an inner product and complete for the corresponding norm).

1.2 Reproducing Kernel Hilbert Spaces (RKHS)

Let \mathcal{X} be a set and $\mathcal{H} \subset \mathbb{R}^{\mathcal{X}}$ a class of functions forming a Hilbert space.

Definition 2

A kernel K is called a **reproducing kernel** (r.k.) of \mathcal{H} if

- \mathcal{H} contains all functions of the form

$$\forall x \in \mathcal{X}, K_x : t \rightarrow K(x, t)$$

- For every $x \in \mathcal{X}$ and $f \in \mathcal{H}$, $f(x) = \langle f, K_x \rangle_{\mathcal{H}}$

If there exists a r.k., \mathcal{H} is called a RKHS.

Definition 3: Equivalent Definition of RKHS

\mathcal{H} is a RKHS if and only if for any $x \in \mathcal{X}$, the mapping

$$\begin{aligned} F : \mathcal{H} &\rightarrow \mathbb{R} \\ f &\mapsto f(x) \end{aligned}$$

is **continuous**.

1.3 Examples

2 Kernel tricks

3 Kernel Methods: Supervised Learning

4 Kernel Methods: Unsupervised Learning

5 The Kernel Jungle

6 Open Problems and Research Topics