

Categories, Proofs, and Processes: Assignment 3

Wira Azmoon Ahmad

11 Feb 2024

Writing this here because I keep mixing them up.

A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is

- **faithful** if each map $F_{A,B} : \mathcal{C}(A, B) \rightarrow \mathcal{D}(F A, F B)$ is injective;
- **full** if each map $F_{A,B} : \mathcal{C}(A, B) \rightarrow \mathcal{D}(F A, F B)$ is surjective

Question 1

For each function $f : C \rightarrow D$ in **Set**, S maps it to the functor

$$S f : \mathcal{C} \rightarrow \mathcal{D}$$

where $\mathcal{C}, \mathcal{D}$ are discrete categories in **Cat** with $|C|, |D|$ objects respectively.

$$\mathbf{Set} \xrightarrow{S} \mathbf{Cat}$$

$$\begin{array}{ccc} C & & \mathcal{C} \\ f \downarrow & \longmapsto & \downarrow S f \\ D & & \mathcal{D} \end{array}$$

Then $\forall a \in C, b \in D, (S a) \in \text{Ob}(\mathcal{C}), (S b) \in \text{Ob}(\mathcal{D})$, the mapping is

$$S f :: S a \mapsto S b \iff f :: a \mapsto b$$

$$S f :: 1_{(S a)} \mapsto 1_{(S b)} \iff f :: a \mapsto b$$

Proposition 1.1. $\forall f \in \text{Ar}(\mathbf{Set}), (S f) : \mathcal{C} \rightarrow \mathcal{D}$ satisfies functoriality.

Proof. Since there is a valid function where $\forall a \in C, f(a) = b$, $(S f)$ will map every object $(S a)$ in $\mathcal{C}$ to an $(S b)$ in $\mathcal{D}$.

There are no arrows between objects in the discrete categories, only identity arrows. For any identity arrow 1_A in $\mathcal{C}$,

$$(S f)(1_A \circ 1_A) = (S f)(1_A) = 1_{(S f)(1_A)}$$

□

Proposition 1.2. *The functor $S : \mathbf{Set} \rightarrow \mathbf{Cat}$ is full and faithful.*

Proof. To show faithfulness, let $f, g : C \rightarrow D$ be arbitrary functions in $\mathbf{Set}$ so $S f, S g : \mathcal{C} \rightarrow \mathcal{D}$. Suppose $S f = S g$. Then since for all $a \in C$,

$$\begin{aligned} S f :: S a \mapsto S b &\iff f :: a \mapsto b \\ S g :: S a \mapsto S b &\iff g :: a \mapsto b \end{aligned}$$

we have

$$\forall a \in C : f :: a \mapsto b \iff S f = S g :: S a \mapsto S b \iff g :: a \mapsto b$$

and $f = g$. S is faithful.

For fullness, let $F : \mathcal{C} \rightarrow \mathcal{D}$ be an arbitrary functor in $\mathbf{Cat}(S A, S B)$, where A, B are sets in $\mathbf{Set}$. Since there exists the function $f : A \rightarrow B$ in $\mathbf{Set}$, then $S f = F$. So S is full. □

Question 2

Definition 2.3. The bivariate Hom-functor is given by

$$\begin{aligned} \mathcal{C}(-, -)(A, B) &:= \mathcal{C}(A, B) \\ \mathcal{C}(-, -)(g : B \rightarrow A, f : C \rightarrow D) &:= h \mapsto f \circ h \circ g, \forall h \in \mathcal{C}(A, C) \end{aligned}$$

for all objects and arrows $A, B, C, D \in \text{Ob}(\mathcal{C})$, $f, g \in \text{Ar}(\mathcal{C})$.

Proposition 2.4. *The bivariate Hom-functor is a functor.*

Proof. Let $A, B, C, D, E, F \in \text{Ob}(\mathcal{C}) = \text{Ob}(\mathcal{C}^{\text{op}})$ be arbitrary objects.

For any object pair A, B , the functor has a mapping to the set $\mathcal{C}(A, B)$.

Let $f : A \rightarrow B, g : B \rightarrow C$ be arbitrary arrows in $\mathcal{C}$, and $h : E \rightarrow D, k : F \rightarrow E$ be arbitrary arrows in $\mathcal{C}^{\text{op}}$.

$$\begin{aligned} \mathcal{C}(-, -)(h, g) \circ \mathcal{C}(-, -)(k, f) &= (p \mapsto g \circ p \circ h) \circ (p' \mapsto f \circ p' \circ k) \\ &= (p \mapsto g \circ (f \circ p \circ k) \circ h) \\ &= (p \mapsto (g \circ f) \circ p \circ (k \circ h)) \\ &= \mathcal{C}(-, -)(h \circ k, g \circ f) \\ &= \mathcal{C}(-, -)((h, g) \circ (k, f)) \end{aligned} \tag{1}$$

So the composition law is satisfied.

For the identity laws,

$$\begin{aligned} \mathcal{C}(-, -)(1_A, 1_B) &= \mathcal{C}(1_A, 1_B) \\ &= h \mapsto 1_B \circ h \circ 1_A \\ &= h \mapsto h \\ &= 1_{\mathcal{C}(A, B)} \\ &= 1_{\mathcal{C}(-, -)(A, B)} \end{aligned} \tag{2}$$

Thus, the bivariate Hom-functor satisfies *functoriality*. □

Question 3

(a)

Fact 3.5. Faithful functors do not, in general, reflect isomorphisms.

Example 3.6.

$$A \xrightarrow{f} B \qquad F A \xrightleftharpoons[g=(F f)^{-1}]{F f} F B$$

The diagram above shows two-object categories $\mathcal{C}, \mathcal{D}$ respectively, with the identities assumed but not drawn. The arrow $g = (F f)^{-1} : F B \rightarrow F A$ in $\mathcal{D}$ has no mapping from an arrow in $\mathcal{C}$.

Functoriality is preserved since

$$\begin{aligned} F(1_A) &= 1_{F A}, & F(1_B) &= 1_{F B} \\ F(f \circ 1_A) &= F(f) = F(f) \circ 1_{F A} = F(F) \circ F(1_A) \\ F(1_B \circ f) &= F(f) = 1_{F B} \circ F(f) = F(1_B) \circ F(F) \end{aligned}$$

The functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is faithful since

$$F f \neq F 1_A = 1_{F A} \neq F 1_B = 1_{F B}$$

It can be noted it is not full because of the arrow g in $\mathcal{D}$.

The arrow $F f$ is isomorphic (to $(F f)^{-1}$) but f is not iso. The functor does not reflect isomorphism.

(b)

Proposition 3.7. *Every full and faithful functor reflects isomorphisms.*

Proof. Let the functor $F : \mathcal{C} \rightarrow \mathcal{D}$ be full and faithful. By fullness, for any arrow g in $\mathcal{D}$, there exists an arrow f in $\mathcal{C}$ such that $F f = g$.

Suppose we have two iso arrows g, g^{-1} in $\mathcal{C}$. Using fullness we find $f : A \rightarrow B, f' : B' \rightarrow A'$ such that $F f = g, F f' = g^{-1}$. Since g, g^{-1} are iso, $F A = F A', F B = F B'$ so

$$g : F A \rightarrow F B, \quad g^{-1} : F B \rightarrow F A$$

where

$$g^{-1} \circ g = 1_{F A}, \quad g \circ g^{-1} = 1_{F B}$$

By functoriality, we have from composition that

$$\begin{aligned} F(f \circ f') &= F f \circ F f' = g \circ g^{-1} = 1_{F B} \\ F(f' \circ f) &= F f' \circ F f = g^{-1} \circ g = 1_{F A} \end{aligned}$$

We also have from the identity law that

$$F(1_A) = 1_{F A}, \quad F(1_B) = 1_{F B}$$

By faithfulness,

$$\begin{aligned} F(1_A) = F(f' \circ f) = 1_{F A} &\implies f' \circ f = 1_A \\ F(1_B) = F(f \circ f') = 1_{F B} &\implies f \circ f' = 1_B \end{aligned}$$

So $f' = f^{-1}$, and f, f^{-1} are iso.

Given an iso arrow $F f$ in $\mathcal{D}$, we then have an iso arrow f in $\mathcal{C}$. The full and faithful functor reflects isomorphisms. □

(c)

Proposition 3.8. *Every equivalence preserves monics.*

Proof. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be an equivalence, and let $f : B \rightarrow C$ be a monic arrow in $\mathcal{C}$. Then $F f : F B \rightarrow F C$. Let $F g, F h : A' \rightarrow F B$ be arbitrary arrows in $\mathcal{D}$.

By equivalence, there exists an object A of $\mathcal{C}$ such that $F(A) \cong A'$.

By fullness, let the arrows $g, h : A \rightarrow B$ be in $\mathcal{C}$.

Suppose $F f \circ F g = F f \circ F h$. Then

$$\begin{aligned} F f \circ F g = F f \circ F h &\implies F(f \circ g) = F(f \circ h) && \text{(by functoriality)} \\ &\implies f \circ g = f \circ h && \text{(by faithfulness)} \\ &\implies g = h && \text{(since } f \text{ is monic)} \\ &\implies F g = F h \end{aligned} \tag{3}$$

Then


$$\forall F g, F h : F f \circ F g = F f \circ F h \implies F g = F h$$

so the equivalence preserves monics. □

Question 4

Proposition 4.9. *If a category $\mathcal{C}$ has a terminal object and pullbacks, it has binary products and equalisers.*

Proof. Let T be the terminal object, and let $\tau_A : A \rightarrow T$ denote the unique arrow from an object A to T .




The morphisms $A \xrightarrow{\tau_A} T \xleftarrow{\tau_B} B$ then have the pullback $A \xleftarrow{p} D \xrightarrow{q} B$.

For any morphisms $A \xleftarrow{p'} D' \xrightarrow{q'} B$, we have by composition,


$$\tau_A \circ p' = \tau_{D'} = \tau_B \circ q' \implies h' : D' \rightarrow D \text{ is unique}$$

Then D with morphisms $A \xleftarrow{p} D \xrightarrow{q} B$ is the product of A and B , since for all pairs $A \xleftarrow{p'} D' \xrightarrow{q'} B$, there is the unique morphism $h' = \langle p', q' \rangle : D' \rightarrow D$ such that $p' = p \circ h'$ and $q' = q \circ h'$.

Using the binary product, we can then show equalisers. Suppose we have the parallel morphisms $A \xrightarrow{f} B$. As shown before, the product $A \times B$ with pullback morphisms $A \xleftarrow{\pi_1} D \xrightarrow{\pi_2} B$ exist. The following diagram with f commutes, and is similar if g is used instead of f , where 1_A is the identity on A .



Let $A \xleftarrow{p} E \xrightarrow{q} A$ be the pullback of $A \xrightarrow{\langle 1_A, f \rangle} A \times B \xleftarrow{\langle 1_A, g \rangle} A$.



Then we show that $e = p = q : E \rightarrow A$ is the equaliser of (f, g) . First, we show $p = q$ using the product,

$$\begin{aligned} p &= \pi_1 \circ \langle 1_A, f \rangle \circ p \\ &= \pi_1 \circ \langle 1_A, g \rangle \circ q \\ &= q \end{aligned} \tag{4}$$

Then we have

$$\begin{aligned}
f \circ e &= \pi_2 \circ \langle 1_A, f \rangle \circ e \\
&= \pi_2 \circ \langle 1_A, g \rangle \circ e \\
&= g \circ e
\end{aligned} \tag{5}$$

Suppose we then have a morphism $h : C \rightarrow A$ such that $f \circ h = g \circ h$. Then

$$\begin{aligned}
\langle 1_A, f \rangle \circ h &= \langle 1_A \circ h, f \circ h \rangle \\
&= \langle 1_A \circ h, g \circ h \rangle \\
&= \langle 1_A, g \rangle \circ h
\end{aligned} \tag{6}$$

Then by the pullback, there is a unique morphism $\hat{h} : C \rightarrow E$ such that $h = e \circ \hat{h}$.

$$\begin{array}{ccccc}
E & \xrightarrow{e} & A & \xrightleftharpoons[g]{f} & B \\
\uparrow \hat{h} & \nearrow h & & & \\
C & & & &
\end{array}$$


Then $e : E \rightarrow A$, the pullback of $(\langle 1_A, f \rangle, \langle 1_A, g \rangle)$, is the equaliser of (f, g) where $A \xrightleftharpoons[g]{f} B$.

Thus, it is shown that the category $\mathcal{C}$ has binary products and equalisers.

□

Question 5

Proposition 5.10. *If $ABCD, BEDF$ are pullback squares, then so is $AECF$.*



Proof. Suppose we have morphisms $C \xleftarrow{p} G \xrightarrow{q} E$ such that $(g \circ f) \circ p = e \circ q$.

By the pullback square $BEDF$,

$$e \circ q = g \circ (f \circ p) \implies \exists! h' : G \rightarrow B, q = b \circ h' \wedge f \circ p = d \circ h'$$

By the pullback square $ABCD$ and using h' ,

$$d \circ h' = f \circ p \implies \exists! h : G \rightarrow A, p = c \circ h \wedge h' = a \circ h$$

From the unique h, h' such that $q = b \circ h', h' = a \circ h$, then h is the unique morphism where $q = b \circ a \circ h$. Then combining $q = b \circ a \circ h$ with $p = c \circ h$ (from pullback square $ABCD$) where h is unique, $AECF$ is also a pullback square. □

Question 6