

Categories, Proofs, and Processes:

Assignment 5

Wira Azmoon Ahmad

26 Feb 2024

Question 1

(a)

The set-indexed product of a set of objects $\{A_i\}_{i \in I}$ is the Cartesian product

$$\prod_{i \in I} A_i = \left\{ g : I \rightarrow \bigcup_{i \in I} A_i \mid \forall i \in I : g(i) \in A_i \right\}$$

with its arrows $\pi_j(g) = g(j)$ for all $j \in I$.


Let C be an object with a set of arrows $f_i : C \rightarrow A_i$. Then the unique arrow $\langle f_i \rangle_{i \in I}$ is defined where for all $c \in C$,

$$\langle f_i \rangle_{i \in I}(c) := i \mapsto f_i(c)$$

For each $j \in I$, and for all $c \in C$,

$$\begin{aligned} \pi_j \circ \langle f_i \rangle_{i \in I}(c) &= \pi_j(i \mapsto f_i(c)) \\ &= (i \mapsto f_i(c))(j) \\ &= f_j(c) \end{aligned} \tag{1}$$

so $\forall j \in I, \pi_j \circ \langle f_i \rangle_{i \in I} = f_j$.



Suppose there is another arrow $h : C \rightarrow \prod_{i \in I} A_i$ such that $\forall j \in I, \pi_j \circ h = f_j$. Then $\forall c \in C$,

$$\begin{aligned} \pi_j \circ h(c) &= \pi_j(h(c)) \\ &= h(c)(j) \\ h(c)(j) &= f_j(c) \\ \implies h(c) &:: j \mapsto f_j(c) \end{aligned} \tag{2}$$

So $h = \langle f_i \rangle_{i \in I}$ and the arrow is unique.

(b)

When $|I| = 2$, we get binary products. The product of 2 objects A_i, A_j for $i, j \in I$ is the object $A_i \times A_j$ with arrows $A_i \xleftarrow{\pi_i} A_i \times A_j \xrightarrow{\pi_j} A_j$ such that for any object C and set of arrows $A_i \xleftarrow{\pi_i} C \xrightarrow{\pi_j} A_j$, there is a unique arrow $\langle f_i, f_j \rangle : C \rightarrow A_i \times A_j$ such that $\pi_i \circ \langle f_i, f_j \rangle = f_i$ and $\pi_j \circ \langle f_i, f_j \rangle = f_j$.

When $|I| = 1$, we get terminal objects. The product of 1 object A_i for $i \in I$ is the terminal object A_i with the identity 1_{A_i} such that for any object C , the arrow $f_i : C \rightarrow A_i$, is the unique arrow from C to A_i .

(c)

Let (L, γ) be the limit of $D : \mathcal{I} \rightarrow \mathcal{C}$. Then the arrows are defined as

$$\{\gamma_J : L \rightarrow D J\}_{J \in \text{Ob}(\mathcal{I})}$$

where the only arrows in $\mathcal{J}$ are identities 1_J so $D 1_J = 1_{D J}$. Since (L, γ) is the terminal object in $\mathbf{Cone}(D)$, for every other cone to D , (A, δ) , there is a unique arrow $g : A \rightarrow L$ such that $\forall J \in \text{Ob}(\mathcal{I}), \gamma_J \circ g = \delta_J$.

$$\begin{array}{ccc} & D J & \\ \delta_J \nearrow & \uparrow \gamma_J & \\ A & \xrightarrow{g} & L \end{array}$$

By using the functor $S : \mathbf{Set} \rightarrow \mathbf{Cat}$ from Assignment 3 Exercise 1, let $S I = \mathcal{I}$, so there's an isomorphism $\phi : I \cong \text{Ob}(\mathcal{I})$. Then set $\forall j \in I$,

- $D \phi(j) := A_j$
- $L := \prod_{j \in I} A_j$
- $\gamma_{\phi(j)} := \pi_j$
- $A := C$
- $\delta_{\phi(j)} := f_j$
- $g := \langle f_j \rangle_{j \in I}$

so the limit of the diagram matches the definition of the set-indexed products.

(d)

Proposition 1.1. *If $\mathcal{D}$ has all set-indexed products, then so too does the functor category $[\mathcal{C}, \mathcal{D}]$.*

Proof. Suppose $\mathcal{D}$ has all set-indexed products.

Let I be the set of indices, and $\{F_i : \mathcal{C} \rightarrow \mathcal{D}\}_{i \in I}$ be a set of functors. First, we construct the product functor $\prod_{i \in I} F_i : \mathcal{C} \rightarrow \mathcal{D}$ and prove its functoriality.

$$\begin{aligned} \prod_{i \in I} F_i(C) &:= \prod_{i \in I} (F_i C) \\ \prod_{i \in I} F_i(f : C \rightarrow C') &:= \prod_{i \in I} (f_i : F_i C \rightarrow F_i C') := \langle f_i \circ \pi_i \rangle_{i \in I} \end{aligned}$$

where $\pi_i : \prod_{i \in I} F_i \rightarrow F_i C$, which exists in $\mathcal{D}$.

Let $C \xrightarrow{f} C' \xrightarrow{g} C''$ be arrows.

$$\begin{aligned} \prod_{i \in I} F_i(g \circ f) &= \langle (g \circ f)_i \circ \pi_i \rangle_{i \in I} \\ &= \prod_{i \in I} (g \circ f)_i \circ \langle \pi_i \rangle_{i \in I} \\ &= \prod_{i \in I} g_i \circ \prod_{i \in I} f_i \circ \langle \pi_i \rangle_{i \in I} \\ &= \langle g_i \circ \pi_i \rangle_{i \in I} \circ \prod_{i \in I} f_i \circ \langle \pi_i \rangle_{i \in I} \\ &= \langle g_i \circ \pi_i \rangle_{i \in I} \circ \langle f_i \circ \pi_i \rangle_{i \in I} \\ &= \prod_{i \in I} F_i(g) \circ \prod_{i \in I} F_i(f) \end{aligned} \tag{3}$$

$$\begin{aligned} \prod_{i \in I} F_i(1_C) &= \langle 1_{F_i C} \circ \pi_i \rangle_{i \in I} \\ &= \langle \pi_i \rangle_{i \in I} \\ &= \prod_{i \in I} 1_{F_i C} \\ &= 1_{\prod_{i \in I} (F_i C)} \\ &= 1_{\prod_{i \in I} F_i(C)} \end{aligned} \tag{4}$$

So the product functor satisfies functoriality.

Next, we define and show the naturality of the functor projections. For an arbitrary natural transformation $t_j : \prod_i F_i \rightarrow F_j$, define $\forall C$ in $\mathcal{C}$,

$$t_{j,C} = \pi_{j,C} : \prod_i F_i C \rightarrow F_j C$$

which exists as a projection in $\mathcal{D}$. Let $f : C \rightarrow C'$ be an arrow in $\mathcal{C}$, and $f_j : F_j C \rightarrow F_j C'$ be an arrow in $\mathcal{D}$. Then by the property of the product, we have

$$f_j \circ \pi_{j,C} = \pi_{j,C'} \circ \langle f_i \circ \pi_i \rangle_i$$

So naturality is satisfied.

$$\begin{array}{ccc}
& G & \\
s_j \swarrow & \downarrow \langle s_i \rangle_i & \searrow \\
& \prod_i F_i & \\
& \downarrow t & \\
& F_j &
\end{array}
\quad
\begin{array}{ccccc}
G & C & \xrightarrow{G f} & G & C' \\
\downarrow \langle s_i \rangle_i & & & & \downarrow \langle s_i \rangle_i \\
s_{j,C} \swarrow & \prod_i F_i & C & \xrightarrow{\langle f_i \circ \pi_i \rangle_i} & \prod_i F_i & C' & \searrow s_{j,C'} \\
& \downarrow \pi_{j,C} & & & \downarrow \pi_{j,C'} & \\
& F_j & C & \xrightarrow{f_j} & F_j & C'
\end{array}$$

Let again $f : C \rightarrow C'$ be an arrow in $\mathcal{C}$, Suppose we have some functor G and a set of natural transformations $s_i : G \rightarrow F_i$. Define the natural transformation $\langle s_i \rangle_i : G \rightarrow \prod_i F_i$ such that

$$\langle s_i \rangle_{i,C} = \langle s_i, C \rangle_i : G C \rightarrow \prod_i F_i C$$

By naturality of s_i , we have

$$f_i \circ s_{i,C} = s_{i,C'} \circ G f$$

From the property of the product, we have that

$$\pi_{j,C} \circ \langle s_i \rangle_{i,C} = s_{j,C}, \quad \pi_{j,C'} \circ \langle s_i \rangle_{i,C'} = s_{j,C'}$$

Then these imply that

$$\begin{aligned}
f_j \circ \pi_{j,C} \circ \langle s_i \rangle_{i,C} &= \pi'_{j,C'} \circ \langle s_i \rangle_{i,C'} \circ G f \\
\pi_{j,C'} \circ \langle f_i \circ \pi_i \rangle_i \circ \langle s_i \rangle_{i,C} &= \pi_{j,C'} \circ \langle s_i \rangle_{i,C'} \circ G f
\end{aligned}$$

Since this is true $\forall j \in I$, then by the property of the product, there is the unique arrow

$$\langle f_i \circ \pi_i \rangle_i \circ \langle s_i \rangle_{i,C} = \langle s_i \rangle_{i,C'} \circ G f : G C \rightarrow \prod_i F_i C'$$

So the naturality of $\langle s_i \rangle_i$ is satisfied. Since $\forall C$ in $\mathcal{C}$, and $\forall j \in I$, we have some $\pi_{j,C}$ that

$$\pi_{j,C} \circ \langle s_i \rangle_{i,C} = s_{j,C}$$

where $t_j = \pi_{j,C}$, then

$$\forall j \in I : t_j \circ \langle s_i \rangle_i = s_j$$

Now, we show the uniqueness of the natural transformation. Suppose there is some other natural transformation $u : G \rightarrow \prod_i F_i$ such that $\forall j \in I, t_j \circ u = s_j$. Then for all C in $\mathcal{C}$, $\forall j \in I, \pi_{j,C} \circ u_C = s_{j,C}$ for some $\pi_{j,C}$. Then there is a set of mappings

$$\pi_j \circ u_C = s_{j,C} : G C \rightarrow F_j C$$

By property of the product, there is a unique mapping

$$\langle \pi_i \circ u_C \rangle_i = \langle s_i, C \rangle_i : G C \rightarrow \prod_i F_i C$$

such that $\forall j \in I, \pi_j \circ \langle \pi_i \circ u_C \rangle_i = s_{j,C}$. Then for all C in $\mathcal{C}$,

$$\begin{aligned}
s_{j,C} &= \pi_j \circ \langle \pi_i \circ u_C \rangle_i \\
&= \pi_j \circ \langle \pi_i \rangle_i \circ u_C \\
&= \pi_j \circ 1_{\prod_i F_i C} \circ u_C \\
&= \pi_j \circ u_C
\end{aligned} \tag{5}$$

By property of the product, there is the unique arrow

$$\langle s_{i,C} \rangle_i = \langle \pi_i \circ u_C \rangle_i = u_C : G C \rightarrow \prod_i F_i C$$

So $\forall C$ in $\mathcal{C}, u_C = \langle s_{i,C} \rangle_i$. Thus, $\langle s_i \rangle_i : G \rightarrow \prod_i F_i$ is the unique natural transformation that satisfies the properties defined above. $\square$

(e)

Proposition 1.2. *The Yoneda embedding $Y : \mathcal{C} \rightarrow [\mathcal{C}^{op}, \mathbf{Set}]$ preserves set-indexed products, when they exist.*

Proof. Suppose $\mathcal{C}$ has set-indexed products. Let $\{C_i\}_i$ be a set of objects in $\mathcal{C}$ with the product, object $\prod_i C_i$, with arrows $\pi_j : \prod_i C_i \rightarrow C_j$. We show that for the set of objects $\{Y C_i\}_i := \mathcal{C}(-, C_i)_i$ in $[\mathcal{C}^{op}, \mathbf{Set}]$, the object with arrows

$$\begin{aligned}
Y \prod_i C_i &:= \mathcal{C}(-, \prod_i C_i) \\
\forall j \in I, Y \pi_j &:= \mathcal{C}(-, \pi_j) : \mathcal{C}(-, \prod_i C_i) \rightarrow \mathcal{C}(-, C_j)
\end{aligned}$$

is their set-indexed product.

By theorem from lectures, Y is full and faithful. Hence, there is an bijection between sets of arrows

$$\phi_{A,B} : \mathcal{C}(A, B) \cong [\mathcal{C}^{op}, \mathbf{Set}](Y A, Y B)$$

Then by the property of the product, since for any object C' and set of arrows $f_i : C' \rightarrow C_i$ in $\mathcal{C}$, there is the unique arrow $\langle f_i \rangle_i : C' \rightarrow \prod_i C_i$ such that $\forall j \in I, \pi_j \circ \langle f_i \rangle_i = f_j$, for the corresponding objects $Y C', \{Y C_i\}_i, Y \prod_i C_i$, with the set of arrows $\{Y f_i\}_i = \{\phi_{C', C_i}(f_i)\}_i$ in $[\mathcal{C}^{op}, \mathbf{Set}]$, there must be the unique arrow

$$Y \langle f_i \rangle_i = (\phi_{C', \prod_i C_i})(\langle f_i \rangle_i) : Y C' \rightarrow Y \prod_i C_i$$

such that $\forall j \in I, (Y \pi_j) \circ (Y \langle f_i \rangle_i) = (Y f_j)$. The embedding preserves set-indexed products when they exist. $\square$

Question 2

Note that for all objects A, B in $\mathcal{P}$,

$$\mathcal{P}(A, B) = \begin{cases} \emptyset & \text{if } A \not\leq B \\ \{\leq\} & \text{if } A \leq B \end{cases}$$

Then for each object $\mathcal{P}(-, A)$ in $[\mathcal{P}, \mathbf{Set}]$, $\forall A, B, C, D \in P$,

$$\mathcal{P}(-, A)(B) \in \{\emptyset, \{\leq\}\}$$

$$\mathcal{P}(-, A)(f : B \rightarrow C) \in \{\emptyset \rightarrow \emptyset, \emptyset \rightarrow \{B \leq A\}, \{C \leq A\} \rightarrow \{B \leq A\}\}$$

And for all arrows $\mathcal{P}(-, f : A \rightarrow B)$,

$$\mathcal{P}(-, f)(g) \in \{\emptyset \rightarrow \emptyset, \emptyset \rightarrow \{C \leq B\}, \{D \leq A\} \rightarrow \{C \leq B\}\}$$

$$\mathcal{P}(-, f)(C) \in \{\emptyset \rightarrow \emptyset, \emptyset \rightarrow \{C \leq B\}, \{C \leq A\} \rightarrow \{C \leq B\}\}$$

These only consist of either the empty set $\emptyset$, or singleton sets $\{*\}$ and arrows between them. Then the objects in the subcategory of $\mathbf{Set}$ mapped to are just the empty set $\emptyset$ and the singleton set $\{*\}$ with the arrows $1_\emptyset, 1_{\{*\}}, \emptyset \rightarrow \{*\}$, corresponding to the full subcategory $\mathcal{B}$.

The functor is full since for any arrow $\mathcal{P}(-, f : A \rightarrow B)$, there is an arrow $f : A \rightarrow B$ in $\mathcal{P}$. It is full since if two arrows $F, G : \mathcal{P}(-, A) \rightarrow \mathcal{P}(-, B)$ are such that $F = G$, so $F = \mathcal{P}(-, f : A \rightarrow B)$, $G = \mathcal{P}(-, g : A \rightarrow B)$, then $f = g$ since P is a preorder.

(a)

(b)

(c)

(d)

(e)

(f)