

MODULE 18 / 40

Red-winged Blackbird

*Agelaius phoeniceus***QUICK STATS**

Length	6.7 – 9.1 in (17 – 23 cm)
Wingspan	12.2 – 15.8 in (31 – 40 cm)
Weight	1.1 – 2.7 oz (32 – 77 g)
Lifespan	Up to 15 years (avg. 2 in wild)

PRESENTATION

The Red-winged Blackbird is one of the most abundant birds in North America — and among the most instantly recognizable. The male's jet-black plumage set against blazing scarlet-and-yellow shoulder patches is one of the most vivid color combinations in North American ornithology. Few sights signal the arrival of spring more dramatically than a male Red-winged Blackbird perched atop a cattail, wings half-spread to display those brilliant epaulets, delivering his liquid, bubbling "conk-la-reeee" call. In contrast, females are so heavily streaked and brown that beginners often fail to recognize them as the same species.

HABITAT & RANGE

The Red-winged Blackbird breeds across virtually all of North America, from Alaska and Canada south through the United States, Mexico, and into Central America. It is one of the most widespread breeding birds on the continent. It shows a strong preference for wetland habitats — cattail marshes, reed beds, wet meadows, and the edges of ponds and lakes — but also nests in dry upland fields and roadsides. In winter, northern populations migrate south and form enormous mixed flocks with other blackbirds and starlings, sometimes numbering in the millions.

DIET & FEEDING

- Black-oil sunflower seeds and cracked corn at feeders
- White millet scattered on the ground
- Wild grass seeds, grain, and weed seeds — the core of the natural diet
- Insects, spiders, and aquatic invertebrates during breeding season
- Wild berries and small fruits in autumn
- Waste grain gleaned from agricultural fields in winter

BEHAVIOR & SOCIAL LIFE

Red-winged Blackbirds are highly polygynous — a single male may hold a territory with up to 15 females nesting within it, though 2–3 is more typical. Males arrive on the breeding grounds weeks before females and spend much of the early season in fierce territorial battles, displaying their red shoulder patches and singing constantly from prominent perches. They are remarkably aggressive defenders of their territories, regularly dive-bombing much larger animals — hawks, herons, crows, and even humans — that venture too close to active nests. Outside the breeding season they are highly gregarious, forming enormous mixed-species flocks that roost communally.

NESTING & BREEDING

Females build deep, woven cup nests of grasses and aquatic vegetation, lashed to cattail stems or low shrubs over or near water. Clutches of 3–4 pale blue-green eggs with dark scrawls are incubated by the female alone for 11–13 days. Both parents feed the nestlings, which fledge at 11–14 days. Pairs raise 2–3 broods per season. Females accept cowbird eggs at a high rate, making them frequent brood parasite hosts.

SONG & COMMUNICATION

The Red-winged Blackbird's song is one of the most evocative sounds of North American wetlands — a liquid, gurgling "conk-la-reeee" or "oak-a-lee" delivered with a dramatic wing-spread display. Males have individual song repertoires used to advertise territory. The sharp "check" call and harsh chattering alarm notes are given by both sexes. Females produce a scolding chatter when disturbed near the nest.

CONSERVATION STATUS

IUCN Red List:	Least Concern	Population trend:	Declining
----------------	---------------	-------------------	-----------

Despite being one of the most abundant birds in North America — with a population estimated at 150 million individuals — the Red-winged Blackbird has declined significantly since 1970, losing an estimated 30% of its population. Wetland drainage, agricultural intensification, and the loss of native grasslands are the primary drivers. Protecting and restoring wetland habitats is critical for this species.

FEEDER TIPS

- Scatter cracked corn and white millet on a large ground platform
- Red-winged Blackbirds arrive in spring flocks — expect multiple birds at once
- They prefer open feeding areas with water or marsh nearby
- In winter, mixed blackbird flocks may descend on feeders by the hundreds
- Planting native cattails and marsh plants near water attracts nesting pairs

COLORING GUIDE

ZONE	RECOMMENDED COLOR
Body (male)	Glossy jet black
Shoulder patch (male)	Brilliant scarlet-red
Shoulder border	Yellow to pale yellow
Female — overall	Dark brown, heavily streaked
Female — face	Pale buff supercilium stripe
Bill	Sharp, pointed, black

COLOR ME!

Red-winged Blackbird • *Agelaius phoeniceus* • Module 18

