

MODULE 26 / 40

Black-capped Chickadee

Poecile atricapillus

QUICK STATS

Length	4.7 – 5.9 in (12 – 15 cm)
Wingspan	6.3 – 8.3 in (16 – 21 cm)
Weight	0.3 – 0.5 oz (9 – 14 g)
Lifespan	Up to 12 years (avg. 2–3 in wild)

PRESENTATION

The Black-capped Chickadee is arguably the most beloved feeder bird in North America — a tiny, round, endlessly cheerful bird whose bold black cap, crisp white cheeks, and fearless curiosity have made it a symbol of winter resilience and backyard birding joy. It is one of the few wild birds that can be trained to take seeds from an outstretched hand, and it often does so with remarkable boldness. Despite its tiny size — barely heavier than two nickels — it is superbly adapted to survive the harshest northern winters, capable of lowering its body temperature at night to conserve energy and maintaining thousands of cached food items whose locations it can remember for weeks.

HABITAT & RANGE

The Black-capped Chickadee is found year-round across the northern United States and most of Canada, from Alaska east to Newfoundland and south through the northern tier of states, the Appalachians, and the Pacific Northwest. It inhabits deciduous and mixed forests, forest edges, suburban parks, and wooded gardens wherever there are trees large enough to provide nesting cavities and foraging opportunities. It is a permanent resident throughout its range, though it may make short irruptive movements in years when northern food supplies fail.

DIET & FEEDING

- Black-oil sunflower seeds — the absolute top choice at feeders
- Sunflower chips and safflower seeds
- Suet cakes — especially important in cold winter months
- Peanut pieces and whole peanuts
- Nyjer seed — taken occasionally
- Insects, spiders, and their eggs gleaned from bark — dominant summer food

BEHAVIOR & SOCIAL LIFE

Black-capped Chickadees live in stable winter flocks of 6–10 birds with a fixed dominance hierarchy. The dominant pair leads the flock and gets priority access to the best food. Subordinate birds are pushed to the flock's periphery, where predator risk is higher. Flock members communicate constantly with a complex vocabulary of calls. They are among the most important "nuclear species" in winter mixed-species flocks — nuthatches, Downy Woodpeckers, and kinglets follow chickadee flocks and respond to their alarm calls. Chickadees are extraordinary food cachers, hiding thousands of seeds in bark crevices, leaves, and soil, and relocating each cache within weeks.

NESTING & BREEDING

Black-capped Chickadees excavate their own nest cavities in soft, rotting wood — a capability unique among small songbirds. They also readily use nest boxes. The female builds a soft cup nest inside the cavity using moss, plant fibers, and a thick lining of mammal fur and plant down. Clutches of 6–8 white eggs with reddish-brown spots are incubated by the female for 12–13 days. Both parents feed the nestlings, which fledge at 16 days. One brood per season.

SONG & COMMUNICATION

The Black-capped Chickadee has one of the most complex vocal systems of any North American bird. Its familiar "chick-a-dee-dee-dee" call varies in the number of "dee" notes to encode the urgency of predator threats — more "dees" mean a more dangerous predator nearby. The male's clear, two-note whistled "fee-bee" song is one of the first sounds of late winter, often heard on warm days in January or February.

CONSERVATION STATUS

IUCN Red List:	Least Concern	Population trend:	Stable
----------------	---------------	-------------------	--------

The Black-capped Chickadee population is estimated at 41 million individuals and is broadly stable. It is one of the most studied birds in North America, with long-term research revealing remarkable cognitive abilities including episodic-like memory and the ability to expand the hippocampus (the brain's memory center) in autumn to accommodate increased caching demands. It benefits strongly from backyard feeders and nest box programs.

FEEDER TIPS

- Black-oil sunflower seeds in any feeder style — chickadees use all types
- They cache seeds — expect rapid, repeated visits as they store for winter
- Hand-feeding is achievable: stand still near the feeder with seeds in palm
- Install a nest box (1.25-inch entrance) on a tree or post — they will use it
- Suet in a cage feeder is especially valued during cold snaps below freezing

COLORING GUIDE

ZONE	RECOMMENDED COLOR
Cap	Glossy black
Bib	Black — extends to upper breast
Cheeks	Bright white
Back & wings	Gray — soft, medium tone
Breast & belly	White to pale buff on flanks
Bill	Short, stout, black

COLOR ME!

Black-capped Chickadee • *Poecile atricapillus* • Module 26

