

Science | Volume 393 | Issue 6818

Issue date: 2026-09-24 | Retrieved: 2026-09-24T21:35:20.050865+00:00 | 13 relevant primary-research papers

Evidence discipline

Paper-supported findings, published enabling evidence, cross-species inference, speculative hypotheses, and proposed pilot experiments are labeled separately. Access notes state when only an abstract or limited preview was available.

Screening notes

- Official publisher issue page returned HTTP 403; no full-page review is claimed.
- Screened 13 biology-relevant journal articles after excluding reviews, perspectives, news, corrections, and editorials by metadata/title rules.
- All summaries are deterministic extractions from the bounded public evidence packet; no language-model API or paid research service was used.

Official issue

<https://www.science.org/toc/science/393/6818>

Why Xenopus now

Early Xenopus combines rapid external development, targeted microinjection, explants, lineage tracing, live imaging, F0 CRISPR, rescue, and scalable replication. It can convert abstract-supported candidate mechanisms into spatially resolved causal tests before committing to slower mammalian models.

Escape of the *BC200* gene to a human poxvirus reveals its persistent transposition in primates

Pu Gao, Ellen J. Pritham, Cedric Feschotte, Cheng Sun

journal-article | Science 393(6818) | 2026-09-24

DOI [10.1126/science.aeh3441](https://doi.org/10.1126/science.aeh3441)

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Although BC200 was co-opted approximately 40 million years ago to regulate neuronal translation, we show that it never lost its mobilization capacity.
WHY IT MATTERS	The study advances mechanistic understanding of Escape of the <i>BC200</i> gene to a human poxvirus reveals its persistent transposition in primates and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• We identified two insertions of the human BC200 noncoding RNA gene in the poxvirus molluscum contagiosum virus (MCV), which were likely acquired through long interspersed nuclear element 1 (LINE-1 or L1)-mediated retrotransposition during modern human history. • Although BC200 was co-opted approximately 40 million years ago to regulate neuronal translation, we show that it never lost its mobilization capacity. • Rather, BC200 functioned as a “master” source of L1-mediated germline retrotransposition events throughout anthropoid evolution, spawning hundreds of lineage-specific insertions.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed <i>Xenopus</i> mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Escape of the BC200 gene to a human poxvirus reveals its persistent...

IMPORTANCE / SIGNIFICANCE

Immunity And Host Response is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-18 to stabilize cell-stress signaling, epithelial integrity, and lineage-specific transcription; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target epidermal and mesendodermal lineages before mature immunity by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: cell-stress signaling, epithelial integrity, and lineage-specific transcription. Secondary readouts: survival, barrier morphology, and developmental timing, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo

FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine *Xenopus* capabilities

RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.aeh3441](https://doi.org/10.1126/science.aeh3441)

Mammals produce *cyclo* -octasulfur to suppress lipid peroxidation and ferroptosis

Uladzimir Barayeu, Seiryu Ogata, Tsuyoshi Takata, Minkyung Jung, Tetsuro Matsunaga, Masanobu Morita, Tomoaki Ida, Mike Lange, et al.

journal-article | Science 393(6818) | 2026-09-24

DOI 10.1126/science.aec5473

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	We show that mammals harbor elemental sulfur in the form of its most stable allotrope, cyclo -octasulfur (S 8).
WHY IT MATTERS	The study advances mechanistic understanding of Mammals produce <i>cyclo</i> -octasulfur to suppress lipid peroxidation and ferroptosis and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• We show that mammals harbor elemental sulfur in the form of its most stable allotrope, cyclo -octasulfur (S 8). • We found that S 8 accumulates to millimolar concentrations in mitochondrial membranes and in lipid droplets in both mouse and human cells. • We further identified lipid droplet-associated nitric oxide synthase as a source of S 8 biosynthesis and found that S 8 accumulation in lipid droplets limits lipid peroxidation and suppresses ferroptosis.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Mammals produce *cyclo*-octasulfur to suppress lipid peroxidation and...

IMPORTANCE / SIGNIFICANCE

Metabolism And Organelle Function is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-13 to stabilize organelle state, metabolite-sensitive reporters, and lineage-marker induction; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target animal-cap and marginal-zone explants by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-13. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: organelle state, metabolite-sensitive reporters, and lineage-marker induction. Secondary readouts: survival, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo

FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine *Xenopus* capabilities

RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.aec5473](#)

Single wildfire smoke exposure causes lasting harm, natural experiment in monkeys hints

Sara Reardon

journal-article | Science 393(6818) | 2026-09-24

DOI [10.1126/science.aem6545](https://doi.org/10.1126/science.aem6545)

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Infant macaques showed immune changes and signs of faster aging years after inhaling smoke
WHY IT MATTERS	The study advances mechanistic understanding of Single wildfire smoke exposure causes lasting harm, natural experiment in monkeys hints and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• Infant macaques showed immune changes and signs of faster aging years after inhaling smoke
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Single wildfire smoke exposure causes lasting harm, natural experiment in...

IMPORTANCE / SIGNIFICANCE

Immunity And Host Response is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-18 to stabilize cell-stress signaling, epithelial integrity, and lineage-specific transcription; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target epidermal and mesendodermal lineages before mature immunity by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: cell-stress signaling, epithelial integrity, and lineage-specific transcription. Secondary readouts: survival, barrier morphology, and developmental timing, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.aem6545](https://doi.org/10.1126/science.aem6545)

Fibroblasts restrain gut inflammation by IGF1-dependent regulation of innate lymphocytes

Qingxia Lin, Ruichao E. Liu, Qiang Wang, Lan Kang, Boyuan Chen, Yu Lu, Yanting Zhang, Zhijie Gu, et al.

journal-article | Science 393(6818) | 2026-09-24

DOI 10.1126/science.aei0062

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Using mouse models of intestinal inflammation, we found cross-talk between IGF1-expressing fibroblasts and group 3 innate lymphoid cells (ILC3s).
WHY IT MATTERS	The study advances mechanistic understanding of Fibroblasts restrain gut inflammation by IGF1-dependent regulation of innate lymphocytes and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• Using mouse models of intestinal inflammation, we found cross-talk between IGF1-expressing fibroblasts and group 3 innate lymphoid cells (ILC3s).
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Fibroblasts restrain gut inflammation by IGF1-dependent regulation of innate...

IMPORTANCE / SIGNIFICANCE

Immunity And Host Response is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-18 to stabilize cell-stress signaling, epithelial integrity, and lineage-specific transcription; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target epidermal and mesendodermal lineages before mature immunity by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: cell-stress signaling, epithelial integrity, and lineage-specific transcription. Secondary readouts: survival, barrier morphology, and developmental timing, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.aei0062](https://doi.org/10.1126/science.aei0062)

Erratum for the Research Article “Modeling of late-onset Alzheimer’s disease neuropathology via direct neuronal reprogramming”

Author list unavailable in retrieved metadata

journal-article | Science 393(6818) | 2026-09-24

DOI [10.1126/science.aem2402](https://doi.org/10.1126/science.aem2402)

Access: Metadata and title only; no abstract or full text was retrieved. | Evidence status: metadata/title-supported

MAIN DISCOVERY	The title and publisher metadata identify a primary-research study of Erratum for the Research Article “Modeling of late-onset Alzheimer’s disease neuropathology via direct neuronal reprogramming”; no abstract was retrieved, so no more specific discovery claim is made.
WHY IT MATTERS	The study advances mechanistic understanding of Erratum for the Research Article “Modeling of late-onset Alzheimer’s disease neuropathology via direct neuronal reprogramming” and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Computational modeling
KEY EVIDENCE	• The retrieved publisher-deposited metadata identifies this item as a journal article in the detected volume and issue. • The DOI provides the canonical path for manual verification of figures and full methods.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Erratum for the Research Article “Modeling of late-onset Alzheimer’s disease...”

IMPORTANCE / SIGNIFICANCE

Neural Development And Function is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 10-18 to stabilize neural-domain geometry, lineage allocation, and reporter dynamics; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target neural plate and neural-border lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 10-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: neural-domain geometry, lineage allocation, and reporter dynamics. Secondary readouts: marker expression, cell trajectories, and neural-tube morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.aem2402](https://doi.org/10.1126/science.aem2402)

Ceramide synthesis mediates colorectal cancer metastasis through a YAP-driven regenerative program

Swagata Goswami, Qiming Zhang, Abdullah Burak Yildiz, Upasana Das Adhikari, Akhouri Kishore Raghawan, Manon Bulliard, Sabhyata Sedhain, Odai Darawshi, et al.

journal-article | Science 393(6818) | 2026-09-24

DOI [10.1126/science.adw8520](https://doi.org/10.1126/science.adw8520)

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	We demonstrate that tumor-intrinsic ceramide metabolism, amplified by dietary fat, initiates colorectal cancer metastasis.
WHY IT MATTERS	The study advances mechanistic understanding of Ceramide synthesis mediates colorectal cancer metastasis through a YAP-driven regenerative program and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• We demonstrate that tumor-intrinsic ceramide metabolism, amplified by dietary fat, initiates colorectal cancer metastasis. • We observed that dietary fat exposure triggers a sustained increase in de novo ceramide biosynthesis, mediated by the dihydroceramide desaturase Degs1 . • Ceramide accumulation activates yes-associated protein (YAP) through protein phosphatase 2A (PP2A)-mediated dephosphorylation, promoting a durable shift toward a distinct YAP-driven regenerative (YAP-DR) program, marked by Basp1 , that promotes metastasis.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Ceramide synthesis mediates colorectal cancer metastasis through a YAP-driven...

IMPORTANCE / SIGNIFICANCE

Metabolism And Organelle Function is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-13 to stabilize organelle state, metabolite-sensitive reporters, and lineage-marker induction; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target animal-cap and marginal-zone explants by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-13. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: organelle state, metabolite-sensitive reporters, and lineage-marker induction. Secondary readouts: survival, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.adw8520](https://doi.org/10.1126/science.adw8520)

Senescence-directed nanotherapy ameliorates fibrosis and overcomes immune exclusion in cancer

Clemens Hinterleitner, Valentin J. A. Barthet, Hailey V. Goldberg, Kristen C. Vogt, Ana Marie Perea, Logan R. Hillger, Stephen Ruiz, Domhnall McHugh, et al.

journal-article | Science 393(6818) | 2026-09-24

DOI [10.1126/science.aeg4791](https://doi.org/10.1126/science.aeg4791)

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Here, we show that P-selectin is expressed by a subset of senescent-like cells in fibrotic tissues and tumors.
WHY IT MATTERS	The study advances mechanistic understanding of Senescence-directed nanotherapy ameliorates fibrosis and overcomes immune exclusion in cancer and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Computational modeling
KEY EVIDENCE	• Fibrotic remodeling of tissues and tumors establishes immunosuppressive microenvironments that drive organ dysfunction and, in cancer, limit response to immunotherapy. • Here, we show that P-selectin is expressed by a subset of senescent-like cells in fibrotic tissues and tumors. • These findings establish SMNPs as a generalizable strategy to target pathogenic senescent cell subsets across fibrosis and cancer.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Senescence-directed nanotherapy ameliorates fibrosis and overcomes immune...

IMPORTANCE / SIGNIFICANCE

Immunity And Host Response is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-18 to stabilize cell-stress signaling, epithelial integrity, and lineage-specific transcription; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target epidermal and mesendodermal lineages before mature immunity by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: cell-stress signaling, epithelial integrity, and lineage-specific transcription. Secondary readouts: survival, barrier morphology, and developmental timing, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.aeg4791](https://doi.org/10.1126/science.aeg4791)

Targeting an atypical G protein-coupled receptor signaling pathway for cardiac fibrosis therapy

Hao Zhang, Rabindra V. Shivnaraine, Lu Ren, Phung N. Thai, Remi Janicot, Wenjuan Zhu, Rihua Huang, Dirk H. Siepe, et al.

journal-article | Science 393(6818) | 2026-09-24

DOI 10.1126/science.aej5896

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	These findings establish AR-driven Gbetagamma signaling as a potential therapeutic target for cardiac fibrosis.
WHY IT MATTERS	These findings establish AR-driven Gbetagamma signaling as a potential therapeutic target for cardiac fibrosis.
MAIN / NEW METHODS	• Organoid or stem-cell model • Computational modeling
KEY EVIDENCE	• To address this unmet clinical need, we used a multidimensional drug discovery pipeline centered on a human induced pluripotent stem cell (iPSC)-based platform. • Mechanistic studies revealed an atypical adenosine receptor (AR)-dependent signaling pathway in which AR subtypes converge on Gbetagamma (the betagamma subunits of heterotrimeric GTP-binding proteins) to activate phosphoinositide 3-kinase (PI3K)-AKT and yes-associated protein (YAP). • These findings establish AR-driven Gbetagamma signaling as a potential therapeutic target for cardiac fibrosis.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Targeting an atypical G protein-coupled receptor signaling pathway for cardiac...

IMPORTANCE / SIGNIFICANCE

Cell Signaling And Fate Control is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.aej5896](https://doi.org/10.1126/science.aej5896)

Neural evidence that dogs segment the speech they hear with a humanlike consonant bias

Boglárka Morvai, Kinga G. Tóth, Marianna Boros, Dorottya S. Rácz, Ivaylo lotchev, Kitti Szabó, Attila Andics

journal-article | Science 393(6818) | 2026-09-24

DOI [10.1126/science.adw7709](https://doi.org/10.1126/science.adw7709)

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Linguistic regularity-based processing biases can thus also emerge in a nonspeaking species exposed to speech.
WHY IT MATTERS	This suggests that dogs, similar to humans, exhibit consonant bias when segmenting the speech they hear.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• Intertrial coherence measures of neural entrainment and event-related potentials revealed enhanced word-level tracking for consonant- compared with vowel-structured streams in both species.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Neural evidence that dogs segment the speech they hear with a humanlike...

IMPORTANCE / SIGNIFICANCE

Neural Development And Function is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 10-18 to stabilize neural-domain geometry, lineage allocation, and reporter dynamics; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target neural plate and neural-border lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 10-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: neural-domain geometry, lineage allocation, and reporter dynamics. Secondary readouts: marker expression, cell trajectories, and neural-tube morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.adw7709](https://doi.org/10.1126/science.adw7709)

An in-body networking system for communication between wearable and implantable therapeutics

Ramy Ghanim, Yoon Jae Lee, Garan Byun, Joy Jackson, Julia Z. Ding, Jihoon Park, Markella Bibidakis, Elaine Feller, et al.

journal-article | Science 393(6818) | 2026-09-24

DOI [10.1126/science.adz5300](https://doi.org/10.1126/science.adz5300)

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	In vivo in rats, we demonstrate coordinated, full-body networks of sensors and neural interfaces that enable wireless dual-limb motor control.
WHY IT MATTERS	Implants are syringe-injectable, require negligible power consumption in listening states, and provide >10x greater tissue communication coverage than Bluetooth.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• In vivo in rats, we demonstrate coordinated, full-body networks of sensors and neural interfaces that enable wireless dual-limb motor control.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of An in-body networking system for communication between wearable and implantable...

IMPORTANCE / SIGNIFICANCE

Cell Mechanics And Morphogenesis is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 10-13 to stabilize cell polarity, convergence-extension, tissue strain, and migration persistence; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target dorsal mesoderm and involuting marginal-zone explants by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 10-13. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: cell polarity, convergence-extension, tissue strain, and migration persistence. Secondary readouts: fate markers, adhesion dynamics, and whole-embryo axis elongation, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.adz5300](https://doi.org/10.1126/science.adz5300)

Ancient introgression drives wild boar expansion and phenotypic diversification of domestic pigs

Jian-Hai Chen, Xiaoyong Du, Zhuqing Zheng, Neena Amatya Gorkhali, Changyi Xiao, Shilin Zhu, Jie Zhong, Liangliang Fu, et al.

journal-article | Science 393(6818) | 2026-09-24

DOI 10.1126/science.adq7553

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Here, using 745 genomes representing populations across Eurasia, we demonstrate that wild boar expanded from South Asia during the Middle Pleistocene.
WHY IT MATTERS	Our findings reveal a long-lasting evolutionary legacy of this ancient introgression, which provided genetic material that was later shaped by natural and artificial selection.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• Here, using 745 genomes representing populations across Eurasia, we demonstrate that wild boar expanded from South Asia during the Middle Pleistocene. • Our findings reveal a long-lasting evolutionary legacy of this ancient introgression, which provided genetic material that was later shaped by natural and artificial selection.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Ancient introgression drives wild boar expansion and phenotypic diversification...

IMPORTANCE / SIGNIFICANCE

Gene Regulation And Chromatin is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.adq7553](https://doi.org/10.1126/science.adq7553)

Alternative immunity

Mitch Leslie

journal-article | Science 393(6818) | 2026-09-24

DOI [10.1126/science.aem6505](https://doi.org/10.1126/science.aem6505)

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	A little-known arm of the immune system respond to lipids, with a protein called CD1 driving reactions to bee stings, bacteria, and possibly poison ivy
WHY IT MATTERS	The study advances mechanistic understanding of Alternative immunity and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• The retrieved publisher-deposited metadata identifies this item as a journal article in the detected volume and issue. • The DOI provides the canonical path for manual verification of figures and full methods.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Alternative immunity

IMPORTANCE / SIGNIFICANCE

Immunity And Host Response is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-18 to stabilize cell-stress signaling, epithelial integrity, and lineage-specific transcription; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target epidermal and mesendodermal lineages before mature immunity by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: cell-stress signaling, epithelial integrity, and lineage-specific transcription. Secondary readouts: survival, barrier morphology, and developmental timing, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.
- High-dose perturbation may cause nonspecific toxicity or developmental delay.

- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.aem6505](#)

Nanotherapy against senescence

Mariantonietta D’Ambrosio, Jesús Gil

journal-article | Science 393(6818) | 2026-09-24

DOI 10.1126/science.ael2714

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Targeting senescence decreases fibrosis and boosts the response of cancer cells to immunotherapy
WHY IT MATTERS	The study advances mechanistic understanding of Nanotherapy against senescence and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• The retrieved publisher-deposited metadata identifies this item as a journal article in the detected volume and issue. • The DOI provides the canonical path for manual verification of figures and full methods.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Nanotherapy against senescence

IMPORTANCE / SIGNIFICANCE

Developmental Conservation And Mechanism is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early *Xenopus* provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use *Xenopus tropicalis* for F0 genetics and *Xenopus laevis* explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test *Xenopus laevis* and *X. tropicalis* orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct *Xenopus* evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine <i>Xenopus</i> capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points
--	--	---

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: [Featured paper \(publisher-linked source\) 10.1126/science.ael2714](https://doi.org/10.1126/science.ael2714)

CROSS-PAPER SYNTHESIS

RECURRING MECHANISMS

- immunity and host response (5 papers)
- metabolism and organelle function (2 papers)
- neural development and function (2 papers)
- cell signaling and fate control (1 paper)
- cell mechanics and morphogenesis (1 paper)

METHODOLOGICAL TRENDS

- Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details (10)
- Computational modeling (3)
- Organoid or stem-cell model (1)

RANKED XENOPUS OPPORTUNITIES

#1 Developmental timing and tissue autonomy of Escape of the *BC200* gene to a human poxvirus reveals its persistent...

Significance: high if the focal mechanism is conserved | Novelty: medium-high | Mechanistic Clarity: medium pending candidate-specific assay refinement | Preliminary Support: featured-paper evidence; Xenopus pilots proposed | Xenopus Advantage: high for temporal, spatial, rescue, and explant experiments | Feasibility: high for the three rapid pilots | Follow On Aims: epistasis, spatial profiling, and cross-species validation

#2 Developmental timing and tissue autonomy of Mammals produce *cyclo* -octasulfur to suppress lipid peroxidation and...

Significance: high if the focal mechanism is conserved | Novelty: medium-high | Mechanistic Clarity: medium pending candidate-specific assay refinement | Preliminary Support: featured-paper evidence; Xenopus pilots proposed | Xenopus Advantage: high for temporal, spatial, rescue, and explant experiments | Feasibility: high for the three rapid pilots | Follow On Aims: epistasis, spatial profiling, and cross-species validation

#3 Developmental timing and tissue autonomy of Single wildfire smoke exposure causes lasting harm, natural experiment in...

Significance: high if the focal mechanism is conserved | Novelty: medium-high | Mechanistic Clarity: medium pending candidate-specific assay refinement | Preliminary Support: featured-paper evidence; Xenopus pilots proposed | Xenopus Advantage: high for temporal, spatial, rescue, and explant experiments | Feasibility: high for the three rapid pilots | Follow On Aims: epistasis, spatial profiling, and cross-species validation

#4 Developmental timing and tissue autonomy of Fibroblasts restrain gut inflammation by IGF1-dependent regulation of innate...

Significance: high if the focal mechanism is conserved | Novelty: medium-high | Mechanistic Clarity: medium pending candidate-specific assay refinement | Preliminary Support: featured-paper evidence; Xenopus pilots proposed | Xenopus Advantage: high for temporal, spatial, rescue, and explant experiments | Feasibility: high for the three rapid pilots | Follow On Aims: epistasis, spatial profiling, and cross-species validation

COHERENT MULTI-AIM PROPOSAL

A coherent multi-aim proposal can unite papers centered on immunity and host response: Aim 1 establishes conservation and timing, Aim 2 resolves cell autonomy and pathway order, and Aim 3 links molecular activity to tissue-scale robustness.

INDEPENDENT PILOTS

- Developmental timing and tissue autonomy of Mammals produce *cyclo*-octasulfur to suppress lipid peroxidation and...
- Developmental timing and tissue autonomy of Single wildfire smoke exposure causes lasting harm, natural experiment in...
- Developmental timing and tissue autonomy of Fibroblasts restrain gut inflammation by IGF1-dependent regulation of innate...

SOURCE / METHOD NOTE

Retrieved 2026-09-24T21:35:20.050865+00:00. Change detection and bibliographic metadata: publisher-deposited Crossref record (<https://api.crossref.org/journals/0036-8075/works>). Abstract fallback: Europe PMC when available. Canonical issue and DOI URLs point to the publishers. Official-page check: Official publisher issue page returned HTTP 403; no full-page review is claimed. The automated evidence boundary is publisher-deposited Crossref metadata plus legally available Europe PMC abstracts. Publisher pages are probed but paywalls and access controls are not bypassed. Figures and full methods are not claimed as reviewed. No OpenAI API, model API, paid literature service, paywall bypass, unpublished user data, or claimed full-text review was used. Grant concepts are speculative templates requiring expert, candidate-specific revision.