CNS MCDB UPDATE

Nature | Volume 658 | Issue 8136

Issue date: 2026-10-15 | Retrieved: 2026-10-15T04:49:03.120206+00:00 | 13 relevant primary-research papers

Evidence discipline

Paper-supported findings, published enabling evidence, cross-species inference, speculative hypotheses, and proposed pilot experiments are labeled separately. Access notes state when only an abstract or limited preview was available.

Screening notes

- Official publisher issue page was reachable and contained the detected volume and issue identifiers.
- Screened 13 biology-relevant journal articles after excluding reviews, perspectives, news, corrections, and editorials by metadata/title rules.
- All summaries are deterministic extractions from the bounded public evidence packet; no language-model API or paid research service was used.

Official issue

https://www.nature.com/nature/volumes/658/issues/8136

Why Xenopus now

Early Xenopus combines rapid external development, targeted microinjection, explants, lineage tracing, live imaging, F0 CRISPR, rescue, and scalable replication. It can convert abstract-supported candidate mechanisms into spatially resolved causal tests before committing to slower mammalian models.

Procognitive restoration of PV neuron plasticity in neurodevelopmental disorders

Yu-Tzu Shih, Jason Bondoc Alipio, Zin-Juan Klaft, Nathaniel Green, Alok Nath Mohapatra, Travis D. Goode, Muthu Panchanatham, Devesh Pathak, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10907-8

Access: Abstract retrieved from Europe PMC; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Here we designed an input-specific translatome screen to identify regulators of experience-dependent PV IN plasticity genes (XPGs) in the CA3/CA2 subregion of adult hippocampus.
WHY IT MATTERS	We demonstrate that a substantial proportion of upregulated candidate XPGs exhibit haploinsufficiency in autism spectrum disorder, epilepsies, bipolar disorder and schizophrenia, which suggests that there is impaired experience-dependent PV IN plasticity in NDDs.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• Here we designed an input-specific translatome screen to identify regulators of experience-dependent PV IN plasticity genes (XPGs) in the CA3/CA2 subregion of adult hippocampus. • We demonstrate that a substantial proportion of upregulated candidate XPGs exhibit haploinsufficiency in autism spectrum disorder, epilepsies, bipolar disorder and schizophrenia, which suggests that there is impaired experience-dependent PV IN plasticity in NDDs.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

SPECIFIC GRANT IDEA 1 | PAPER 1

Developmental timing and tissue autonomy of Procognitive restoration of PV neuron plasticity in neurodevelopmental disorders

IMPORTANCE / SIGNIFICANCE

Neural Development And Function is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 10-18 to stabilize neural-domain geometry, lineage allocation, and reporter dynamics; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target neural plate and neural-border lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 10-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: neural-domain geometry, lineage allocation, and reporter dynamics. Secondary readouts: marker expression, cell trajectories, and neural-tube morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10907-8

Temporal uncoupling of radial glia lineage progression in cortical organoids

Melissa Stouffer, Osvaldo A. Miranda, Florian M. Pauler, Fabrizia Pipicelli, Carmen Streicher, Giselle Cheung, Simon Hippenmeyer

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10916-7

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	We found that RGPs exhibit a high level of plasticity in proliferative potential in organoids rather than strict temporally stereotyped lineage progression as observed in vivo.
WHY IT MATTERS	We found that RGPs exhibit a high level of plasticity in proliferative potential in organoids rather than strict temporally stereotyped lineage progression as observed in vivo.
MAIN / NEW METHODS	• Single-cell transcriptomics • Organoid or stem-cell model
KEY EVIDENCE	• Mosaic analysis with double markers (MADM)-based lineage tracing in vivo has revealed a quantitative framework of RGP lineage progression 1 . • Here we established MADM technology 2,3 in mouse embryonic stem cells to probe RGP lineage progression in a self-organizing cortical organoid system. • We found that RGPs exhibit a high level of plasticity in proliferative potential in organoids rather than strict temporally stereotyped lineage progression as observed in vivo.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

SPECIFIC GRANT IDEA 1 | PAPER 2

Developmental timing and tissue autonomy of Temporal uncoupling of radial glia lineage progression in cortical organoids

IMPORTANCE / SIGNIFICANCE

Neural Development And Function is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 10-18 to stabilize neural-domain geometry, lineage allocation, and reporter dynamics; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target neural plate and neural-border lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 10-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: neural-domain geometry, lineage allocation, and reporter dynamics. Secondary readouts: marker expression, cell trajectories, and neural-tube morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10916-7

Biomarkers of nivolumab benefit in resectable non-small cell lung cancer

Tina Cascone, Mark M. Awad, Jonathan D. Spicer, Jie He, Shun Lu, Fumihiro Tanaka, Robin Cornelissen, Lubos B. Petruzelka, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10925-6

Access: Abstract retrieved from Europe PMC; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	These findings provide insights into predictive markers for outcomes with perioperative nivolumab in resectable NSCLC.
WHY IT MATTERS	These findings provide insights into predictive markers for outcomes with perioperative nivolumab in resectable NSCLC.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• These findings provide insights into predictive markers for outcomes with perioperative nivolumab in resectable NSCLC.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

SPECIFIC GRANT IDEA 1 | PAPER 3

Developmental timing and tissue autonomy of Biomarkers of nivolumab benefit in resectable non-small cell lung cancer

IMPORTANCE / SIGNIFICANCE

Gene Regulation And Chromatin is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10925-6

Family genetic designs in MoBa provide insights into health and functioning

Elizabeth C. Corfield, Alexey A. Shadrin, Oleksandr Frei, Zillur Rahman, Bayram Cevdet Akdeniz, Tahir Tekin Filiz, Aihua Lin, Isabella Badini, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10926-5

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	We then use trio data to illustrate how family-based genomic designs can identify distinct direct and indirect sources of genetic influence and structural confounding.
WHY IT MATTERS	These demonstrations highlight MoBa as a broadly valuable resource for advancing understanding of health and functioning across the lifecourse and generations.
MAIN / NEW METHODS	• Genome-wide association or QTL mapping
KEY EVIDENCE	• Abstract Genome-wide association studies using large, population-based samples of unrelated individuals have discovered thousands of genetic associations with health and disease 1 . • However, increasing evidence suggests that population-based estimates, while precise, can also reflect confounding that affects their use and interpretation. • This confounding can be overcome using data from genotyped family members, such as nuclear mother-father-child trios 2,3 .
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

SPECIFIC GRANT IDEA 1 | PAPER 4

Developmental timing and tissue autonomy of Family genetic designs in MoBa provide insights into health and functioning

IMPORTANCE / SIGNIFICANCE

Gene Regulation And Chromatin is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10926-5

Creating bottom-up RNA transfer vehicles from synthetic protein assemblies

Maren Kirstin Schuhmacher, Christoph Gruber, Christopher M. R. Lang, Ricardo M. W. Ruijpers, Lyupka Mazneykova, Brice Beinsteiner, Ariane Krus, Barbara Tremmel, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10952-3

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	In addition, we demonstrate that their tropism can be programmed by incorporation of computationally designed peptide binders and use them to deliver therapeutically relevant cargo RNAs into a wide range of cellular models.
WHY IT MATTERS	The study advances mechanistic understanding of Creating bottom-up RNA transfer vehicles from synthetic protein assemblies and creates a defined basis for testing conservation and developmental function in a vertebrate embryo.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• In addition, we demonstrate that their tropism can be programmed by incorporation of computationally designed peptide binders and use them to deliver therapeutically relevant cargo RNAs into a wide range of cellular models. • We show the in vivo biodistribution of one of these vehicles in a mouse at near-single-cell resolution, confirm its safety, and use it to perform a gene-editing treatment strategy for Duchenne muscular dystrophy in patient-derived cells and a pig. • Our work demonstrates how proteins created by generative artificial intelligence can be harnessed for the rational engineering of RNA transport systems with the desired properties by overcoming the limitations of natural protein diversity.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

SPECIFIC GRANT IDEA 1 | PAPER 5

Developmental timing and tissue autonomy of Creating bottom-up RNA transfer vehicles from synthetic protein assemblies

IMPORTANCE / SIGNIFICANCE

Gene Regulation And Chromatin is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10952-3

Rewiring the ribosome to translate proteins encoded in its own RNA

Kasra Alizadeh, Dorota Klepacki, Nora Vázquez-Laslop, Alexander S. Mankin

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10962-1

Access: Abstract retrieved from Europe PMC; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Here we show that bacterial ribosomes can synthesize proteins encoded within their own RNA.
WHY IT MATTERS	One potential solution is integrating the protein-coding sequence into the ribosome itself, thereby committing the ribosome to synthesis of a single polypeptide.
MAIN / NEW METHODS	• Biochemical reconstitution
KEY EVIDENCE	• Here we show that bacterial ribosomes can synthesize proteins encoded within their own RNA. • These findings establish the mechanistic feasibility of a ribosome translating its own rRNA in vivo and in vitro, offering a versatile platform for orthogonal protein production and insights into the origin of translation.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Rewiring the ribosome to translate proteins encoded in its own RNA

IMPORTANCE / SIGNIFICANCE

Gene Regulation And Chromatin is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10962-1

Aberrant excitatory neuronal ERBB4 promotes Alzheimer’s disease pathology

Se Young Lee, Eunseok Park, Ha-Eun Lee, Seongbin Kim, Yeji Yeo, Juwon Park, Young-Jin Choi, Kiheon Lee, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10964-z

Access: Abstract retrieved from Europe PMC; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Here we demonstrate that, during AD progression in mice, astrocytes and microglia increase phagocytic elimination of excitatory synapses while reducing elimination of inhibitory synapses, suggesting that neuroinflammation alone may be dispensable for early AD synapse loss.
WHY IT MATTERS	These findings identify aberrant Erbb4 expression in excitatory neurons as an early driver of AD pathophysiology and a potential therapeutic target across neurodegenerative diseases.
MAIN / NEW METHODS	• Single-nucleus multi-omics
KEY EVIDENCE	• Here we demonstrate that, during AD progression in mice, astrocytes and microglia increase phagocytic elimination of excitatory synapses while reducing elimination of inhibitory synapses, suggesting that neuroinflammation alone may be dispensable for early AD synapse loss. • Subsequent transcriptomic analyses showed that excitatory neuronal Erbb4 is both necessary and sufficient to induce EREN and reactive gliosis. • These findings identify aberrant Erbb4 expression in excitatory neurons as an early driver of AD pathophysiology and a potential therapeutic target across neurodegenerative diseases.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

SPECIFIC GRANT IDEA 1 | PAPER 7

Developmental timing and tissue autonomy of Aberrant excitatory neuronal ERBB4 promotes Alzheimer’s disease pathology

IMPORTANCE / SIGNIFICANCE

Neural Development And Function is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 10-18 to stabilize neural-domain geometry, lineage allocation, and reporter dynamics; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target neural plate and neural-border lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 10-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: neural-domain geometry, lineage allocation, and reporter dynamics. Secondary readouts: marker expression, cell trajectories, and neural-tube morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10964-z

A binding-to-release strategy for targeted anticancer drug delivery

Zihao Wen, Mengxin Xu, Zijun Yan, Ziren Kong, Yupeng Wang, Pei Liu, Junyi Chen, Yanzhao Liu, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10971-0

Access: Abstract retrieved from Europe PMC; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	This work establishes a framework that overcomes the internalization barrier, broadening the target scope for therapeutic and diagnostic conjugates.
WHY IT MATTERS	This work establishes a framework that overcomes the internalization barrier, broadening the target scope for therapeutic and diagnostic conjugates.
MAIN / NEW METHODS	• Biochemical reconstitution
KEY EVIDENCE	• Drug conjugates, such as antibody-drug conjugates (ADCs) and small molecule-drug conjugates (SMDCs), are often dependent on efficient receptor-mediated endocytosis for payload release 1-3 -supported by about 10% of targets 4-7 . • This platform demonstrated high specificity from in vitro to clinical specimens, achieving precise detection of fibroblast activation protein (FAP) expression in patient-derived lymph nodes. • This work establishes a framework that overcomes the internalization barrier, broadening the target scope for therapeutic and diagnostic conjugates.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of A binding-to-release strategy for targeted anticancer drug delivery

IMPORTANCE / SIGNIFICANCE

Gene Regulation And Chromatin is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10971-0

Synergistic degradation of fucoidans in the ocean

Andreas Sichert, Shaul Pollak, Taylor Priest, Akshit Goyal, Samuel Miravet-Verde, Shinichi Sunagawa, Otto X. Cordero, Uwe Sauer

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10980-z

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Here we show that fucoidans are degraded through synergistic interactions between specialized bacteria with complementary metabolic functions.
WHY IT MATTERS	This work provides a mechanistic framework for understanding carbon cycling in the ocean and for engineering synthetic microbial consortia to degrade recalcitrant polysaccharides.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• Here we show that fucoidans are degraded through synergistic interactions between specialized bacteria with complementary metabolic functions. • Our findings suggest that the environmental turnover of complex biopolymers depends not only on individual metabolic capabilities of degraders but also on ecological interactions shaped by substrate architecture.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

SPECIFIC GRANT IDEA 1 | PAPER 9

Developmental timing and tissue autonomy of Synergistic degradation of fucoidans in the ocean

IMPORTANCE / SIGNIFICANCE

Metabolism And Organelle Function is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-13 to stabilize organelle state, metabolite-sensitive reporters, and lineage-marker induction; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target animal-cap and marginal-zone explants by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-13. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: organelle state, metabolite-sensitive reporters, and lineage-marker induction. Secondary readouts: survival, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10980-z

Intracellular complement factor H protects neurons during CNS inflammation

Christina Mayer, Marcel S. Woo, Jana K. Sonner, Lars Binkle-Ladisch, Felix Fischbach, Patricia Sekol, Matthew D. Smith, Fernando Lucas-Ruiz, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10981-y

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Here we identify a neuroprotective role for intracellular complement factor H (CFH) in neurons.
WHY IT MATTERS	Together, our results identify CFH as a key mediator of neuronal resilience across the CNS in mice and humans and provide mechanistic insight into inflammatory neurodegeneration with implications for MS therapy and neuroprotection more broadly.
MAIN / NEW METHODS	• Single-nucleus multi-omics
KEY EVIDENCE	• Here we identify a neuroprotective role for intracellular complement factor H (CFH) in neurons. • Using single-nucleus RNA-sequencing analysis of RGCs from donors with MS and control individuals, we found that CFH expression was strongly correlated with intrinsic resilience to RGC degeneration. • These findings reveal a non-canonical intracellular function of CFH in neurons.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Intracellular complement factor H protects neurons during CNS inflammation

IMPORTANCE / SIGNIFICANCE

Neural Development And Function is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 10-18 to stabilize neural-domain geometry, lineage allocation, and reporter dynamics; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target neural plate and neural-border lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 10-18. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: neural-domain geometry, lineage allocation, and reporter dynamics. Secondary readouts: marker expression, cell trajectories, and neural-tube morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10981-y

3D epigenome of glial cell types in developing human cortex

Ian R. Jones, Li Wang, Michael Kosicki, Stephanie L. Battle, Vivek JJ Narayan, Qiuli Bi, Kaila Gemenes, Yuxi Liu, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-10987-6

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	Finally, oRG cCREs are enriched for human accelerated regions compared with other cCREs and a subset of human accelerated regions show activity differences from their chimpanzee orthologues that interact with genes involved in neuronal development.
WHY IT MATTERS	Our findings advance the understanding of human-specific gene regulation during corticogenesis.
MAIN / NEW METHODS	• Chromatin profiling • Computational modeling
KEY EVIDENCE	• Our prior study of neurogenesis, including radial glia (RG), intermediate progenitor cells, excitatory neurons and interneurons demonstrated that chromatin looping underlies transcriptional regulation for lineage-specific genes, shedding light on how non-coding genetic variants contribute to neuropsychiatric disorders by means of cell-type-specific gene regulation 3 . • Finally, oRG cCREs are enriched for human accelerated regions compared with other cCREs and a subset of human accelerated regions show activity differences from their chimpanzee orthologues that interact with genes involved in neuronal development. • Our findings advance the understanding of human-specific gene regulation during corticogenesis.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of 3D epigenome of glial cell types in developing human cortex

IMPORTANCE / SIGNIFICANCE

Gene Regulation And Chromatin is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-10987-6

Scalable near-real-time Bayesian phylogenetics for outbreaks with Delphy

Patrick Varilly, Mark Schifferli, Katherine Yang, Paul Cronan, Ivan Specht, Tim Burcham, Olivia Glennon, Olivia Jacks, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-11012-6

Access: Abstract retrieved from Europe PMC; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	23,24), we demonstrate state-of-the-art accuracy with up to 2-3 orders of magnitude improvements in speed.
WHY IT MATTERS	Pathogen genomic analysis is central to tracking, understanding and containing outbreaks 1-13 , but the complexity and cost of state-of-the-art phylogenetic tools limit global access and impact.
MAIN / NEW METHODS	• Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details
KEY EVIDENCE	• 23,24), we demonstrate state-of-the-art accuracy with up to 2-3 orders of magnitude improvements in speed. • Assessing Delphy's scalability, we show that a simulated dataset of 100,000 sequences can be analysed within a day. • Delphy automatically identifies key viral lineages and mutations, as well as their emergence and prevalence through time, with quantified uncertainties grounded in Bayesian theory.
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of Scalable near-real-time Bayesian phylogenetics for outbreaks with Delphy

IMPORTANCE / SIGNIFICANCE

Gene Regulation And Chromatin is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-11012-6

The gene-regulatory evolution of the human skeleton

Yizhi Yan, Nadav Mishol, Katharina Lange, Zicong Zhang, Gal Bodek, Aya Kigel, Noam Priel, Nachshon Egyes, et al.

journal-article | Nature 658(8136) | 2026-10-15

DOI 10.1038/s41586-026-11053-x

Access: Publisher-deposited abstract retrieved through Crossref; no full-text review is claimed. | Evidence status: abstract-supported

MAIN DISCOVERY	These atlases reveal an extensive rewiring of the extracellular matrix (ECM), including a marked suppression of glycosaminoglycan (GAG) biosynthesis, leading to an approximately three-to-fourfold reduction in joint GAG content in humans compared with non-human apes.
WHY IT MATTERS	Together, our results reveal a coordinated evolutionary remodelling of the human skeletal ECM, and establish a comprehensive framework for dissecting the genetic basis of human skeletal biology.
MAIN / NEW METHODS	• Computational modeling
KEY EVIDENCE	• Using massively parallel reporter assays (MPRAs) in chondrocytes, we assayed 561,410 human-derived substitutions in promoters and enhancers, identifying 15,077 loci with human-specific regulatory activity. • These atlases reveal an extensive rewiring of the extracellular matrix (ECM), including a marked suppression of glycosaminoglycan (GAG) biosynthesis, leading to an approximately three-to-fourfold reduction in joint GAG content in humans compared with non-human apes. • We find that this human-specific shift bears signatures of selection, and is likely to be a key contributor to the exceptional susceptibility of humans to degenerative skeletal diseases 3-5 .
LIMITATIONS	• This automated report used title, DOI metadata, and an available abstract; it does not claim manual full-text, figure, or complete methods review. • The deterministic extractor cannot assess experimental quality, effect size, statistics, or unreported negative results beyond the retrieved source text. • The proposed Xenopus mechanism is new speculation and has not been established by the featured paper.

Developmental timing and tissue autonomy of The gene-regulatory evolution of the human skeleton

IMPORTANCE / SIGNIFICANCE

Gene Regulation And Chromatin is central to how embryonic cells convert molecular state into reproducible tissue organization. Establishing whether the featured mechanism operates during early vertebrate development could reveal a conserved control point and a tractable route to explain developmental variability without overstating clinical translation.

KNOWLEDGE GAP

The featured paper does not establish whether the focal regulator or mechanism described by the featured paper acts during early vertebrate development, whether its effect is cell-autonomous, which developmental window is sensitive, or whether molecular and morphogenetic phenotypes can be rescued independently.

RATIONALE AND XENOPUS ADVANTAGE

The paper supplies a candidate mechanism, while early Xenopus provides rapid external development, targeted blastomere injection, explants, lineage tracing, live imaging, inexpensive biological replication across clutches, and direct comparison of molecular and tissue-scale outcomes.

CENTRAL HYPOTHESIS

Speculative: the focal regulator or mechanism described by the featured paper acts cell-autonomously during stages 8-14 to stabilize pathway activity, lineage allocation, and spatial marker-domain size; acute loss will reduce the predicted molecular readout and increase pattern variance, while stage-matched wild-type rescue will restore both.

EXPERIMENTAL DESIGN

Use Xenopus tropicalis for F0 genetics and Xenopus laevis explants. Target animal-cap, organizer, and mesendodermal lineages by injecting two independent CRISPR RNPs at the 4- to 8-cell stage with a fluorescent lineage tracer; analyze stages 8-14. Groups: uninjected, non-targeting RNP, each guide alone, pooled guides, dose series, wild-type mRNA rescue, perturbation-resistant rescue, and temporally delayed perturbation where a validated small molecule or inducible construct exists. Primary readouts: pathway activity, lineage allocation, and spatial marker-domain size. Secondary readouts: cell behavior, developmental timing, and gastrulation morphology, amplicon sequencing, RT-qPCR or in situ hybridization, and an orthogonal biochemical or reporter assay. Use at least three independent spawnings, 20-30 embryos per group per spawning for morphology, and 8-10 pooled or individually profiled samples per molecular condition; randomize embryos, blind image scoring, and analyze clutch as a random effect with multiplicity-controlled planned contrasts. Decision point: advance to epistasis and spatial single-cell profiling only if both guides reproduce the effect and rescue restores at least 50% of the molecular and developmental phenotype.

EXPECTED RESULTS AND INTERPRETATIONS

Support would be a guide-concordant loss of the predicted molecular readout, increased spatial or embryo-to-embryo variance, and rescue of both molecular and tissue phenotypes. This would place the mechanism upstream of developmental robustness. A molecular phenotype without morphogenesis would suggest buffering or a parallel tissue effector and would redirect the next aim to sensitized or explant conditions. A morphogenetic phenotype without the predicted reporter change would argue for an alternative biochemical function. No reproducible phenotype after confirmed editing would refute an obligate early role and shift the model toward redundancy, later function, or species-specific deployment.

POTENTIAL PITFALLS

- F0 mosaicism may weaken or spatially blur phenotypes.
- Paralog redundancy or maternal stores may mask an early requirement.

- High-dose perturbation may cause nonspecific toxicity or developmental delay.
- The featured mechanism may not be conserved or may use different effectors in amphibians.
- Abstract-only evidence may omit critical context needed to choose the most specific assay.

ALTERNATIVE STRATEGIES

- Map editing per embryo and enrich analysis for high-editing specimens; establish stable alleles if mosaicism remains limiting.
- Target expressed paralogs jointly, use maternal-zygotic strategies where feasible, or test dominant-negative and degron-based acute perturbations.
- Use lower doses, later induction, tissue-restricted injection, explants, and matched survival/staging controls to separate mechanism from toxicity.
- Test Xenopus laevis and X. tropicalis orthologs, then validate the key interaction biochemically or in a complementary cell model.
- Before full-scale work, manually review the linked paper's figures and methods and revise the reporter or perturbation to match the demonstrated mechanism.

PUBLISHED ENABLING EVIDENCE AND PROPOSED PRELIMINARY-DATA PLAN

Published enabling evidence and proposed preliminary-data plan: no unpublished user-laboratory data are assumed. The featured paper is the cited cross-species enabling source; direct Xenopus evidence was not automatically verified. Pilot 1: map candidate expression or pathway activity at 5-6 early stages using RT-qPCR and in situ hybridization (three spawnings; expected Figure 1A-B); go if a reproducible spatial or temporal signal exceeds twofold over the lowest stage. Pilot 2: test two CRISPR RNPs plus non-targeting control in three spawnings with at least 25 embryos per group, amplicon sequencing, blinded morphology, and the primary molecular readout (Figure 1C-E); go if both guides produce the same directional effect of at least 20% with acceptable survival. Pilot 3: perform wild-type mRNA rescue in three spawnings with at least 20 embryos per group (Figure 1F-G); go if rescue reverses at least 50% of the molecular and phenotype effect. Failure at any threshold triggers the paired alternative strategy rather than a claim of preliminary support.

NOVELTY medium-high - transfers a recent mechanism into a spatially and temporally tractable vertebrate embryo	FEASIBILITY high for the pilots - injection, explants, imaging, genotyping, and rescue are routine Xenopus capabilities	RISK medium - conservation, maternal contribution, and mosaicism are explicit decision points

GRANT FIT

Exploratory Aim 1 for conservation and causality; successful pilots support Aim 2 epistasis and Aim 3 spatial mechanism.

STRONGEST REVIEWER-FACING PREMISE

Xenopus can test when, where, and whether a newly reported cellular mechanism controls intact vertebrate development in one experimentally integrated system.

SUPPORTING SOURCES

featured paper: Featured paper (publisher-linked source) 10.1038/s41586-026-11053-x

CROSS-PAPER SYNTHESIS

RECURRING MECHANISMS

- gene regulation and chromatin (8 papers)
- neural development and function (4 papers)
- metabolism and organelle function (1 paper)

METHODOLOGICAL TRENDS

- Publisher-deposited title/abstract analysis; consult the linked paper for complete technical details (5)
- Biochemical reconstitution (2)
- Single-nucleus multi-omics (2)
- Computational modeling (2)
- Single-cell transcriptomics (1)
- Organoid or stem-cell model (1)

RANKED XENOPUS OPPORTUNITIES

#1 Developmental timing and tissue autonomy of Procognitive restoration of PV neuron plasticity in neurodevelopmental disorders

Significance: high if the focal mechanism is conserved | Novelty: medium-high | Mechanistic Clarity: medium pending candidate-specific assay refinement | Preliminary Support: featured-paper evidence; Xenopus pilots proposed | Xenopus Advantage: high for temporal, spatial, rescue, and explant experiments | Feasibility: high for the three rapid pilots | Follow On Aims: epistasis, spatial profiling, and cross-species validation

#2 Developmental timing and tissue autonomy of Temporal uncoupling of radial glia lineage progression in cortical organoids

Significance: high if the focal mechanism is conserved | Novelty: medium-high | Mechanistic Clarity: medium pending candidate-specific assay refinement | Preliminary Support: featured-paper evidence; Xenopus pilots proposed | Xenopus Advantage: high for temporal, spatial, rescue, and explant experiments | Feasibility: high for the three rapid pilots | Follow On Aims: epistasis, spatial profiling, and cross-species validation

#3 Developmental timing and tissue autonomy of Biomarkers of nivolumab benefit in resectable non-small cell lung cancer

Significance: high if the focal mechanism is conserved | Novelty: medium-high | Mechanistic Clarity: medium pending candidate-specific assay refinement | Preliminary Support: featured-paper evidence; Xenopus pilots proposed | Xenopus Advantage: high for temporal, spatial, rescue, and explant experiments | Feasibility: high for the three rapid pilots | Follow On Aims: epistasis, spatial profiling, and cross-species validation

#4 Developmental timing and tissue autonomy of Family genetic designs in MoBa provide insights into health and functioning

Significance: high if the focal mechanism is conserved | Novelty: medium-high | Mechanistic Clarity: medium pending candidate-specific assay refinement | Preliminary Support: featured-paper evidence; Xenopus pilots proposed | Xenopus Advantage: high for temporal, spatial, rescue, and explant experiments | Feasibility: high for the three rapid pilots | Follow On Aims: epistasis, spatial profiling, and cross-species validation

COHERENT MULTI-AIM PROPOSAL

A coherent multi-aim proposal can unite papers centered on gene regulation and chromatin: Aim 1 establishes conservation and timing, Aim 2 resolves cell autonomy and pathway order, and Aim 3 links molecular activity to tissue-scale robustness.

INDEPENDENT PILOTS

- Developmental timing and tissue autonomy of Temporal uncoupling of radial glia lineage progression in cortical organoids
- Developmental timing and tissue autonomy of Biomarkers of nivolumab benefit in resectable non-small cell lung cancer
- Developmental timing and tissue autonomy of Family genetic designs in MoBa provide insights into health and functioning

SOURCE / METHOD NOTE

Retrieved 2026-10-15T04:49:03.120206+00:00. Change detection and bibliographic metadata: publisher-deposited Crossref record (https://api.crossref.org/journals/0028-0836/works). Abstract fallback: Europe PMC when available. Canonical issue and DOI URLs point to the publishers. Official-page check: Official publisher issue page was reachable and contained the detected volume and issue identifiers. The automated evidence boundary is publisher-deposited Crossref metadata plus legally available Europe PMC abstracts. Publisher pages are probed but paywalls and access controls are not bypassed. Figures and full methods are not claimed as reviewed. No OpenAI API, model API, paid literature service, paywall bypass, unpublished user data, or claimed full-text review was used. Grant concepts are speculative templates requiring expert, candidate-specific revision.